



**The external PASTA domain of the essential
serine/threonine protein kinase PknB regulates
mycobacterial growth**

Journal:	<i>Open Biology</i>
Manuscript ID:	RSOB-15-0025.R2
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Turapov, Obolbek; University of Leicester, Loraine, Jessica; University of Leicester, Jenkins, Christopher; University of Leicester, Barthe, Philippe; CNRS, McFeely, Daniel; University of Leicester, Forti, Francesca; Università degli Studi di Milano, Ghisotti, Daniella; Università degli Studi di Milano, Hesek, Dusan; University of Notre Dame, Lee, Mijoon; University of Notre Dame, Bottrill, Andrew; University of Leicester, Vollmer, Waldemar; Newcastle University, Mobashery, Shahriar; University of Notre Dame, Cohen-Gonsaud, Martin; CNRS, Mukamolova, Galina; University of Leicester,
Subject:	microbiology, biochemistry
Keywords:	Mycobacterium tuberculosis, serine/threonine protein kinase, magnesium, muropeptides, PknB, PASTA domain

SCHOLARONE™
Manuscripts

The external PASTA domain of the essential serine/threonine protein kinase PknB regulates mycobacterial growth

Obolbek Turapov¹, Jessica Loraine^{1§}, Christopher H. Jenkins¹, Philippe Barthe^{2,3}, Daniel McFeely^{1&}, Francesca Forti⁴, Daniela Ghisotti⁴, Dusan Hese⁵, Mijoon Lee⁵, Andrew R. Bottrill¹, Waldemar Vollmer⁶, Shahriar Mobashery⁵, Martin Cohen-Gonsaud^{2,3} and Galina V. Mukamolova^{1*}

¹Department of Infection, Immunity and Inflammation, University of Leicester, Leicester, LE1 9HN, UK; ²Centre de Biochimie Structurale, CNRS UMR 5048, 29, rue de Navacelles 34090 Montpellier, France; ³INSERM U1054, Université Montpellier I et II, Montpellier, France; ⁴Dipartimento di BioScienze, Università degli Studi di Milano, Italy; ⁵Department of Chemistry and Biochemistry, 423 Nieuwland Science Center, University of Notre Dame, Notre Dame, IN 46556 USA; ⁶ Centre for Bacterial Cell Biology, Institute for Cell and Molecular Biosciences, Newcastle University, Newcastle upon Tyne, NE2 4AX, United Kingdom.

[§] Current address: Department of Biology, University of York, Wentworth way, York, YO10 5DD

[&]Current address: School of Life Sciences, The University of Warwick, Coventry, CV4 7AL

*To whom correspondence should be addressed. E-mail: gvm4@le.ac.uk

Abstract

PknB is an essential serine/threonine protein kinase required for mycobacterial cell division and cell wall biosynthesis. Here we demonstrate that over-expression of the external PknB_PASTA domain in mycobacteria results in delayed regrowth, accumulation of elongated bacteria and increased sensitivity to β -lactam antibiotics. These changes are accompanied by altered production of certain enzymes involved in cell wall biosynthesis as revealed by proteomics studies. The growth inhibition caused by over-expression of the PknB_PASTA domain is completely abolished by enhanced concentration of magnesium ions but not muropeptides. Finally, we show that the addition of recombinant PASTA domain could prevent regrowth of *Mycobacterium tuberculosis* and therefore offers an alternative opportunity to control replication of this pathogen. These results suggest that the PknB_PASTA domain is involved in regulation of peptidoglycan biosynthesis and maintenance of cell wall architecture.

Keywords: *Mycobacterium tuberculosis*, serine/threonine protein kinase, magnesium, muropeptides, PknB, PASTA domain.

Introduction

Serine/Threonine protein kinases (STPKs) are widely distributed in Gram-positive and Gram-negative bacteria [1]. *Mycobacterium tuberculosis*, the causative agent of tuberculosis, possesses 11 STPKs [2] and two of them, PknA and PknB, are indispensable for growth in laboratory culture [3-5], while PknE [6], PknG [7, 8] and PknH [9] have been implicated in *M. tuberculosis* virulence. The

essential PknB kinase belongs to a distinct family of STPKs found only in Gram-positive bacteria [10]. The important feature of these kinases is the presence of the so-called PASTA (Penicillin-binding protein and serine/threonine kinase associated) domains in the surface-exposed region [11]. In Firmicutes, PASTA domain containing kinases are not essential for growth. In *Staphylococcus aureus*, an Stk1 mutant was impaired in virulence and had higher resistance to Triton X-100 and fosmidomycin [12], while in *S. pneumoniae* StkP kinase was important for competence, biofilm formation and virulence [13]. More detailed investigation of the role of StkP in *S. pneumoniae* established that it has a crucial role in coordinating cell division and peptidoglycan synthesis during growth [14]. In contrast, PrkC in *Bacillus subtilis* was shown to be important for survival in stationary phase. A *prkC* deletion mutant had a significantly lower optical density in stationary phase compared with the wild-type bacilli [15]. However, in later studies it was demonstrated that PrkC regulated a novel muropeptide-mediated germination pathway [16] and possibly remodeling the cell wall via controlling expression of YocH muralytic enzyme [17].

In *Streptomyces coelicolor* a PASTA-domain containing kinase PknB regulates carbon flux and antibiotic production and is not essential for growth [18]. In *Corynebacterium glutamicum* PknB is also dispensable for growth and a significant change in replication and cell shape could only be detected in mutants missing several STPKs [19]. In this bacterium PknB apparently regulates polymerization of FtsZ, however the precise mechanism and biological significance of this observation require further investigation.

Mycobacteria appear to be a unique bacterial group in which PknB is essential for growth [3-5]. Its overexpression or partial depletion in *M. smegmatis* and *Mycobacterium bovis* BCG caused dramatic alterations of cellular morphology and growth inhibition; the enzymatic activity of PknB was demonstrated to be essential for the observed effects [4, 5]. Over the past decade great progress has been achieved in the identification of PknB substrates [10]. They include proteins from various functional categories; cell-wall enzymes - InhA [20], PbpA [21] and MabA [22]; regulatory proteins - SigH [23], GarA [24] and FhaA [25]; proteins involved in cell division - Wag31 [4] and MviN [26]. Also, PknB regulates an “oxygen-mediated replication switch” and hence transition to dormancy and resuscitation [27]. Therefore it is not surprising that PknB has been considered as a critical drug target for development of anti-tuberculosis antimicrobials. Possible application of PknB kinase inhibitors for killing of *M. tuberculosis* was investigated. Yet only limited success has been achieved, mainly due to poor penetration of these agents into mycobacterial cells [28].

PknB consists of several domains (figure 1); functions for some of them have been established, while the precise role of others remain unknown [29]. The components of PknB include a conserved catalytic kinase domain, a juxtamembrane part attached to a membrane-spanning region and surface-exposed sensory component, consisting of PASTA, designated as PknB_PASTA domain [30, 31]. PASTA domains have been proposed to recognize growing strands of nascent peptidoglycan and activate serine/threonine protein kinases. Structural studies showed that *B. subtilis* PASTA domain binds synthetic

muropeptides at relatively high affinity, with the presence of the diaminopimelic acid in the muropeptide being crucial for this binding [32]. The PknB_PASTA domain from mycobacteria can also bind, synthetic muropeptides, however it remains unclear whether this binding influences activation of PknB, bacterial growth and resuscitation [33].

In this study we investigated whether the extracellular PknB_PASTA domain itself plays a distinct role in mycobacterial growth and therefore can be potentially used as a drug target or a novel chemotherapeutic agent. The results presented here suggest that the PASTA domain likely recognizes growing peptidoglycan strands, regulates production of penicillin-binding proteins and distribution of proteins involved in division and generation of septum. Furthermore, our data indicate that the interactions of the PknB_PASTA domain with the cell wall are important and that its abundance may serve as a controlling mechanism for integrity of cell wall and bacterial growth.

Materials and Methods

Bacterial strains and growth media. *Mycobacterium smegmatis* mc2155, *Mycobacterium bovis* BCG Glaxo strain and *Mycobacterium tuberculosis* H37Rv were grown in Sauton's medium or Middlebrook 7H9 (Becton, Dickinson and Company) medium supplemented with Albumin-Dextrose Complex (designated as supplemented 7H9 medium). *M. smegmatis* was also grown in Lysogeny broth (LB) with addition of 0.05% (w/v) Tween 80 to prevent bacterial aggregation. All bacterial strains were cultivated at 37°C with shaking in BD Falcon™ conical tubes or in conical flasks. Where needed antimicrobials were

added at the following concentrations ($\mu\text{g/ml}$): hygromycin 50; kanamycin 50; tetracycline 0.02.

Bacterial growth was followed by measurement of absorbance at 580 nm in a Jenway spectrophotometer. Growth of *M. smegmatis* in various media was performed using a Bioscreen Growth Analyser as described previously [34]. Presented Bioscreen data are mean values of two independent experiments done in quintuplicates. Three independent transformants were used for all growth experiments. Viability was assayed by estimation of Colony-Forming Unit (CFU) on 7H10 agar (Becton, Dickinson and Company). Supernatants (SN) were prepared from logarithmic phase cultures of *M. tuberculosis* ($\text{OD}_{580}=0.6-0.8$) grown in Sauton's medium, filter sterilised twice and used for experiments immediately. For dose-dependency experiments culture supernatants were freeze-dried and reconstituted in sterile water. For low magnesium control experiments concentration of MgSO_4 in Sauton's medium was 0.2 mM.

Microscopy. *M. smegmatis* bacilli were mounted on PBS and observed by phase contrast microscopy using a Diphot 300 inverted microscope with a 100 W mercury light source. Images were recorded using a 12/10bit, high-speed Peltier-cooled CCD camera (FDI, Photonic Science) using Image-Pro Plus (Media Cybernetics) software. Labeling of nascent peptidoglycan was done as described previously (35). Briefly, mycobacteria were grown in Sauton's medium in the presence of 20 ng/ml of tetracycline to exponential growth phase ($\text{OD}_{600}\sim 0.5$). A mixture of Vancomycin-BODIPY (Life Technologies) and unlabeled vancomycin (1:1) was added to cultures to the final concentration of 2 $\mu\text{g/ml}$, followed by a

two-hour incubation at 37°C with shaking. Bacteria were washed twice with PBS containing 0.1% Tween 80 and used for fluorescence microscopy.

Generation PknB constructs for overexpression studies. The *pknB* gene and its variants (table S1) were amplified from *M. tuberculosis* H37Rv DNA and cloned into the *Bam*HI and *Spe*I sites of the pMind plasmid [36], for detail of primers see table S2. Constructs were confirmed by sequencing and electroporated into *M. smegmatis* or *M. tuberculosis*. Over-expression of *pknB* or its variants was induced by the addition of tetracycline and confirmed by qRT-PCR.

Preparation of muropeptides. Peptidoglycan from *Escherichia coli* and mycobacteria was isolated and purified as described previously [37, 38]. For growth assays peptidoglycan (2 mg) was digested with mutanolysin, lysozyme, RpfB or MltA at 37°C for up to 72 hours. Muramidases were used at the final concentration of 100 µg/ml. Peptidoglycan digestion with mutanolysin and MltA was carried out in buffer containing 80 mM NaH₂PO₄, pH 4.8; with lysozyme in 25 mM NaH₂PO₄, pH 6.0, and with RpfB in 40 mM sodium citrate, pH 6.5. The presence of muropeptides was confirmed by HPLC. In separate experiments peptidoglycan was sonicated 3 times for 30 s, briefly span down at 100×g to remove big fragments and used for growth assays. Synthetic muropeptides were prepared as described previously [39].

Determination of minimum inhibitory concentration (MIC) for selected antimicrobials. Tests were performed in 96-well microtitre plates using a dilution method [40]. Briefly *M. smegmatis* (1×10⁵ CFU) were inoculated in 100 µl of

Sauton's medium supplemented with kanamycin and tetracycline and a range of concentrations of antimicrobial tested. Plates were incubated three days at 37°C. MIC was determined as the lowest concentration at which no visible growth was detected after 3 days of incubation. Fresh antimicrobial stocks were prepared for each experiment; four independent experiments were done.

Effect of recombinant PASTA protein on growth of *M. tuberculosis*.

Recombinant PASTA was purified as described previously [31] and filter sterilised prior to growth experiments. Mycobacteria (10^3 cells/ml) were inoculated in 16-well microtitre plates containing supplemented 7H9 medium with different concentration of sterile rPknB_PASTA. For each concentration four replicates were inoculated. Sealed plates were incubated at 37°C without shaking. Optical density was measured after 15 and 25 days of incubation. The experiment was repeated twice. Similar results were obtained when mycobacteria were grown in Falcon tubes or conical plastic flasks. *M. tuberculosis* stationary phase culture was stored at 4°C for 2 months and 2×10^4 cells/ml were inoculated in conical flasks. The flasks were incubated at 37°C for up to 12 weeks. Optical density was measured at regular intervals using a Jenway spectrophotometer. The assay was performed in triplicate, three times.

Transcriptional profiling. Total RNA was isolated from 10 ml of mycobacterial cultures from exponential growth phase using the Trizol method [41]. DNA contamination was removed with Turbo DNA-free DNAase (Ambion) before cDNA was generated using Superscript Reverse Transcriptase II (Invitrogen) and gene-specific primers (table S2). Q-PCR was performed in a

Corbett Rotor Gene 6000 real time thermocycler using Absolute QPCR SYBR Green mix (Thermo) and gene-specific primers. Levels of expression were normalized to *16s rRNA* [42].

Protein electrophoresis and western blot. Mycobacteria were collected by centrifugation, washed in PBS and resuspended in buffer, containing, 50 mM Tris HCl, pH 8.0 and 150 mM NaCl. Bacteria were lysed in FastPrep®-24 Instrument (MP Biomedicals, United Kingdom) using glass beads. Lysates were centrifuged at $14000 \times g$ for 20 min to separate soluble and insoluble fractions. Proteins were separated, using 12% SDS PAGE and transferred on nitrocellulose membrane. Primary antibodies were anti-polyhistidine antibodies (Sigma); alkaline-phosphatase conjugate anti-mouse antibodies were used as secondary antibodies. Sigma Fast BCIP/NBT was use as a substrate to visualize recognized proteins.

Isolation of membrane protein fraction for proteomics analysis. Membrane fractions were prepared as described previously [43]. Briefly, *M. smegmatis* TMP and MIND strains were inoculated in 2 L conical flasks containing 500 ml Sauton's medium. The cultures were incubated at 37°C with shaking (200 rpm) for 3 hours. Cultures from two flasks were used for preparation of membrane fractions of each strain. The bacteria were harvested by centrifugation at $6000 \times g$ for 20 min and washed with PBS. The pellets were resuspended in 10 ml of extraction buffer containing 50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 10 mM $MgCl_2$, and stirred on ice for 10 minutes. The cell suspension was sonicated before the addition of 100 µg of DNase I per gram of

cells and stirring on ice for 5 minutes to decrease viscosity. The suspension was centrifuged at $25,000 \times g$ for 25 minutes at 4°C . The pellet was discarded and the supernatant centrifuged at $100,000 \times g$ for 60 min at 4°C . After discarding supernatant the pellet was carefully resuspended in 100 mM sodium carbonate buffer, pH 11.5 and centrifuged at $100,000 \times g$ for 60 min. The centrifugation and resuspension in carbonate buffer steps were repeated a minimum of 3 times before a final resuspension and centrifugation step using sterile MilliQ water. The membrane fractions were resuspended in 1 ml sterile deionized water and freeze dried.

Analysis of membrane proteins. Proteomics was carried out by the University of Leicester Proteomics Facility (PNAFL, University of Leicester, <http://www2.le.ac.uk/colleges/medbiopsych/facilities-and-services/cbs/protein-and-dna-facility/pnacl>). Trypsin digestion of membrane proteins was performed using a filter-aided sample preparation method [44]. LC-MS/MS was carried out using an RSLCnano HPLC system (Dionex, UK) and an LTQ-Orbitrap-Velos mass spectrometer (Thermo Scientific). Peptides were eluted from the trap column at a flow rate of $0.3 \mu\text{l}/\text{min}$ and through a reverse-phase PicoFrit capillary column ($75 \mu\text{m}$ i.d. \times 400 mm) containing Symmetry C18 100 Å media (Waters, UK) that was packed in-house using a high pressure device (Proxeon Biosystems, Denmark) over a period of four hours, with the output of the column sprayed directly into the nanospray ion source of the LTQ-Orbitrap-Velos mass spectrometer.

The raw data file obtained from each LC-MS/MS acquisition was processed using Proteome Discoverer (version 1.4.0.288, Thermo Scientific), searching each file in turn using Mascot (45) (version 2.2.04, Matrix Science Ltd.) against the UniProtKB-Swissprot database. The peptide tolerance was set to 10 ppm and the MS/MS tolerance was set to 0.02 Da. Fixed modifications were set as carbamidomethyl (C) and variable modifications set as oxidation (M). A decoy database search was performed. The output from Proteome Discoverer was further processed using Scaffold Q+S⁴ (version 4.0.5, Proteome Software). For quantitative experiments the digested peptides were labeled with the TMTsixplexTM Label Reagent kit (Life Sciences) according to the manufacturer instruction. Scaffold Q+ (version Scaffold_4.3.4, Proteome Software Inc., Portland, OR) was used to quantify Label Based Quantitation (iTRAQ, TMT, SILAC) peptide and protein identifications. Both protein and peptide identifications were accepted if they could be established at greater than 95.0% probability and contained at least 2 identified peptides. Peptide probabilities from X! Tandem were assigned by the Peptide Prophet algorithm (46) with Scaffold delta-mass correction. Peptide probabilities from Mascot were assigned by the Scaffold Local FDR algorithm. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD002120 and 10.6019/PXD002120

Analytical gel-filtration. 50 µl of PknB_PASTA at a concentration of 50 µM were injected on a S75 10/300 column (GE Life Science) at a 0.4 ml/min flow rate. The column was either equilibrated with 25 mM Tris HCl (pH 8.5), and 100

mM NaCl with or without 25 mM MgSO₄. For the injection in the column in the buffer without MgSO₄ the protein was pre-incubated with 5 mM EDTA. In the case of gel-filtration in the presence of MgSO₄ the protein was pre incubated with 25 mM MgSO₄.

Nuclear Magnetic Resonance Experiments. All NMR experiments were carried out at 20°C on Bruker Avance III 700 (¹H-¹⁵N double resonance experiments) or Avance III 500 (¹H-¹³C-¹⁵N triple-resonance experiments) spectrometer equipped with 5 mm z-gradient TCI cryoprobe, using the standard pulse sequences. NMR samples consist of approximately 50 µM ¹⁵N-labeled protein dissolved in 10 mM phosphate buffer (pH 6.8), 100 mM NaCl, with 5% D₂O for the lock prepared as described elsewhere [31].

Results

The external PASTA domain is important for PknB function. It has been previously shown that overexpression of enzymatically active PknB in mycobacteria results in growth inhibition and alteration of bacterial shape [4]. Additionally the external PknB_PASTA domain is indispensable for functional complementation of PknB conditional mutants of *M. smegmatis* and *M. tuberculosis* [47]. However the precise role of the PknB_PASTA domain in regulation of mycobacterial growth and peptidoglycan biosynthesis remains unclear. In our initial experiments we investigated growth patterns of *M. smegmatis* overexpressing *pknB*.

A full-length *pknB* or its truncated form missing the region encoding the external PASTA domain (*pknB*-ΔPASTA) were cloned into pMind plasmid [35],

containing a tetracycline-regulated promoter (figure 1; table S1). The plasmids were transformed into *M. smegmatis* and the resultant strains, designated as PknB1 (*pknB*), PknB5 (*pknB-ΔPASTA*) and MIND (empty plasmid), grew similarly in all media tested in the absence of the inducer tetracycline (data not shown). However, addition of tetracycline to the PknB1 or PknB5 strains affected growth in liquid medium (figure 2). This inhibitory effect was probably caused by the induction of expression of *pknB* and *pknB-ΔPASTA* from the pMind plasmid. When grown in the presence of tetracycline PknB1 and PknB5 expressed the *pknB* versions from pMind at relatively high level, corresponding to 1% ($\pm 0.3\%$) of all *16S-rRNA* transcripts as judged by quantitative RT-PCR; no expression of the target gene was detected in mycobacteria in the absence of tetracycline. Interestingly, overexpressing the *pknB* versions resulted in different growth characteristics. In all media the PknB1 strain had a longer apparent lag-phase (time to detectable turbidity) and slower growth rate (figure 2; table S3), indicating that overexpression of the truncated PknB missing the external PASTA domain was less toxic than overexpression of the full-length kinase.

To further investigate the role of the PknB_PASTA domain in bacterial growth directly, we cloned the *pknB* region encoding the membrane-anchored external domain of *M. tuberculosis* (designated as *tmp*, figure 1) into the pMind plasmid and transformed this plasmid into *M. smegmatis*. In the absence of tetracycline, both MIND and TMP strains grew similarly (figure 3a, filled symbols) and their cDNA analysed by q-RT-PCR contained similar levels of *tmp* transcripts. However, addition of tetracycline led to overexpression of the *tmp* transcript

(reaching $9.6 \pm 0.8\%$ from the total *16s rRNA* transcripts) in the TMP strain, which resulted in growth inhibition especially in Sauton's medium (table S3; figure 3a, open symbols). The TMP strain required a longer time to initiate bacterial growth, but its growth rate was identical to the one observed in the MIND control strain. This phenotype was highly reproducible and did not change after passage *in vitro* (figure s1). Expression of the PASTA domain without the transmembrane domain (PknB13), the transmembrane domain on its own (PknB9) or the transmembrane domain with one PASTA unit, PknB10 (figure 1) had no significant effect on bacterial growth (figure 3b). Strains overexpressing the PASTA domain with two or three PASTA units (PknB11 and PknB12, respectively) displayed a marginal growth defect. All PASTA variants were expressed at similar levels according to qRT-PCR (data not shown). We generated Myc-6xHis-tagged versions of TMP (designated as TMPH strain; table S1) and of the soluble PASTA domain (designated as PknB14) and investigated their localization by Western blot. TMPH was detected in the envelope fraction, while PknB14 was found in the cytoplasm *M. smegmatis* (figure 3c,d). Importantly, TMPH and TMP strains showed no difference in growth under all conditions described here (data not shown). These results confirm that only the surface exposed domain containing four PASTA units significantly impaired mycobacterial growth.

Overproduction of the PknB_PASTA domain also influenced cellular morphology. Microscopic examination of the strains revealed accumulation of elongated TMP cells of irregular shape (figure 4b, left panel), similar to cells with partially depleted PknB [4], suggesting that the overexpressed PASTA domain

possibly interfered with the PknB function. Furthermore, Vancomycin-BODIPY labelling revealed anomalous distribution of nascent peptidoglycan in TMP mycobacteria (figure 4*b*, middle panel). While in control MIND cells newly synthesized peptidoglycan was mainly localized at the poles and mid-cell (figure 4*a*), TMP cells displayed a diffused staining across the entire cell surface (figure 4*b*). We also observed significant proportions of unlabeled or weakly labeled TMP cells (figure 4*b*). TMP mycobacteria tend to aggregate, suggesting a modified cell surface, delayed cell separation or increased lysis.

Elevated Mg²⁺ ion level eliminates TMP-mediated growth inhibition. The results described above support the functional importance of the PknB_PASTA domain for PknB activity. The structural studies of the PknB_PASTA suggested that this domain could dimerise due to conformational flexibility between the different PASTA domains [31]. It was proposed that the binding of putative ligands could induce the dimerization of kinase domains and consequent trans-activation of PknB [31]. Thus, the PASTA domain could serve as a receptor for signaling molecules (for example, muropeptides) and activate bacterial resuscitation and growth, employing the mechanism previously proposed for *B. subtilis* spore germination [16, 39]. The PknB-PASTA domain missing the functional kinase domain may compete with the native PknB for the ligands and dysregulate the PknB-mediated signaling pathways. Therefore, we investigated whether culture supernatant from growing mycobacteria (potentially containing the putative ligand) would eliminate the growth defect caused by TMP overexpression. Indeed, the growth inhibition was abolished by the addition of

342 culture supernatant in a dose-dependent manner (figure 5a). Further tests
343 showed that the active compound was a low-molecular-weight chemical resistant
344 to heating and it was present in Sauton's medium prior to exposure to bacteria
345 (data not shown). Eventually, we demonstrated that the active entity was Mg^{2+}
346 from $MgSO_4$ or $MgCl_2$. Elevated concentration of Mg^{2+} could completely abolish
347 the inhibition by PASTA-domain overexpression (figure 5b) and the dose-
348 dependent effect of conditioned medium was largely due to increased Mg^{2+}
349 concentrations (data not shown). Supernatants obtained from cultures grown in
350 lower $MgSO_4$ (0.2 mM) medium had only a moderate growth stimulatory effect on
351 both control and TMP strains (figure 5c). The increased Mg^{2+} concentration did
352 not influence expression of *tmp* ($p > 0.05$ t-test) and the production of TMPH, a
353 His-tagged version of TM-PASTA (data not shown). Magnesium also improved
354 growth of PknB1 and PknB5 strains (data not shown).

355 To investigate whether the PknB_PASTA domain was able to bind Mg^{2+} ions,
356 we used nuclear magnetic resonance (NMR) methodology [31]. A ^{15}N labeled
357 sample of recombinant PknB_PASTA (50 μM) was titrated with increasing
358 amounts of $MgCl_2$ (or $MgSO_4$) up to 50 mM. The ion binding either with a lateral
359 chain or the protein backbone would generate chemical-shift perturbation on
360 the 1H - ^{15}N resonance measured in the HSQC experiment [48]. However, no
361 chemical-shift perturbation was observed in the PknB_PASTA upon addition of
362 $MgCl_2$ or EDTA, suggesting the lack of interaction between the protein and Mg^{2+}
363 ions (figure S2), and the addition of $MgSO_4$ to recombinant PASTA did not
364 change the oligomerization state of recombinant PASTA (figure S3). These

results show that the ameliorating effect of Mg^{2+} on TMP-mediated growth inhibition was indirect.

Muropeptides do not eliminate TMP-mediated growth inhibition. In separate experiments we tested the effects of muropeptides on the growth inhibition in the TMP strain. We used peptidoglycan from *Escherichia coli* digested with various muramidases, sonicated or RpfB-digested peptidoglycan from *M. smegmatis*, and synthetic muropeptides, *N*-acetylglucosamine, *N*-acetylmuramic acid and *N*-acetylglucosaminyl-*N*-acetylmuramic acid (table S4). We only observed a minor reduction in the duration of lag phase of TMP strain from 65 to 52 h in the presence of relatively high concentrations of digested or sonicated peptidoglycan (0.5 mg/ml and higher). The observed reduction in lag phase did not depend on the type of peptidoglycan used. Synthetic muropeptides, previously employed for germination of *B. subtilis* spores [39], did not reduce growth inhibition at any concentration tested (10 pM – 50 mM) and did not stimulate growth of *M. smegmatis* (figure 5d). Among other divalent metals tested Ca^{2+} improved the TMP growth, however it could not be used at concentrations above 5 mM due to poor solubility. Since Mg^{2+} ions are known to stabilize cell wall and are frequently used for cultivation of bacteria with defective cell wall [49], we investigated whether the TMP strain was more susceptible to various antimicrobials compared with control.

Over-expression of PknB-PASTA increases sensitivity of *M. smegmatis* to β -lactam antibiotics and alters the abundance of cell wall proteins. We tested minimum inhibitory concentrations (MIC) of various antimicrobial agents

for PknB1, TMP and MIND strains. MIC of rifampicin (inhibition of transcription), ethambutol (arabinogalactan biosynthesis) and streptomycin (protein biosynthesis) were identical in all the strains tested (table 1). However, the TMP strain was remarkably more sensitive to several β -lactam antibiotics: meropenem, ampicillin and clavulanate, an inhibitor of β -lactamase. In the presence of sub-inhibitory concentration of clavulanate (100 μ M) MIC of meropenem for the TMP strain decreased below 0.15 μ M compared to 6.5 μ M in the control strain. Interestingly, overexpression of PknB did not alter antimicrobial susceptibility, suggesting that the PknB-PASTA domain on its own may perturb peptidoglycan synthesis causing higher sensitivity to β -lactams. The perturbation of peptidoglycan synthesis could also explain the prolonged lag-phase observed in the TMP strain. To address this possibility membrane proteins isolated from the MIND and TMP strains in lag-phase were investigated.

In total we identified 1071 proteins in the membrane fractions of both strains using LTQ-Orbitrap Velos mass spectrometry. The TMPH strain had 90 proteins specific for this sample and 95 proteins were detected in the MIND strain only. The majority of proteins (886) were identified in both strains. Importantly, production of the PknB_PASTA domain in the membrane fraction was confirmed by detection of 22 unique fragments with 86% peptide coverage in the TMPH strain (table S5). Most of the proteins detected (705) were assigned as hypothetical with unknown function, while 267 were predicted as membrane or cell surface proteins. This result confirms that the extraction procedure selectively enriched the membrane proteins. A high number of membrane

proteins implicated in transport of metals and nutrients and enzymes was found in the membrane fractions of both strains. Next, peptides were labeled using the TMPsixplexTM kit to compare the relative abundance of membrane proteins in the TMP and MIND (table S5). We focused our analysis on cell division proteins and enzymes involved in cell wall biosynthesis. Most of cell division proteins were equally present in the TMP and MIND strains (data not shown). However, CwsA was more abundant in TMP strain (table 2). The precise function of CwsA is unknown but it has been shown to interact with CrgA, which is annotated as an inhibitor of septum formation [50, 51]. CrgA was detected only in the membrane fractions of TMP but not in those of MIND and its relative abundance could not be calculated. Another CrgA interaction partner, Wag31 was detected at similar levels in both strains. Interestingly, the TMP strain had more enzymes involved in biosynthesis of mycolyl-arabinogalactan (table 2 and table s5), including several mycolyl- and galactofuranosyl-transferases. Regarding peptidoglycan-related enzymes the levels of two metallo- β -lactamases, L, D-transpeptidase LdtB and D-alanyl-D-alanine carboxypeptidase DacB1 were increased in the TMP membrane fraction (table 2). Complete data are available via ProteomeXchange with identifier PXD002120. Collectively these results may suggest that the overexpression of the PASTA domain influences the biosynthesis and architecture of mycobacterial cell wall.

The addition of recombinant PASTA protein abolishes growth of *M. tuberculosis*. The results described above suggest that overexpression of the PknB_PASTA domain can be employed for growth inhibition and increase of

susceptibility to antimicrobials in medically important pathogen *M. tuberculosis*. We therefore investigated whether the PknB_PASTA domain would have a similar effect in *M. tuberculosis*. We transformed pMind *pknB7* and empty pMind plasmids (table s1) into *M. tuberculosis* H37Rv and investigated growth of the resultant strains (TMP and MIND) in the presence of tetracycline. As shown in figure 6a overexpression of the external membrane attached PASTA (TMP) indeed inhibited initiation of *M. tuberculosis* growth. We next studied if addition of recombinant PASTA protein containing PASTA units only (residues 354-626, designated as rPASTA) may alter the growth. We noted that the addition of rPASTA to *M. tuberculosis* resulted in growth defect at concentration of 10 µg/ml (figure 6b) and increased apparent lag phase from 5 to 14 days. This growth inhibition was temporary and the cultures eventually produced normal growth, while addition of recombinant PASTA to logarithmic phase culture had no effect on growth (not shown). However, the growth inhibitory effect of rPASTA was more pronounced in experiments when mycobacteria were stored at 4°C for 2 months (figure 6c). We found that these cells retained culturability and produced normal growth both in liquid and solid media. Nevertheless, their growth initiation was significantly delayed by addition of rPASTA at concentration 10 µg/ml. The mycobacteria were not killed by rPASTA and after two month incubation generated normal stationary phase culture (figure 6c). In all experiments control buffer did not have any inhibitory effect. These results suggest that the PASTA domain may directly interfere with mycobacterial regrowth and therefore presents a plausible target for the design of specific drugs altering bacterial regrowth and

457 resuscitation.

458 **Discussion and conclusions**

459 PknB has been shown to regulate bacterial division, growth and cell wall
460 biosynthesis [10]. The data presented in this study suggest that the PknB-
461 mediated signaling pathway can be targeted through the external PknB_PASTA
462 domain. Overexpression of the PknB_PASTA domain delayed the regrowth of
463 both *M. tuberculosis* and *M. smegmatis*. The growth inhibition was ameliorated
464 by addition of high concentrations of Mg^{2+} but not by mucopeptides. Mg^{2+} ions are
465 crucial for all living organisms and the concentration may vary and reach as high
466 as 20 mM in eukaryotic [52] and a massive 100 mM in prokaryotic cells [53]. In
467 Gram-negative bacteria magnesium ions stabilize the outer membrane [54] and it
468 is found in high concentrations in cell wall of Gram-positive bacteria, which bind
469 divalent ions to their teichoic acids [54]. Mycobacteria do not possess teichoic
470 acids and the binding of Mg^{2+} ions has not been studied in the mycobacterial cell
471 wall in detail, although Mg^{2+} ions improved growth of mycobacteria in media with
472 low pH [56]. Growth of *B. subtilis* mutants missing certain peptidoglycan binding
473 proteins was improved at high Mg^{2+} levels, suggesting that divalent metals may
474 compensate growth defects due to altered peptidoglycan biosynthesis [57].

475 The precise mechanism of the magnesium effect on growth of TMP strain is
476 unknown. Mg^{2+} ions may stabilize the cellular membrane and protein complexes
477 associated with the membrane disturbed by over-expression of PknB_PASTA
478 domain. Though it is also possible that Mg^{2+} ions directly bind to the growing
479 strands of stems of peptidoglycan and promotes their polymerization. Further

dissection of molecular mechanisms underlying inhibitory effect of the PknB_PASTA overexpression will facilitate our understanding of biological function of PknB and its extracellular domain as well as the roles of Mg^{2+} ions in cell wall architecture. Three possible scenarios may explain the observed effects associated with overexpression of PknB_PASTA (figure 7). (i) The overexpressed PknB_PASTA competes for the ligand (peptidoglycan) with a native PknB and therefore interferes with the PknB-mediated signaling pathway (figure 7a). (ii) The PknB_PASTA domain interacts with penicillin-binding proteins and dysregulates peptidoglycan biosynthesis (figure 7b). (iii) Finally, highly abundant PknB_PASTA may directly disrupt the structure of peptidoglycan by binding to the growing strand and physically prevent extension and cross-linking (figure 7c).

The regulatory function of PknB (phosphorylation of cell-wall enzymes and division factors) is evolved to coordinate cell-wall biosynthesis and division. In fact, a recently published study established that PknB coordinated assembly of peptidoglycan synthesizing complex via recruiting FhaA and phosphorylating MviN, a protein essential for mycobacterial growth [26].

In *Listeria monocytogenes* the inactivation of a PASTA-containing kinase, a homologue of PknB, resulted in increased susceptibility to β -lactam antimicrobials [58], as in the TMP strain which presumably shows high susceptibility to β -lactam antimicrobials due to partial inactivation of PknB function. However, while the PknB mutant lyses and is not able to grow, the TMP strain only shows an initial impairment of growth in form of a prolonged lag-phase

but is able to grow normally in exponential phase. Thus, other factors apart from interference with PknB function may contribute to the growth phenotype of the TMP strain.

A possible direct interaction of PknB_PASTA with penicillin-binding proteins is in accordance with recent findings that the PASTA domain of StkP kinase from *S. pneumoniae* was able to bind penicillin-binding protein 2x [59]. We also cannot exclude that the PASTA domain changes the architecture of the cell wall by interacting directly with peptidoglycan (figure 7c) as it was previously observed *in vitro* [16, 60]. The results presented in this study may suggest that the overexpressed PknB_PASTA domain interacts with growing strands of peptidoglycan and possibly physically interferes with its cross-linking (figure 7c), culminating in higher susceptibility of the overexpressing strain to meropenem, clavulanate and ampicillin. Meropenem has been shown to inhibit both DD-carboxypeptidase and LD-transpeptidase activities [61] and in combination with clavulanate it is highly active against growing and persisting mycobacteria [62].

Our findings are consistent with the previously proposed peptidoglycan-sensing role of PknB_PASTA [11, 30]. Further investigation is required to clarify whether the PknB_PASTA domain recognizes the specific structures present in the region of peptidoglycan growth and therefore activates dimerisation of the catalytic domains or whether it plays a more structural role in supporting growing peptidoglycan and ensuring proper localization of PknB. Strains over-expressing truncated forms of the PASTA domain did not have pronounced growth defect (figure 3b), suggesting that four PASTA units and the transmembrane region are

required for PknB functionality. Removal of the transmembrane part resulted in mislocalization of the PASTA domain. Although the precise role of the individual PASTA units awaits elucidation our data are consistent with the previous observation that the truncated PknB missing one or more PASTA domains was not able to complement the conditional *pknB* mutant [47]. It also remains unclear whether the PASTA-domain can sense short muropeptides released by mycobacteria during growth. In our experiments we obtained a very modest effect of peptidoglycan on the growth inhibition caused by PASTA overproduction and no stimulation of growth or resuscitation by externally added muropeptides. It is possible that specific muropeptides are required for activation of PknB and subsequent resuscitation, since muropeptides have been shown to possess resuscitation-stimulatory activity [63]. It is conceivable that the PASTA domain may be directly involved in resuscitation by sensing *in vivo* rearrangement of peptidoglycan, possibly due to the action of resuscitation-promoting factors [34, 64] and other muramidases, and regulation of cell-wall biosynthesis. Additional experiments on monitoring rearrangement of peptidoglycan and interaction between the external PASTA-domain and stem peptidoglycan, using probes for microscopy of living bacteria such as highly sensitive atomic force microscopy cantilevers will allow us to establish the precise PknB-mediated sensing mechanism [65].

The datasets supporting this article have been uploaded as part of the Supplementary Material.

Conflict of interest

549 The authors declare that they have no conflict of interest.

550 **Authors' contribution**

551 OT conceived of the study, carried out molecular work and growth experiments,
552 analysed data and drafted the manuscript; JL carried out molecular work,
553 prepared peptidoglycan and muropeptides; CHJ carried out molecular work and
554 prepared samples for proteomics study; PB carried out protein purification and
555 NMR experiments; DM performed and analysed growth experiments; FF and DG
556 contributed to molecular work and helped draft the manuscript; DH and ML
557 synthesized and purified muropeptides; AB carried out proteomics experiments
558 and analysed proteomics data; WV coordinated peptidoglycan and muropeptide
559 experiments, provided muropeptides and helped draft the manuscript; SM
560 coordinated synthetic muropeptide experiments, provided synthetic muropeptides
561 and helped draft the manuscript; MCG conceived of the study, designed the
562 study and coordinated protein work; GM conceived of the study, designed the
563 study, analysed data, coordinated the study and drafted the manuscript.

564 **Acknowledgements**

565 We are grateful to Roger Buxton for critical reading of this manuscript and to
566 Brian Robertson for the pMind plasmid. *Mtb* DNA was provided by Colorado
567 State University (Contract No.HHSN266200400091C; NIH, NIAID N01-AI-
568 40091). We acknowledge the Centre for Core Biotechnology Services at the
569 University of Leicester for support with Containment Level 3 experiments and the
570 University of Leicester Proteomics Facility for analysis of membrane proteins.

Funding

This study was supported by BBSRC grant BB/H008586/1 (OT and GVM), MRC-DTG (JL), the National Institutes of Health grant AI090348 (DH, ML and SM) and the EC DIVINOCELL project (WV).

References

1. Pereira SFF, Goss L, Dworkin J. 2011 Eukaryote-Like Serine/Threonine Kinases and Phosphatases in Bacteria. *Microbiol. Mol. Biol. R* **75**,192-212. (doi: 10.1128/MMBR.00042-10)
2. Chao J, Wong D, Zheng X, Poirier V, Bach H, Hmama Z, Av-Gay Y. 2010 Protein kinase and phosphatase signaling in *Mycobacterium tuberculosis* physiology and pathogenesis. *Biochim. Biophys. Acta* **1804**, 620-627. (doi: 10.1016/j.bbapap.2009.09.008)
3. Fernandez P, Saint-Joanis B, Barilone N, Jackson M, Gicquel B, Cole ST, Alzari PM. 2006 The Ser/Thr protein kinase PknB is essential for sustaining mycobacterial growth. *J. Bacteriol.* **188**, 7778-7784. (doi: 10.1128/JB.00963-06)
4. Kang CM, Abbott DW, Park ST, Dascher CC, Cantley LC, Husson RN. 2005 The *Mycobacterium tuberculosis* serine/threonine kinases PknA and PknB: substrate identification and regulation of cell shape. *Genes Dev.* **19**, 1692-1704. (doi: 10.1101/gad.1311105)
5. Forti F, Crosta A, Ghisotti D. 2009 Pristinamycin-inducible gene regulation in mycobacteria. *J. Biotechnol.* **140**, 270-277. (doi:10.1016/j.jbiotec.2009.02.001)

- 594 6. Kumar D, Narayanan S. 2012 PknE, a serine/threonine kinase of
595 *Mycobacterium tuberculosis* modulates multiple apoptotic paradigms.
596 *Infect. Genet. Evol.* **12**, 737-747. (doi: 10.1016/j.meegid.2011.09.008)
- 597 7. Walburger A, Koul A, Ferrari G, Nguyen L, Prescianotto-Baschong C,
598 Huygen K, Klebl B, Thompson C, Bacher G, Pieters J. 2004 Protein kinase
599 G from pathogenic mycobacteria promotes survival within macrophages.
600 *Science* **304**, 1800-1804. (doi: 10.1126/science.1099384)
- 601 8. Cowley S, Ko M, Pick N, Chow R, Downing KJ, Gordhan BG, Betts JC,
602 Mizrahi V, Smith DA, Stokes RW, Av-Gay Y. 2004 The *Mycobacterium*
603 *tuberculosis* protein serine/threonine kinase PknG is linked to cellular
604 glutamate/glutamine levels and is important for growth *in vivo*. *Mol.*
605 *Microbiol.* **52**, 1691-1702. (doi: 10.1111/j.1365-2958.2004.04085.x)
- 606 9. Gómez-Velasco A, Bach H, Rana AK, Cox LR, Bhatt A, Besra GS, Av-Gay
607 Y. 2013 Disruption of the serine/threonine protein kinase H affects
608 phthiocerol dimycocerosates synthesis in *Mycobacterium tuberculosis*.
609 *Microbiology* **159**, 726-736. (doi: 10.1099/mic.0.062067-0)
- 610 10. Molle V, Kremer L .2010 Division and cell envelope regulation by Ser/Thr
611 phosphorylation: Mycobacterium shows the way. *Mol. Microbiol.* **75**, 1064-
612 1077. (doi: 10.1111/j.1365-2958.2009.07041)
- 613 11. Jones G, Dyson P. 2006 Evolution of transmembrane protein kinases
614 implicated in coordinating remodeling of gram-positive peptidoglycan:
615 inside versus outside. *J. Bacteriol.* **188**, 7470-7476.
616 (doi: 10.1128/JB.00800-06)

- 617 12. Debarbouille M, Dramsi S, Dusserget O, Nahori MA, Vaganay E, Jouvion
618 G, Gazzzone A, Msadek T, Duclos B. 2009 Characterization of a
619 serine/threonine kinase involved in virulence of *Staphylococcus aureus*. *J.*
620 *Bacteriol.* **191**: 4070-4081. (doi: 10.1128/JB.01813-08)
- 621 13. Banu LD, Conrads G, Rehrauer H, Hussain H, Allan E, van der Ploeg JR.
622 2010 The *Streptococcus mutans* serine/threonine kinase, PknB, regulates
623 competence development, bacteriocin production, and cell wall
624 metabolism. *Infect. Immun.* **78**, 2209-2220. (doi: 10.1128/IAI.01167-09)
- 625 14. Beilharz K, Nováková L, Fadda D, Branny P, Massidda O, Veening JW.
626 2012 Control of cell division in *Streptococcus pneumoniae* by the
627 conserved Ser/Thr protein kinase StkP. *Proc. Natl. Acad. Sci. U S A* **109**:
628 E905-913. (doi: 10.1073/pnas.1119172109)
- 629 15. Gaidenko TM, Kim TJ, Price CW. 2002 The PrpC serine-threonine
630 phosphatase and PrkC kinase have opposing physiological roles in
631 stationary-phase *Bacillus subtilis* cells. *J. Bacteriol.* **184**, 6109-6114.
632 (doi: 10.1128/JB.184.22.6109-6114.2002)
- 633 16. Shah IM, Laaberki MH, Popham DL, Dworkin J. 2008 A eukaryotic-like
634 Ser/Thr kinase signals bacteria to exit dormancy in response to
635 peptidoglycan fragments. *Cell* **135**, 486-496.
636 (doi: 10.1016/j.cell.2008.08.039)
- 637 17. Shah IM, Dworkin J. 2010 Induction and regulation of a secreted
638 peptidoglycan hydrolase by a membrane Ser/Thr kinase that detects

- 639 muropeptides. *Mol. Microbiol.* **75**, 1232-1243. (doi: 10.1111/j.1365-
640 2958.2010.07046.x)
- 641 18. Jones G, Del Sol R, Dudley E, Dyson P. 2011 Forkhead-associated
642 proteins genetically linked to the serine/threonine kinase PknB regulate
643 carbon flux towards antibiotic biosynthesis in *Streptomyces coelicolor*.
644 *Microbial. Biotechnol.* **4**, 263-274. (doi: 10.1111/j.1751-
645 7915.2010.00237.x)
- 646 19. Schultz C, Niebisch A, Schwaiger A, Viets U, Metzger S, Bramkamp M,
647 Bott M. 2009 Genetic and biochemical analysis of the serine/threonine
648 protein kinases PknA, PknB, PknG and PknL of *Corynebacterium*
649 *glutamicum*: evidence for non-essentiality and for phosphorylation of OdhI
650 and FtsZ by multiple kinases. *Mol. Microbiol.* **74**, 724-741.
651 (doi: 10.1111/j.1365-2958.2009.06897.x)
- 652 20. Molle V, Gulten G, Vilch  ze C, Veyron-Churlet R, Zanella-Cl  on I,
653 Sacchettini JC, Jacobs WR Jr, Kremer L. 2010 Phosphorylation of InhA
654 inhibits mycolic acid biosynthesis and growth of *Mycobacterium*
655 *tuberculosis*. *Mol. Microbiol.* **78**, 1591-1605.
656 (doi: 10.1111/j.1365-2958.2010.07446.x)
- 657 21. Dasgupta A, Datta P, Kundu M, Basu J. 2006 The serine/threonine kinase
658 PknB of *Mycobacterium tuberculosis* phosphorylates PBPA, a penicillin-
659 binding protein required for cell division. *Microbiology* **152**, 493-504.
660 (doi: 10.1099/mic.0.28630-0)

- 661 22. Veyron-Churlet R, Zanella-Cleon I, Cohen-Gonsaud M, Molle V, Kremer L.
662 2010 Phosphorylation of the *Mycobacterium tuberculosis* beta-ketoacyl-
663 acyl carrier protein reductase MabA regulates mycolic acid biosynthesis.
664 *J. Biol. Chem.* **285**, 12714-12725. (doi: 10.1074/jbc.M110.105189)
- 665 23. Park ST, Kang CM, Husson RN. 2008 Regulation of the SigH stress
666 response regulon by an essential protein kinase in *Mycobacterium*
667 *tuberculosis*. *Proc. Natl. Acad. Sci. U S A* **105**, 13105-13110.
668 (doi: 10.1073/pnas.0801143105)
- 669 24. Ventura M, Rieck B, Boldrin F, Degiacomi G, Bellinzoni M, Barilone N,
670 Alzaidi F, Alzari PM, Manganelli R, O'Hare HM. 2013 GarA is an essential
671 regulator of metabolism in *Mycobacterium tuberculosis*. *Mol. Microbiol.* **90**,
672 356-366. (doi: 10.1111/mmi.12368)
- 673 25. Roumestand C, Leiba J, Galoppe N, Margeat E, Padilla A, Bessin Y,
674 Barthe P, Molle V, Cohen-Gonsaud M. 2011 Structural insight into the
675 *Mycobacterium tuberculosis* Rv0020c protein and its interaction with the
676 PknB kinase. *Structure* **19**, 1525-1534. (doi: 10.1016/j.str.2011.07.011)
- 677 26. Gee CL, Papavinasasundaram KG, Blair SR, Baer CE, Falick AM, King
678 DS, Griffin JE, Venghatakrishnan H, Zukauskas A, Wei JR, Dhiman RK,
679 Crick DC, Rubin EJ, Sassetti CM, Alber T. 2012 A phosphorylated
680 pseudokinase complex controls cell wall synthesis in mycobacteria. *Sci.*
681 *Signal.* **5**, ra7. (doi: 10.1126/scisignal.2002525)
- 682 27. Ortega C, Liao R, Anderson LN, Rustad T, Ollodart AR, Wright AT,
683 Sherman DR, Grundner C. 2014 *Mycobacterium tuberculosis* Ser/Thr

- 684 protein kinase B mediates an oxygen-dependent replication switch. *PLoS*
685 *Biol.* **12**, e1001746. (doi: 10.1371/journal.pbio.1001746)
- 686 28. Loughheed KE ,Osborne SA, Saxty B, Whalley D, Chapman T, Bouloc N,
687 Chugh J, Nott TJ, Patel D, Spivey VL, Kettleborough CA, Bryans JS,
688 Taylor DL, Smerdon SJ, Buxton RS. 2011 Effective inhibitors of the
689 essential kinase PknB and their potential as anti-mycobacterial agents.
690 *Tuberculosis* **91**, 277-286. (doi: 10.1016/j.tube.2011.03.005)
- 691 29. Mieczkowski C, Iavarone AT, Alber T. 2008 Auto-activation mechanism of
692 the *Mycobacterium tuberculosis* PknB receptor Ser/Thr kinase. *EMBO J.*
693 **27**,3186-3197. (doi: 10.1038/emboj.2008.236)
- 694 30. Yeats C, Finn RD, Bateman A .2002 The PASTA domain: a β -lactam-
695 binding domain. *Trends Biochem. Sci.* **27**, 438-440.
696 (doi: 10.1016/S0968-0004(02)02164-3)
- 697 31. Barthe P, Mukamolova GV, Roumestand C, Cohen-Gonsaud M. 2010 The
698 structure of PknB extracellular PASTA domain from *Mycobacterium*
699 *tuberculosis* suggests a ligand-dependent kinase activation. *Structure* **18**,
700 606-615. (doi: 10.1016/j.str.2010.02.013)
- 701 32. Squeglia F, Marchetti R, Ruggiero A, Lanzetta R, Marasco D, Dworkin J,
702 Petoukhov M, Molinaro A, Berisio R, Silipo A. 2011 Chemical basis of
703 peptidoglycan discrimination by PrkC, a key kinase involved in bacterial
704 resuscitation from dormancy. *J. Am. Chem. Soc.* **133**, 20676-20679.
705 (doi: 10.1021/ja208080r)
- 706 33. Mir M, Asong J, Li X, Cardot J, Boons GJ, Husson RN. 2011 The

- 707 extracytoplasmic domain of the *Mycobacterium tuberculosis* Ser/Thr
708 kinase PknB binds specific mucopeptides and is required for PknB
709 localization. *PLoS Pathog.* **7**: e1002182.
710 (doi:10.1371/journal.ppat.1002182)
- 711 34. Mukamolova GV, Turapov OA, Young DI, Kaprelyants AS, Kell DB, Young
712 M. 2002 A family of autocrine growth factors in *Mycobacterium*
713 *tuberculosis*. *Mol. Microbiol.* **46**, 623-635. (doi: 10.1046/j.1365-
714 2958.2002.03184.x35)
- 715 35. Joyce G, Williams KJ, Robb M, Noens E, Tizzano B, Shahrezaei V,
716 Robertson BD. 2012 Cell division site placement and asymmetric growth
717 in mycobacteria. *PLoS ONE* **7**: e44582.
718 (doi:10.1371/journal.pone.0044582)
- 719 36. Blokpoel MC, Murphy HN, O'Toole R, Wiles S, Runn ES, Stewart GR,
720 Young DB, Robertson BD. 2005. Tetracycline-inducible gene regulation in
721 mycobacteria. *Nucleic Acids Res.* **33**, e22. (doi: 10.1093/nar/gni023)
- 722 37. Glauner B, Holtje JV, Schwarz U. 1988 The composition of the murein of
723 *Escherichia coli*. *J. Biol. Chem.* **263**, 10088-10095.
- 724 38. Mahapatra S, Crick DC, McNeil MR, Brennan PJ. 2008 Unique structural
725 features of the peptidoglycan of *Mycobacterium leprae*. *J. Bacteriol.* **190**:
726 655-661. (doi: 10.1128/JB.00982-07)
- 727 39. Lee M, Hesek D, Shah IM, Oliver AG, Dworkin J, Mobashery S. 2010
728 Synthetic peptidoglycan motifs for germination of bacterial spores. *Chem.*
729 *Bio. Chem.* **11**, 2525-2529. (doi: 10.1002/cbic.201000626)

- 730 40. Jorgensen JH, Turnidges JD, Washington JA. 1999 Antibacterial
731 susceptibility tests: dilution and disk diffusion methods. In *Manual of*
732 *Clinical Microbiology* (Murray PR, Baron EJ, Pfaller MA, Tenover FC,
733 Yolken RH. Washington Eds), ASM, DC, 1526-1543.
- 734 41. Waddell SJ, Butcher PD. 2010 Use of DNA arrays to study transcriptional
735 responses to antimycobacterial compounds. In *Antibiotic Resistance*
736 *Protocols* (SH Gillespie and TD McHugh Eds.) *Methods Mol. Biol.* **642**, 75-
737 91. (doi: 10.1007/978-1-60327-279-7_6)
- 738 42. Turapov O, Waddell SJ, Burke B, Glenn S, Sarybaeva A, Tudo G,
739 Labesse G, Young DI, Young M, Andrew PW, Butcher PD, Cohen-
740 Gonsaud M, Mukamolova GV. 2014 Antimicrobial treatment improves
741 mycobacterial survival in non-permissive growth conditions. *Antimicrobial*
742 *Agents Chemother.* **58**, 2798-806.
743 (doi: 10.1128/AAC.02774-13)
- 744 43. Lee BY, Hefta SA, Brennan PJ. 1992 Characterization of the major
745 membrane protein of virulent *Mycobacterium tuberculosis*. *Infect. Immun.*
746 **60**, 2066-2074.
- 747 44. Wisniewski JR, Zougman A, Nagaraj N, Mann M. 2009 Universal sample
748 preparation method for proteome analysis. *Nat. Methods* **6**, 359-362.
749 (doi:10.1038/nmeth.1322)
- 750 45. Perkins DN, Pappin DJ, Creasy DM, Cottrell JS. 1999 Probability-based
751 protein identification by searching sequence databases using mass
752 spectrometry data. *Electrophoresis* **20**, 3551-3567.

- 753 (doi: 10.1002/(SICI)1522-2683(19991201)20:18<3551::AID-
754 ELPS3551>3.0.CO;2-2)
- 755 46. Keller A, Nesvizhskii AI, Kolker E, Aebersold R. 2002 Empirical statistical
756 model to estimate the accuracy of peptide identification made by MS/MS
757 and database search. *Anal. Chem.* **74**, 5383-5392.
758 (doi: 10.1021/ac025747h)
- 759 47. Chawla Y, Upadhyay S, Khan S, Nagarajan SN, Forti F, Nandicoori VK.
760 2014 Protein kinase B (PknB) of *Mycobacterium tuberculosis* is essential
761 for growth of the pathogen in vitro as well as for survival within the host. *J.*
762 *Biol. Chem.* **289**,13858-13875. (doi: 10.1074/jbc.M114.563536)
- 763 48. Ohashi W, Inouye S, Yamazaki T, Hirota H. 2005 NMR analysis of the
764 Mg²⁺-binding properties of aequorin, a Ca²⁺-binding photoprotein. *J.*
765 *Biochem.* **138**, 613-620. (doi: 10.1093/jb/mvi164)
- 766 49. Horwitz AH, Casida LE, Jr. 1978 Survival and reversion of a stable L form
767 in soil. *Can. J. Microbiol.* **24**, 50-55.
- 768 50. Del Sol R, Mullins JG, Grantcharova N, Flardh K, Dyson P. 2006 Influence
769 of CrgA on assembly of the cell division protein FtsZ during development
770 of *Streptomyces coelicolor*. *J. Bacteriol.* **188**,1540-1550.
771 (doi: 10.1128/JB.188.4.1540-1550.2006)
- 772 51. Plocinski P, Arora N, Sarva K, Blaszczyk E, Qin H, Das N, Plocinska R,
773 Ziolkiewicz M, Dziadek J, Kiran M, Gorla P, Cross TA, Madiraju M,
774 Rajagopalan M. 2012 *Mycobacterium tuberculosis* CwsA interacts with
775 CrgA and Wag31, and the CrgA-CwsA complex is involved in

- 776 peptidoglycan synthesis and cell shape determination. *J. Bacteriol.* **194**,
777 6398-6409. (doi: 10.1128/JB.01005-12)
- 778 52. Jahnen-Dechent W, Kettler M. 2012 Magnesium basics. *Clin. Kidney J* **5**,
779 I3-i14. (doi: 10.1093/ndtplus/sfr163)
- 780 53. Moncany ML, Kellenberger E. 1981 High magnesium content of
781 *Escherichia coli* B. *Experientia* **37**: 846-847.
- 782 54. Yamaguchi A, Yanai M, Tomiyama N, Sawai T. 1986 Effects of
783 magnesium and sodium ions on the outer membrane permeability of
784 cephalosporins in *Escherichia coli*. *FEBS Lett.* **208**, 43-47.
- 785 55. Kern T, Giffard M, Hediger S, Amoroso A, Giustini C, Bui NK, Joris B,
786 Bougault C, Vollmer W, Simorre JP. 2010 Dynamics characterization of
787 fully hydrated bacterial cell walls by solid-state NMR: evidence for
788 cooperative binding of metal ions. *J. Am. Chem. Soc.* **132**: 10911-10919.
789 (doi: 10.1021/ja104533w)
- 790 56. Piddington DL, Kashkouli A, Buchmeier NA. 2000 Growth of
791 *Mycobacterium tuberculosis* in a defined medium is very restricted by acid
792 pH and Mg²⁺ levels. *Infect. Immun.* **68**:4518-4522.(doi:
793 10.1128/IAI.68.8.4518-4522.2000)
- 794 57. Murray T, Popham DL, Setlow P. 1998 *Bacillus subtilis* cells lacking
795 penicillin-binding protein 1 require increased levels of divalent cations for
796 growth. *J. Bacteriol.* **180**,4555-4563.
- 797 58. Pensinger DA, Aliota MT, Schaenzer AJ, Boldon KM, Ansari IU, Vincent
798 WJ, Knight B, Reniere ML, Striker R, Sauer JD. 2014 Selective

- 799 pharmacologic inhibition of a PASTA kinase increases *Listeria*
800 *monocytogenes* susceptibility to β -lactam antibiotics. *Antimicrobial Agents*
801 *Chemother.* **58**, 4486-4494. (doi: 10.1128/AAC.02396-14)
- 802 59. Morlot C, Bayle L, Jacq M, Fleurie A, Tourcier G, Galisson F, Vernet T,
803 Grangeasse C, Di Guilmi AM. 2013 Interaction of Penicillin-Binding
804 Protein 2x and Ser/Thr protein kinase StkP, two key players in
805 *Streptococcus pneumoniae* R6 morphogenesis. *Mol. Microbiol.* **90**, 88-
806 102.
807 (doi: 10.1111/mmi.12348)
- 808 60. Maestro B, Novaková L, Hesek D, Lee M, Leyva E, Mobashery S, Sanz
809 JM, Branny P. 2011 Recognition of peptidoglycan and β -lactam antibiotics
810 by the extracellular domain of the Ser/Thr protein kinase StkP from
811 *Streptococcus pneumoniae*. *FEBS Lett.* **585**, 357-363.
812 (doi: 10.1016/j.febslet.2010.12.016)
- 813 61. Kumar P, Arora K, Lloyd JR, Lee IY, Nair V, Fischer E, Boshoff HI, Barry
814 CE 3rd. 2012 Meropenem inhibits D,D-carboxypeptidase activity in
815 *Mycobacterium tuberculosis*. *Mol. Microbiol.* **86**, 367-381.
816 (doi: 10.1111/j.1365-2958.2012.08199.x)
- 817 62. England K, Boshoff HI, Arora K, Weiner D, Dayao E, Schimel D, Via LE,
818 Barry CE 3rd. 2012 Meropenem-clavulanic acid shows activity against
819 *Mycobacterium tuberculosis in vivo*. *Antimicrobial Agents Chemother.* **56**,
820 3384-3387. (doi: 10.1128/AAC.05690-11)

821 63. Nikitushkin VD, Demina GR, Shleeva MO, Kaprelyants AS. 2013
822 Peptidoglycan fragments stimulate resuscitation of "non-culturable"
823 mycobacteria. *Antonie van Leeuwenhoek* **103**, 37-46.
824 (doi: 10.1007/s10482-012-9784-1)

825 64. Kana BD, Mizrahi V. 2010 Resuscitation-promoting factors as lytic
826 enzymes for bacterial growth and signaling. *FEMS Immunol. Med.*
827 *Microbiol.* **58**, 39-50. (doi: 10.1111/j.1574-695X.2009.00606.x)

828 65. Longo G, Alonso-Sarduy L, Rio LM, Bizzini A, Trampuz A, Notz J, Dietler
829 G, Kasas S. 2013 Rapid detection of bacterial resistance to antibiotics
830 using AFM cantilevers as nanomechanical sensors. *Nat. Nanotechnol.* **8**,
831 522-526. (doi: 10.1038/nnano.2013.120)

832

833 **Table captions**

834 **Table 1.** Effect of PknB-PASTA over-expression on antimicrobial susceptibility of
835 *M. smegmatis*

836 **Table 2.** Cell wall enzymes and proteins differently abundant in TMP compared
837 to MIND

838

839 **Figure legends**

840 **Figure 1.** Schematic representation of PknB constructs and strains generated in
841 this study. Kinase: N-terminal kinase domain (aa 1-279); TM: transmembrane
842 region (aa 331-354); TM-PASTA: Penicillin And Serine or Threonine kinase

Associated domain (aa 331-627); TM-P1 (aa 331-423); MYC-HIS: His-Tag; PASTA (aa 354-627).

Figure 2. Effect of *pknB* and *pknB-ΔPASTA* overexpression on growth of *M. smegmatis*. Mycobacteria (5×10^6) from early stationary phase were inoculated in 100 well honeycomb plates, containing various media and incubated in Bioscreen Growth analyser at 37°C with shaking for 5 days in the presence of tetracycline. Presented data are mean values of 5 replicates from 2 independent experiments. Standard deviations were 10% or less of average values and are not shown for clarity.

Figure 3. Effect of PknB_PASTA overexpression on *M. smegmatis* growth. (a) Comparative growth kinetics of *M. smegmatis* strains in Sauton's medium. The same inoculum ($\sim 2 \times 10^6$ cells/well) was used to seed induced (+ tetracycline) and control cultures, respectively. (b) Over-expression of non-secreted PASTA domains or trans-membrane domain on its own does not affect mycobacterial growth. (c) Detection of PknB_TM-PASTA-6×HIS in membrane fraction of *M. smegmatis* strains by western blotting. (d) Detection of the PASTA domain missing the transmembrane region in cytoplasmic fraction of *M. smegmatis* strains by western blotting.

Figure 4. Overexpression of PknB_PASTA results in alteration of bacterial morphology and vancomycin labelling. Mycobacteria from logarithmic growth phase were labeled with Vancomycin-BODIPY conjugate, washed and mounted in PBS and examined using a Diphot 300 inverted microscope. Three independent fields for each strain are shown. Bars indicate 5 μm. A- MIND; B –

866 TMP. Left panel – phase contrast, middle panel – fluorescence microscopy, right
867 panel – merged images. Arrows indicated labeled peptidoglycan.

868 **Figure 5.** Culture supernatant and high concentration of Mg^{2+} abolish the
869 inhibitory effect caused by PknB-PASTA over-expression. (a) Culture
870 supernatants abolish growth inhibitory effect of PknB_TM-PASTA over-
871 expression in a dose-dependent manner. Relative concentration of culture
872 supernatant added is expressed as “fold concentration”. A value of 1-fold
873 corresponds to undiluted culture supernatant. The apparent lag phase was
874 calculated as a period of time when culture reached OD_{600nm} 0.1. (b) Mg^{2+} (10
875 mM) relieves growth inhibition of TMP in the manner comparable to undiluted
876 culture supernatant. (c) Effect of low Mg^{2+} supernatant on apparent lag-phase of
877 TMP strain. (d) Effect of synthetic muropeptides on growth of *M. smegmatis*.
878 Mycobacteria from early stationary phase were washed in Sauton’s medium
879 twice and 5×10^6 bacteria were inoculated in Sauton’s medium, containing culture
880 supernatant or synthetic muropeptides at final concentration of 10 μ M. MPP –
881 MurNAc – Ipentapeptide, DPP – GlcNAc-MurNAc-peptapeptide, ADPP –
882 GlcNAc-1,6 anhydromuramyl-pentapeptide, TPP – tetra-saccharide-peptide, SN
883 – 50% (v/v) culture supernatant. GlcNAc, *N*-acetylglucosamine; MurNAc, *N*-
884 acetylmuramic acid. pentapeptide L-Ala- γ -D-Glu-*m*-DAP-D-Ala-D-Ala. (b) and (d)
885 Presented data are mean values of 5 replicates from 2 independent experiments.
886 Standard deviations were 10% or less of average values and are not shown for
887 clarity. SN - media were supplemented with culture supernatant.

Figure 6. PknB_PASTA reduces growth of *Mycobacterium tuberculosis*. (a) Overexpression of PknB_PASTA domain delays growth initiation. Mycobacteria ($\sim 1 \times 10^6$) were inoculated in supplemented 7H9 medium containing hygromycin and tetrocycline. (b) Effect of different concentrations of rPASTA on *M. tuberculosis* growth. Mycobacteria (10^3 cells/ml) were inoculated in supplemented 7H9 medium containing different concentration of sterile rPASTA. The optical density was measured after 21 days of incubation. (c) Regrowth of stored *M. tuberculosis* was delayed by the addition of rPASTA. *M. tuberculosis* (2×10^4 cells/ml) were inoculated in supplemented 7H9 medium. Buffer was used as control and rPASTA was added at final concentration of 10 μ g/ml. B, C Each data point is an average value of 3 biological replicates, error bars indicate standard deviations. Experiments were performed three times and results of one typical experiment are shown.

Figure 7. Schematic representation of possible mechanisms of PknB_PASTA mediated growth inhibition. (a) In the TMP strain the overexpressed PknB_PASTA competes with native PknB for peptidoglycan ligands and interferes with PknB-mediated signalling. (b) Overexpressed PknB_PASTA binds penicillin-binding proteins and dysregulates peptidoglycan polymerization. (c) Overexpressed PknB_PASTA binds to peptidoglycan and destabilizes its structure. PGS – peptidoglycan-synthesis complex.

Table 1. Effect of PknB-PASTA over-expression on antimicrobial susceptibility of *M. smegmatis*

Strain	Minimum Inhibitory concentration against antibiotic (µM)						
	Rifampicin	Streptomycin	Ethambutol	Meropenem	Clavulanate	Meropenem + Clavulanate*	Ampicillin
MIND	1.2	0.4	2.5	13-19.5	>1000	6.5-13.0	140-170
PknB1	1.2	0.4	2.5	13-19.5	>1000	6.5-13.0	140-170
TMP	1.2	0.4	2.5	6.5	150-200	0.13	30-45

*Clavulanate was added at a concentration of 100 µM.

Table 2. Cell wall enzymes and proteins differently abundant in TMP compared to MIND

<i>M. tb</i> protein	<i>M. smegmatis</i> protein	Protein description	Ratio TMP/MIND Log2 fold
Rv0014c	MSMEG_0028	His-Tagged TM-PASTA	5.9
Rv0008c	MSMEG_0023	Cell wall protein CwsA	1.0
Rv3804c	MSMEG_6398	FbpA mycolyl transferase	1.0
Rv3790	MSMEG_6382	DprE1 decaprenylphosphoryl- β -D-ribose 2'oxidase	0.9
Rv3577	MSMEG_6071	Metallo- β -lactamase	0.9
Rv0129c	MSMEG_3580	FbpB mycolyl transferase	0.6
Rv3782	MSMEG_6367	Galactofuranosyl transferase	0.8
Rv3808c	MSMEG_6403	Galactofuranosyl transferase	0.8
Rv3265	MSMEG_1826	dTDP-RhA:a-D-GlcNAc-diphosphoryl polyprenol, a-3-L-rhamnosyl transferase	0.7
Rv0906	MSMEG_5638	Metallo- β -lactamase	0.6

Rv2518c	MSMEG_4745	LdtB L,D -transpeptidase	0.6
Rv0237	MSMEG_0361	Glycosyl hydrolase	0.6
Rv3330	MSMEG_1661	DacB1D-alanyl-D-alanine carboxypeptidase	0.5
Rv2748c	MSMEG_2690	DNA translocase FtsK	0.5
Rv2171	MSMEG_4239	Conserved lipoprotein	-0.8
Rv2721c	MSMEG_2739	Transmembrane alanine and glycine rich protein	-0.5




Figure 1. Schematic representation of PknB constructs and strains generated in this study. Kinase: N-terminal kinase domain (aa 1-279); TM: transmembrane region (aa 331-354); TM-PASTA: Penicillin And Serine or Threonine kinase Associated domain (aa 331-627); TM-P1 (aa 331-423); MYC-HIS: His-Tag; PASTA (aa 354-627).

146x121mm (300 x 300 DPI)





Figure 2. Effect of *pknB* and *pknB*- Δ PASTA overexpression on growth of *M. smegmatis*. Mycobacteria (5×10^6) from early stationary phase were inoculated in 100 well honeycomb plates, containing various media and incubated in Bioscreen Growth analyser at 37°C with shaking for 5 days in the presence of tetracycline. Presented data are mean values of 5 replicates from 2 independent experiments. Standard deviations were 10% or less of average values and are not shown for clarity.
196x330mm (300 x 300 DPI)



177x175mm (300 x 300 DPI)




Figure 4. Overexpression of PknB_PASTA results in alteration of bacterial morphology and vancomycin labelling. Mycobacteria from logarithmic growth phase were labeled with Vancomycin-BODIPY conjugate, washed and mounted in PBS and examined using a Diphot 300 inverted microscope. Three independent fields for each strain are shown. Bars indicate 5 μ m. A- MIND; B - TMP. Left panel - phase contrast, middle panel - fluorescence microscopy, right panel - merged images. Arrows indicated labeled peptidoglycan. 248x485mm (300 x 300 DPI)




Figure 5. Culture supernatant and high concentration of Mg²⁺ abolish the inhibitory effect caused by PknB-PASTA over-expression. (a) Culture supernatants abolish growth inhibitory effect of PknB_{TM}-PASTA over-expression in a dose-dependent manner. Relative concentration of culture supernatant added is expressed as "fold concentration". A value of 1-fold corresponds to undiluted culture supernatant. The apparent lag phase was calculated as a period of time when culture reached OD_{600nm} 0.1. (b) Mg²⁺ (10 mM) relieves growth inhibition of TMP in the manner comparable to undiluted culture supernatant. (c) Effect of low Mg²⁺ supernatant on apparent lag-phase of TMP strain. (d) Effect of synthetic mucopeptides on growth of *M. smegmatis*. Mycobacteria from early stationary phase were washed in Sauton's medium twice and 5×10⁶ bacteria were inoculated in Sauton's medium, containing culture supernatant or synthetic mucopeptides at final concentration of 10 μM. MPP – MurNAc – Ipentapeptide, DPP – GlcNAc-MurNAc-peptapeptide, ADPP – GlcNAc-1,6 anhydromuramyl-pentapeptide, TPP – tetra-saccharide-peptide, SN – 50% (v/v) culture supernatant. GlcNAc, N-acetylglucosamine; MurNAc, N-acetylmuramic acid. pentapeptide L-Ala-γ-D-Glu-m-DAP-D-Ala-D-Ala. (b) and (d) Presented data are mean values of 5 replicates from 2 independent experiments. Standard deviations were 10% or less of average values and are not shown for clarity. SN - media were supplemented with culture supernatant.

144x113mm (300 x 300 DPI)




Figure 6. PknB_PASTA reduces growth of Mycobacterium tuberculosis. (a) Overexpression of PknB_PASTA domain delays growth initiation. Mycobacteria ($\sim 1 \times 10^6$) were inoculated in supplemented 7H9 medium containing hygromycin and tetrocycline. (b) Effect of different concentrations of rPASTA on M. tuberculosis growth. Mycobacteria (10^3 cells/ml) were inoculated in supplemented 7H9 medium containing different concentration of sterile rPASTA. The optical density was measured after 21 days of incubation. (c) Regrowth of stored M. tuberculosis was delayed by the addition of rPASTA. M. tuberculosis (2×10^4 cells/ml) were inoculated in supplemented 7H9 medium. Buffer was used as control and rPASTA was added at final concentration of 10 µg/ml. B, C Each data point is an average value of 3 biological replicates, error bars indicate standard deviations. Experiments were performed three times and results of one typical experiment are shown.

136x108mm (300 x 300 DPI)




Figure 7. Schematic representation of possible mechanisms of PknB_PASTA mediated growth inhibition. (a) In the TMP strain the overexpressed PknB_PASTA competes with native PknB for peptidoglycan ligands and interferes with PknB-mediated signalling. (b) Overexpressed PknB_PASTA binds penicillin-binding proteins and dysregulates peptidoglycan polymerization. (c) Overexpressed PknB_PASTA binds to peptidoglycan and destabilizes its structure. PGS – peptidoglycan-synthesis complex.

183x435mm (300 x 300 DPI)