

No evidence for an evolutionary trade-off between learning and immunity in a social insect

A. Alghamdi¹, N.E. Raine², E. Rosato³ and E.B. Mallon^{1*}

1. Biology Department, University of Leicester, LE1 7RH, UK

2. Research Centre for Psychology, School of Biological and Chemical Sciences, Queen Mary, University of London, E1 4NS, UK

3. Genetics Department, University of Leicester

The first two authors contributed equally to this work.

* ebm3@le.ac.uk, 01162523488, Fax: 01162523330

Summary

The immune response affects learning and memory in insects. Given this and the known fitness costs of both the immune system and learning, does an evolutionary trade-off exist between these two systems? We tested this by measuring the learning ability of twelve bumble-bee (*Bombus terrestris*) colonies in a free-flying paradigm. We then tested their immune response using the zone of inhibition assay. We found a positive relationship between colony learning performance and immune response, that is, fast learning colonies also show high levels of antimicrobial activity. We conclude that there is no *a priori* reason to demand an evolutionary relationship between two traits that are linked physiologically.

Keywords

Psychoneuroimmunology, cross-talk, social insects, learning speed.

Short title

No evolutionary trade-off between immunity and learning

Introduction

There is extensive communication between the nervous system and immune system in mammals (Dantzer, 2004). Many responses to parasites, such as fever, increased slow-wave sleep, reduced activity, exploration and sexual behaviour in mammals are orchestrated by immune products (proinflammatory cytokines) released in response to the detection of antigens (Maier & Watkins, 1998). Links between nervous and immune systems are not unique to vertebrates. We have shown that both honeybees *Apis mellifera* (Mallon et al., 2003) and bumble-bees *Bombus terrestris* (Riddell & Mallon, 2006, Alghamdi et al., 2008) perform poorly in learning assays when their immune systems have been challenged by lipopolysaccharide (LPS). LPS is a component of gram-negative bacterial cell walls which is a non-pathogenic elicitor of the immune response (Moret & Schmid-Hempel, 2000). That is, we found that learning and memory are impaired by the immune response directly with no parasite present.

Given this physiological link between learning and immunity, and that learning and immunity have demonstrated fitness costs in insects (Raine & Chittka, 2008, Mery & Kawecki, 2003, Kraaijeveld & Godfray, 1997), it seems reasonable to hypothesise an evolutionary trade-off between learning and immunity. An evolutionary trade-off is where the evolution of an increase in a given trait leads to a reduction in a different trait. This could be due to the pleiotropic effects of the genes involved, linkage disequilibrium with deleterious mutations or resource allocation during development (Schmid-Hempel, 2005). Whilst a recent paper showed no evidence of a trade-off between immunity and learning in different artificially selected *Drosophila* lines in

the laboratory (Kolss et al., 2006), it might be more ecological relevant to examine natural levels of variation in these traits.

Bumble-bees are an obvious candidate for this approach as colonies show natural variation in learning performance (Raine et al., 2006b) and the physiological relationship between learning and immunity has been demonstrated in *B. terrestris* (Riddell & Mallon, 2006, Alghamdi et al., 2008). Also there are ecological reasons to believe that learning and immunity could be more costly in bumble-bees than *Drosophila*, potentially leading to a higher likelihood of a trade-off.

The demands of foraging from many different flower species, which can vary dramatically in the quantity and quality of rewards they offer, and the need to find the nest after each foraging bout, mean that bees have highly developed cognitive abilities. Bumble-bees also learn from conspecifics, so-called social learning (Leadbeater & Chittka, 2007). Furthermore, we would also expect that immunity would be a more important trait in social species that have high contact rates with genetically close individuals leading to a greater chance of infection (Cremer et al., 2007).

In this study we used a free-flying floral choice assay to test the learning abilities of bumble-bee colonies. We took workers from these colonies and tested their immune response using the antibacterial zone of inhibition assay. This allowed us to identify any evolutionary relationship between these two traits.

Materials and Methods

We obtained twelve bumble-bee (*Bombus terrestris dalmatinus*) colonies from Koppert Biological Systems (Berkel en Rodenrijs, The Netherlands). All workers were uniquely marked with Opalith tags (Christian Graze KG, Germany).

Learning assay

Results from this associative learning assay are reported in a previous paper (Raine & Chittka, 2008). Bees were pre-trained to forage from 20 bicoloured, blue and yellow, artificial flowers in a flight arena. During pre-training all flowers were rewarded with 50% (w/w) sucrose solution providing previously colour-naïve bees with an equal chance to associate both colours with reward (Raine et al., 2006b). Bees completing at least 5 consecutive foraging bouts were selected for training. These foragers were trained individually, in a flight arena containing 10 blue (Perspex® Blue 727) and 10 yellow (Perspex® Yellow 260) artificial flowers (each 24 x 24mm). Yellow flowers were rewarding (each contained 15µl of 50% (w/w) sucrose solution), whilst blue flowers were empty (unrewarding). Bees were regarded as choosing a flower when they either approached (inspected), or landed on it. Choosing a yellow flower was regarded as ‘correct’, whilst choosing a blue flower was deemed to be an ‘error’. We recorded the choice sequence made by each bee from the time it first entered the flight arena. Recording flower choices ceased once a bee made 99 flower choices after the first time it probed a rewarding (yellow) flower (Raine et al., 2006b). Flowers were changed and their positions re-randomized between foraging bouts to prevent bees using scent marks or previous flower positions as predictors of reward. Flower colours were selected so that bees had to overcome their strong, unlearned preference for blue, before associating one of their innately least favoured colours (yellow) with

reward (Raine et al., 2006a, Chittka et al., 2004).

Learning curves

The starting point for each bee's learning curve was the proportion of errors made (blue flowers chosen) before the bee first probed a rewarding (yellow) flower. Flower choices made by each bee after (and including) the first time it probed a rewarding (yellow) flower were evaluated as the number of errors (blue flowers chosen) in each group of 10 choices. Learning curves (exponential decay functions: $y = y_0 + Ae^{-x/t}$) were fitted to these eleven data points (i.e. the start pointing and subsequent 10 groups of ten flower choices) for each individual bee, using Microcal Origin (Chittka *et al.* 2004; Raine *et al.* 2006b). Here, x is the number of flower choices the bee made, starting with the first time it probed a yellow flower, and y is the number of errors. The saturation performance level (y_0) is the number of errors made by a bee after finishing the learning process, i.e. when reaching a performance plateau. The decay constant (t) is a measure of learning speed: with lower t values corresponding to faster learning speeds. A is the curve amplitude: the maximum displacement (height) of the curve above y_0 . Both amplitude (A) and saturation performance (y_0) were constrained between 0-10 for curve fitting.

Zone of inhibition assay

This assay measures antibacterial activity: it is based on the ability of immune proteins to inhibit bacterial growth when placed onto an agar plate seeded with bacteria (*Arthrobacter globiformis* 10^5 bacteria per ml of agar). Workers from all twelve colonies were sacrificed after the learning assay and stored at -20°C for later analysis. Each thorax was homogenised in 300 μL of sodium cacodylate solution. 2

μL of the supernatant from the centrifuged solution (1300g for 10 minutes at 4°C) was pipetted into a hole on the agar plate. This was incubated over night (28°C). The resultant zones of inhibition (mm) were measured as the mean of its longest and shortest axis (ZOI value).

Results

As reported in Raine & Chittka (2008) there was significant variation among colonies in learning speed (t value: one way ANOVA: $F_{11, 160} = 1.900$, $p = 0.043$).

We tested the immune response of 55 bees from 12 colonies (mean number per colony (\pm S.D.) = 4.58 ± 0.67) using the zone of inhibition assay. There was a significant difference between colonies in their immune response ($F_{11,43} = 2.30$, $p = 0.026$), which could not be attributed to the effect of body size (headwidth $F_{1,43} = 0.73$, $p = 0.397$).

There was a significant negative correlation between the median t value of a colony and its mean ZOI value (Spearman's rank: $r = -0.608$, $n = 12$, $p = 0.036$; Figure 1). As high t values correspond to slower learning speeds, this is a positive relationship between the ability of a colony to learn and the strength of its immune response.

Discussion

We found a positive correlation between the ability of a colony's workers to learn and the strength of their immune response. Our initial hypothesis that learning ability and immune response would be in an evolutionary trade-off was not supported. Our result is in broad agreement with that of Kolss *et al.*'s (2006) artificial selection experiment, and expands our understanding of this potential learning and immunity trade-off by examining natural variation in both traits in unmanipulated organisms.

Foraging activity has been shown to decrease the immune response of bumble-bee workers (König & Schmid-Hempel, 1995). All workers tested in the learning assay and subsequently used for ZOI assays had similar levels of foraging experience in the laboratory flight arena. Hence, as all our bees were foragers this could not explain variation in immune response. Potential exposure to pathogens which could induce stimulation of the immune system was identical for all twelve colonies which came directly from the bee breeder and were not exposed to field foraging conditions before this experiment.

Phenotypic correlations are generally seen as weak evidence for evolutionary trade-offs (Reznick *et al.*, 2000). However along with Kolss *et al.*'s selection experiment, we can ask why is there no evidence for an evolutionary trade-off when a physiological connection has been found repeatedly? Below we discuss three mutually nonexclusive possibilities.

Firstly, it could be argued that we have incorrectly generalised antimicrobial response to some measure of overall immunocompetence. The various parts of the insect

immune system (antimicrobial peptides, encapsulation, NO production, etc.) are known not to necessarily correlate (Mallon *et al.* 2003). Hence, other parts of the immune response may show an evolutionary trade-off with memory if tested. However as the physiological trade-off has been found repeatedly with ZOI measures, we felt this was the most likely place to find an evolutionary trade-off.

Secondly, there is the possibility that genetic variation exists not only in resource allocation but also in resource acquisition (Reznick *et al.*, 2000). If there was more variation in allocation and less in acquisition, we would expect to see a negative correlation between any two life history traits. Vice versa, we would expect to see a positive correlation (Van Noordwijk & Dejong, 1986).

Thirdly, Schmid-Hempel outlined the differences between the evolutionary and the activation cost of the immune system (Schmid-Hempel, 2005). Evolutionary costs are the fitness effects of possessing an immune system of a given strength. Evolutionary costs can occur due to the pleiotropic effects of resistance genes, linkage disequilibrium with deleterious alleles or changes in resource allocation during development. Activation costs are simply the effect on other physiological systems of generating the immune response from an organism's immune system. Although a physiological connection may lead us to look for an evolutionary trade-off there is no *a priori* reason to demand one.

Acknowledgements

We thank Oscar Ramos-Rodriguez for help with data collection, and Lars Chittka for useful comments. AA is funded by a Saudi government scholarship, and this work was supported by NERC grant (NER/A/S/2003/00469) to NER.

Figures

Figure 1. The relationship between median learning speed (negative t values) and mean zone of inhibition response of the twelve colonies. As high t values correspond to slower learning speeds we have plotted negative t values to make clear the positive relationship between the colony learning ability and the strength of its immune response. Each point represents a colony. Vertical error bars represent median absolute deviation. Horizontal error bars represent standard error.

References

- Alghamdi, A., Dalton, L., Phillis, A., Rosato, E. & Mallon, E. B. 2008. Immune response impairs learning in free-flying bumble-bees. *Biol. Lett.* **4**: 479-481. (doi:10.1098/rsbl.2008.0331)
- Chittka, L., Ings, T. C. & Raine, N. E. 2004. Chance and adaptation in the evolution of island bumblebee behaviour. *Popul. Ecol.* **46**: 243-251. (doi:10.1007/s10144-004-0180-1)
- Cremer, S., Armitage, S. A. O. & Schmid-Hempel, P. 2007. Social immunity. *Curr. Biol.* **17**: R693-R702. (doi:10.1016/i.cub.2007.06.008)
- Dantzer, R. 2004. Innate immunity at the forefront of psychoneuroimmunology. *Brain Behav. Immun.* **18**: 1-6. (doi:10.1016/j.bbi.2003.09.008)

- Kolss, M., Kraaijeveld, A. R., Mery, F. & Kawecki, T. J. 2006. No trade-off between learning ability and parasitoid resistance in *Drosophila melanogaster*. *J. Evol. Biol.* **19**: 1359-1363. (doi:10.1111/j.1420-9101.2005.01068.x)
- König, C. & Schmid-Hempel, P. 1995. Foraging activity and immunocompetence in workers of the bumble bee *Bombus terrestris* L. *Proc. R. Soc. B* **260**: 225-227.
- Kraaijeveld, A. R. & Godfray, H. C. J. 1997. Trade-off between parasitoid resistance and larval competitive ability in *Drosophila melanogaster*. *Nature* **389**: 278-280.
- Leadbeater, E. & Chittka, L. 2007. Social learning in insects - from miniature brains to consensus building. *Curr. Biol.* **17**: R703-R713. (doi:10.1016/j.cub.2007.06.012)
- Maier, S. F. & Watkins, L. R. 1998. Cytokines for psychologists: implications of bi-directional immune to brain communication for understanding behaviour, mood, and cognition. *Psychol. Rev.* **105**: 83-107.
- Mallon, E. B., Brockmann, A. & Schmid-Hempel, P. 2003. Immune response inhibits associative learning in insects. *Proc. R. Soc. B* **270**: 2471-2473. (doi:10.1098/rspb.2003.2456)
- Mallon, E. B., Loosli, R. & Schmid-Hempel, P. 2003. Specific versus nonspecific immune defense in the bumblebee, *Bombus terrestris* L. *Evolution* **57**: 1444-1447. (doi:10.1554/02-715)
- Mery, F. & Kawecki, T. J. 2003. A fitness cost of learning ability in *Drosophila melanogaster*. *Proc. R. Soc. B* **270**: 2465-2469. (doi:10.1098/rspb.2003.2548)
- Moret, Y. & Schmid-Hempel, P. 2000. Survival for immunity: The price of immune system activation for bumblebee workers. *Science* **290**: 1166-1168. (doi:10.1126/science.290.5494.1166)

- Raine, N. E. & Chittka, L. 2008. The correlation of learning speed and natural foraging success in bumble-bees. *Proc. R. Soc. B* **275**: 803-808. (doi:10.1098/rspb.2007.1652)
- Raine, N. E., Ings, T. C., Dornhaus, A., Saleh, N. & Chittka, L. 2006a. Adaptation, genetic drift, pleiotropy, and history in the evolution of bee foraging behavior. *Adv. Stud Behav* **36**: 305-354. (doi:10.1016/S0065-3454(06)36007-X)
- Raine, N. E., Ings, T. C., Ramos-Rodriguez, O. & Chittka, L. 2006b. Intercolony variation in learning performance of a wild British bumblebee population (Hymenoptera: Apidae: *Bombus terrestris audax*). *Entomologia Generalis* **28**: 241-256.
- Reznick, D., Nunney, L. & Tessier, A. 2000. Big houses, big cars, superfleas and the costs of reproduction. *Trends Ecol. Evol.* **15**: 421-425.
- Riddell, C. E. & Mallon, E. B. 2006. Insect psychoneuroimmunology: Immune response reduces learning in protein starved bumblebees (*Bombus terrestris*). *Brain Behav. Immun.* **20**: 135-138. (doi:10.1016/j.bbi.2005.06.008)
- Schmid-Hempel, P. 2005. Evolutionary ecology of insect immune defenses. *Ann. Rev. Entomol.* **50**: 529-551. (doi:10.1146/annurev.ento.50.071803.130420)
- Van Noordwijk, A. J. & Dejong, G. 1986. Acquisition and allocation of resources - their influence on variation in life-history tactics. *Am. Nat.* **128**: 137-142.

