

**The three-spined stickleback - *Schistocephalus solidus* system:
an experimental model for investigating host-parasite interactions in fish**

I. Barber^{1*} & J. P. Scharsack²

¹ *Department of Biology, University of Leicester, University Road, Leicester LE1 7RH, UK*

² *Animal Evolutionary Ecology Group, Institute for Evolution and Biodiversity, University of
Münster, Huefferstr. 1, D-48149 Münster, GERMANY*

*Corresponding author, email: ib50@le.ac.uk, phone: +44(0)116 252 3462, fax: +44(0)116
252 3330

SUMMARY

Plerocercoids of the pseudophyllidean cestode *Schistocephalus solidus* infect three-spined stickleback *Gasterosteus aculeatus*, with important consequences for the biology of host fish. Techniques for culturing the parasite *in vitro* and generating infective stages that can be used to experimentally infect sticklebacks have been developed, and the system is increasingly used as a laboratory model for investigating aspects of host-parasite interactions. Recent experimental laboratory studies have focused on the immune responses of hosts to infection, the consequences of infection for the growth and reproductive development of host fish and the effects of infection on host behaviour. Here we introduce the host and the parasite, review the major findings of these recent experimental infection studies and identify further aspects of host parasite interactions that might be investigated using the system.

Key words: threespine, Cestoda, Pseudophyllidea, laboratory model, growth, immunology, *Ligula intestinalis*, behaviour, fitness

INTRODUCTION

Plerocercoids of the pseudophyllidean cestode *Schistocephalus solidus* are common parasites of three-spined sticklebacks *Gasterosteus aculeatus* in freshwater and brackish habitats throughout the geographical range of the fish. The three-spined stickleback-*S. solidus* host-parasite system has become an important model in experimental parasitology and is increasingly used to investigate a wide range of questions about host-parasite interactions and co-evolution. Here we present a review of recent studies that have used controlled experimental infections to investigate host-parasite interactions in this system.

We begin our review with background information on the parasite's lifecycle, on the host fish and the 'typical' phenotype of infected sticklebacks in nature, and briefly discuss emerging variation in infection phenotype. We then examine how aspects of the life cycle can be experimentally manipulated in the lab to allow experimental infections of sticklebacks to be undertaken. The remainder of our review focuses on how experimental infection studies have been used to illuminate host-parasite interactions in the stickleback-*S. solidus* system, including the immune responses of the fish host, the energetic consequences of infection and the consequences of infections for fish behaviour and fitness.

Life cycle of S. solidus in nature

Schistocephalus solidus is a trophically transmitted pseudophyllidean cestode with a three-host life cycle. The definitive host can be any warm-blooded vertebrate; most typically these are fish-eating birds though other endotherms can harbour adult worms, including otters (Hoberg *et al.*, 1997) and – though presumably only rarely – humans (Coombs and Crompton, 1991). *Schistocephalus solidus* does not grow in the gut of the definitive host but undergoes the final stages of sexual maturation there, reproducing sexually either by selfing (if singly infected) or by cross-fertilization (in multiple infections). Eggs released into the water with

the bird's faeces hatch to produce free-swimming coracidia that are transmitted trophically to a wide range of cyclopoid copepods, the 1st intermediate hosts. Here the parasites develop in the copepod haemocoel into proceroids, becoming infective to three-spined sticklebacks, the obligatory specific 2nd intermediate hosts (Bråten, 1966), with the formation of a hooked cercomer. Sticklebacks acquire infections when they feed on parasitized copepods, and in the stickleback digestive tract infective proceroids shed their outer layer, together with the cercomer, and penetrate the wall of the intestine. The parasite then develops into a plerocercoid, which grows to a large size in the fish host's body cavity. The life cycle is completed when sticklebacks harbouring infective plerocercoids are ingested by a definitive host (Clarke, 1954).

The geographical distribution of the parasite is limited by the distribution of the only obligate host in the life cycle, the three-spined stickleback, which is restricted to the Northern hemisphere and occurs around the margins of the Atlantic and Pacific Oceans (Bell and Foster, 1994). In this geographical region *S. solidus* is a regular parasite of three-spined stickleback populations inhabiting freshwater and brackish ecosystems, and is most common found in those in lacustrine or slow flowing habitats (Kennedy, 1974; Wootton, 1976; Barber, 2007).

Specificity of stickleback host

Three-spined sticklebacks are the only recognised fish host of *S. solidus*, although other *Schistocephalus* spp. infect nine-spined stickleback *Pungitius pungitius* and sculpins (Hyslop and Chubb, 1983; Chubb *et al.*, 2006; Seppala *et al.*, 2007; French and Muzzall, 2008). Experimental exposure of nine-spined sticklebacks (*Pungitius pungitius*) to infective stages of *S. solidus* led to much slower plerocercoid growth and infections were cleared after 14 days, while plerocercoids kept on growing in three-spined sticklebacks (Orr *et al.*, 1969).

Heterotransplants of *S. solidus* plerocercoids from *G. aculeatus* to other species of fish (*Cottus gobio*, *Nemacheilus barbatula*, *Phoxinus phoxinus*, *Salmo trutta*, *Coregonus clupeoides*, *Perca fluviatilis*, *Rutilus rutilus*, *Esox lucius*), including *P. pungitius*, died within 2-10 days after transfer, while homotransplants between *G. aculeatus* survived (Bråten 1966). These observations indicate that in principle *S. solidus* plerocercoids can be cleared by a fish immune system, but obviously *S. solidus* plerocercoids are able to avoid an effective immune response in *G. aculeatus*, their specific second intermediate host.

Sticklebacks as experimental model hosts

A major attraction of the stickleback-*S. solidus* host-parasite system for ecological and evolutionary biologists is the rich history of studies investigating the natural history, behaviour and evolutionary biology of the host fish, and a correspondingly substantial literature that has been regularly and thoroughly reviewed (Wootton, 1976; Wootton, 1984; Bell and Foster, 1994; Östlund -Nilsson *et al.*, 2006). This background permits a wide range of ecologically and evolutionarily relevant questions to be addressed. Furthermore, the three-spined stickleback has, in recent years, assumed even more importance as a model species in biology; the publication of linkage and chromosome maps (Peichel *et al.*, 2001; Kingsley *et al.*, 2004) and the sequencing of its genome (Kingsley, 2003) has greatly enhanced its utility in molecular studies of evolution and development (McKinnon *et al.*, 2004; Shapiro *et al.*, 2004; Colosimo *et al.*, 2005; Gibson, 2005; Shapiro *et al.*, 2006). Sticklebacks are also readily bred in laboratory aquaria (Barber and Arnott, 2000), facilitating the challenge of naïve individuals and thus fulfilling both scientific and local ethical requirements for experimental infection studies.

*Field studies of *S. solidus* infected sticklebacks*

A number of studies have examined *S. solidus* infection prevalence and intensity in natural stickleback populations, and the phenotype (including the appearance, energetic condition, reproductive capacity and behaviour) of naturally infected sticklebacks has been well documented. Observations on the phenotype of infected fish from populations where *S. solidus* is endemic are summarised in Table 1. The proportion of fish harbouring infections can be extremely high, in some cases approaching 100% (Dick, 1816; Smyth, 1947; Hopkins and Smyth, 1951), but this varies considerably between populations (MacColl, 2009), and temporally within them (Chappell, 1969). Typical features of classical ‘schistocephalosis’ include characteristic distension of the fish’s abdomen, an altered swimming gait, increased risk-taking behaviour, reduced body condition and functional (if not physiological) castration. However, as more host populations are studied it is becoming clear that there is significant variation in infection phenotype, and there are a number of exceptions to these ‘typical’ infection phenotypes. Notably, intensive studies of some Alaskan populations suggest that the traditional view of *S. solidus* as an absolute castrator of female sticklebacks may need to be revised (Heins and Baker, 2008), and in a small number of populations infection is additionally associated with almost complete demelanisation (Lobue and Bell, 1993; Ness and Foster, 1999).

In vitro culture of S. solidus

Pioneering work by the parasite physiologist J.D. Smyth in the 1940-50s developed protocols for the *in vitro* culture of adult helminths, including *S. solidus* (Smyth, 1946; Smyth and McManus, 1989; Smyth, 1990). In brief, plerocercoids recovered from infected sticklebacks are removed and placed into sterilised culture vessels containing a buffered medium that provides a suitable physico-chemical environment for parasite development (such as horse serum and / or a cell culture medium such as RPMI-1640). Antibiotics may be added to

reduce contamination. The plerocercoid(s) are not placed directly into the medium but are instead constrained within narrow diameter semi-permeable dialysis tubing, which mimics the small intestine of the definitive host and provides the constriction needed for successful fertilisation (Smyth, 1990). Culture vessels are placed into a water bath set to 40°C and shaken gently to aid the dissolution of metabolic products from developing worms. Due to the progenetic development of pseudophyllidean plerocercoids, sexual maturation occurs rapidly and within 48h the adult worms begin producing eggs. The fact that *S. solidus* attains its final size in the fish body cavity and matures so rapidly to the adult form made it possible for the first time to maintain adult cestodes under sterile experimental conditions and undertake detailed physiological studies. Such studies had been impossible with most other tapeworm species, which typically grow and mature over prolonged time periods in the host intestine, and *S. solidus* became an extremely valuable model in parasite physiology (Smyth and McManus, 1989).

Using Smyth's techniques – or slightly modified versions of them – large numbers of eggs can be generated and hatched to yield coracidia, which in turn can be used to infect lab-bred copepods (Wedekind, 1997). After a period of development to the infective cercaria-bearing stage, proceroids can be fed to sticklebacks inside infected copepods, either by gavage or by natural feeding, to generate experimentally infected fish hosts and allowing detailed experimental studies of fish-parasite interactions.

MECHANISMS OF RESISTING *S. SOLIDUS* INFECTION: HOST BEHAVIOUR, IMMUNE RESPONSES AND HOST MANIPULATION

Animals have evolved three major types of mechanisms to avoid, or reduce the likelihood of developing, debilitating parasite infections; behavioural mechanisms that limit contact with

infectious agents, physical barriers to invasive stages and immune systems. The ability to experimentally infect intermediate hosts in large numbers means that it is possible to examine the responses of hosts to controlled, experimental challenge. In recent years, experimental infection studies have been used to examine both behavioural and immunological aspects of stickleback responses to infective stages of *S. solidus*.

Behavioural resistance

The strong selection pressure placed on host organisms to avoid debilitating parasite infections, together with the typically high costs of mounting immune responses against invading pathogens, has led to the evolution of a wide range of strategies of behavioural resistance in animals (Hart, 1990; Hart, 1992; Hart, 1997). For parasites that are transmitted trophically between hosts, avoiding infected prey intuitively reduces the level of exposure to infection, but this requires that infected prey are identifiable, and that the benefits of avoiding parasitized prey outweigh the costs of ignoring them. In some cases, feeding on parasitized prey that are easy to catch may benefit potential hosts if the risks of becoming infected, and/or the costs of infection, are low (Lafferty, 1992).

Copepods infected with procercoids of *S. solidus* and other related pseudophyllidean cestodes behave differently to those that are non-infected, providing the potential for avoidance by discriminating fish. However, the behaviours that are altered by infection – which include activity patterns, swimming ability and responses to disturbance – make infected copepods more susceptible to human ‘predators’ armed with pipettes (Pasternak *et al.*, 1995; Urdal *et al.*, 1995; Wedekind and Milinski, 1996) and so potentially make them more visible to, and / or more easily caught by, fish predators. Infected copepods have also been reported to actively approach sticklebacks (Jakobsen and Wedekind, 1998). It is therefore likely that behaviour changes in infected copepods are adaptations of the parasite to facilitate

transmission (Parker *et al.*, 2009), and recent evidence examining temporal aspects of their behaviour change supports this (Hammerschmidt *et al.*, 2009; Hammerschmidt and Kurtz, 2009). So do sticklebacks avoid eating infected copepods? The results of two critical experimental tests suggest that sticklebacks have no behavioural defence against *S. solidus*, and may even feed on infected copepods preferentially (Urdal *et al.*, 1995; Wedekind and Milinski, 1996). However, there is still further work to be done in this area. One possibility for the apparent non-evolution of avoidance behaviour is that sticklebacks are simply unable to discriminate infected from non-infected copepods, so avoiding infection would mean excluding an important prey type, which is just too costly. Also, because few tests have been carried out, it is not known whether all populations are equally non-discriminating, or whether some populations have evolved to be more selective in their prey choice. Further, individual sticklebacks are known to vary in key personality traits (Bell and Stamps, 2004; Bell, 2005) so are all individuals within populations as likely as others to approach infected copepods, or do fish exhibiting particular personality types suffer increased exposure? Because the probability of acquiring infections after feeding appears to be relatively high (at least in laboratory studies, see below), and the consequences of infections are typically severe, it seems unlikely that feeding on more easily caught infected copepods could be beneficial to sticklebacks. However, it is possible that wild fish develop better immune responses than those reared under sterile laboratory conditions and face a lower risk of infection per infective stage ingested, reducing the pressure to evolve discrimination (see also Hammerschmidt and Kurtz, 2009).

The presence of debilitating parasites in the environment can also drive the evolution of mate preferences, either by the avoidance of mates harbouring directly transmissible parasites, or by the selection of individuals with genes that confer parasite resistance on offspring (Keymer and Read, 1991; Andersson, 1994). In sticklebacks, preferences for males with the brightest red nuptial colouration have evolved at least in part as a mechanism for avoiding parasitised

males (Milinski and Bakker, 1990; Bakker and Milinski, 1991). To test the hypothesis that females selecting brightly ornamented males gain indirect genetic benefits by producing more resistant offspring, Barber *et al.* (2001) produced clutches of maternal half-sibling fish that differed only the brightness of the male parent, and exposed them to infective *S. solidus* proceroids. The results showed that male brightness significantly affected the proportion of half-sibships that developed infections, with brighter males producing the most resistant offspring, suggesting indirect benefits of ornamentation based mate choice.

Dynamics of S. solidus transmission from copepods to sticklebacks

Interactions of *S. solidus* with its first intermediate copepod host were recently reviewed in detail by Hammerschmidt and Kurtz (2009). With no strong evidence that sticklebacks are capable of adopting behaviours to avoid ingesting infected copepods, those in populations with endemic infection are likely to be exposed regularly to infective *S. solidus* proceroids. After the ingestion of a *S. solidus* infected copepod the prey is digested in the stomach and proceroids are released from copepod tissues. Proceroids retain their outer layer – which is rich in PNA-binding sugars (GalNac, D-galactose) – in the stickleback stomach, and presumably this protects the parasite from enzymatic digestion (Hammerschmidt and Kurtz, 2007). During passage through the stomach, the outer layer is shed (or digested) together with the cercomer and in the intestine the underlying tegument with microtriches is exposed. Typically for vertebrate cell surfaces this cuticle is rich in sialic acid residues, which may help the parasite evade the stickleback's immune system (Hammerschmidt and Kurtz, 2005). From the intestine, the parasites penetrate the gut wall and enter the body cavity. The period from ingestion to establishment in the stickleback body cavity is critically important in determining infection success, with 50-75% of ingested parasites failing to complete this phase (Hammerschmidt and Kurtz, 2007). In an experimental infection

experiment, the majority (>90%) of parasites recovered from sticklebacks were alive in the stomach 16h post-exposure (p-e). After 22h p-e, approximately 40% were still alive in the stomach, 40% had entered the body cavity and about 20% of parasites recovered (by dissection and histological analysis) were dead. By 24h p-e, the majority (>90%) of detectable parasites were alive in the body cavity whereas dead or damaged parasites were no longer detectable, presumably due to progressing degradation in the digestive tract (Hammerschmidt and Kurtz, 2007). This indicates that parasites are vulnerable to the aggressive environment in the digestive tract after losing their outer layer, and progression to the intestine and passage through the gut wall must be achieved quickly to avoid significant losses in viability. In terms of preventing *S. solidus* establishment, the relative contribution of hostile conditions in the fish digestive tract and the host immune system is difficult to estimate; however, no attachment of phagocytic cells to, or encapsulation of, *S. solidus* stages during gut wall penetration or in the body cavity was observed (Hammerschmidt and Kurtz, 2007).

There is little evidence yet for a prominent reduction of *S. solidus* viability once the body cavity is reached (i.e. clearance by the immune system). In another infection experiment, relatively few dead parasites (n = 4) were found in the body cavity of infected sticklebacks at 7d and 17d p-e, while 78 *S. solidus* plerocercoids were recovered alive after these times (Scharsack *et al.*, 2007). During this experiment, infection rates changed from >60% at 7d and 17d p-e to approximately 50% after 27-67d p-e (Scharsack *et al.*, 2007). This suggests that while clearance of *S. solidus* plerocercoids in the body cavity is possible during early stages of infection it occurs less frequently later on.

Stimulation of immunity soon after *S. solidus* infection seems to reduce the infection success of *S. solidus*. Wedekind & Little (2004) triggered activation of the host immune system by tissues injury through spine clipping at 7d p-e to *S. solidus*. At 90d p-e, the spine-clipped sticklebacks showed significantly lower infection rates compared to controls without spine

clipping (Wedekind and Little, 2004). At which time plerocercoids were cleared was not recorded in this study, but the results might indicate that immune stimulation was most efficient in the early (1-2w) stage of infection, when *S. solidus* is still vulnerable to immune attack.

Lymphatic organs and leukocytes during S. solidus infection Responses from different immunologically active organs have been recorded in sticklebacks infected experimentally with *S. solidus*. The spleens of *S. solidus* infected sticklebacks were enlarged compared to non infected fish (Arnott *et al.*, 2000). Enlargement of the spleen is often observed during parasite infections of fish, e.g. in common carp *Cyprinus carpio* infected with the blood flagellate *Trypanoplasma borreli*, due to proliferation of leukocytes and increased amounts of antigen-antibody immune complexes, which are removed from the blood stream by spleen macrophages (Bunnajirakul *et al.*, 2000). In the stickleback-*S. solidus* system, specific reasons for the enlargement of spleens are to date unclear and await further investigation. In the blood of *S. solidus* infected sticklebacks, distinct changes of leukocyte subsets have been observed. Early in the infection the proportion of granulocytes increased, while the proportion of lymphocytes decreased in the peripheral blood, with both trends levelling out after 60-96d p-e (Scharsack *et al.*, 2004). The offspring of more brightly ornamented male sticklebacks showed elevated white blood cell counts and were less susceptible to *S. solidus* infection (Barber *et al.*, 2001). These observations suggest that peripheral blood leukocytes may indeed play a role in the *S. solidus* infection, but the underlying mechanisms are not yet well understood. Most information about leukocyte responses to *S. solidus* has been generated from experiments with stickleback head kidney leukocytes (see later). To date, limited information is available on the interplay of lymphatic tissues during *S. solidus* infection, in particular the role of (cellular) immune defence at the site of infection, the body cavity.

However, because the teleost head kidney is a site of antigen presentation, leukocyte activation, proliferation and maturation - and consequently interacts closely with immunological activity in the periphery (Manning, 1994; VanMuiswinkel, 1995) - information derived from head kidney leukocyte studies can be regarded as representative for immune activity in the periphery, even if specific interactions at the site of infection might remain concealed.

Cellular innate immunity

Respiratory burst and monocyte proliferation The respiratory burst activity of head kidney leukocytes (HKL) is one of the most important mechanisms of cellular innate immunity, so may be expected to be up-regulated at an early stage in exposed sticklebacks. However, the HKL respiratory burst from *S. solidus* exposed sticklebacks 7-37d after an experimental challenge did not differ from that of sham-exposed controls, suggesting the mechanism is not important in early defence against infection (Scharsack *et al.*, 2007). Interestingly, the respiratory burst of HKLs was up-regulated from 47-67d p-e, but as neither the survival nor the growth rates of *S. solidus* plerocercoids were affected during this period this appears to be an ineffective defence mechanism.

Nevertheless, the proliferation of head kidney monocytes – a component of the cellular innate immune response – was up-regulated among exposed sticklebacks at 7d p-e, suggesting that the mobilisation of monocytes could play a role in early defence against *S. solidus*. Among fish that developed infections, monocyte proliferation dropped then below sham-exposed controls at 17 d p-e, recovered and dropped again, indicating possible immune-manipulation by *S. solidus* (Scharsack *et al.*, 2007) (see later). Interestingly the kinetics of monocyte proliferation in exposed fish that did not develop infections followed a similar pattern, suggesting an early priming of monocyte responses.

303

304 *Monocyte manipulation?* The idea that *S. solidus* is capable of substantially manipulating
305 stickleback monocyte responses is supported by *in vitro* experiments. Monocytic leucocytes
306 (granulocytes and macrophages) isolated from the head kidney of experimentally infected
307 sticklebacks at 45d p-e failed to respond to *S. solidus* antigens *in vitro* (Scharsack *et al.*,
308 2004). This was not a general anergy, as monocytes from the same sticklebacks responded to
309 stimulation with a non-specific antigen (poke weed mitogen, PWM) in a comparable manner
310 as cells from sham-exposed controls (Scharsack *et al.*, 2004). Thus *S. solidus* does not appear
311 to immune-compromise its stickleback host, but is apparently capable of manipulating
312 (evading) immune traits that are specifically directed against parasite antigens.

313

314 *Immune priming and susceptibility* Priming of the immune system by *S. solidus* does not
315 induce resistance in *G. aculeatus*, as super infections are possible by sequential exposures (i.e.
316 there is no ‘vaccination effect’). Experimentally infected nine-spined sticklebacks (*Pungitius*
317 *pungitius*) have been shown to reject *S. solidus* plerocercoids more rapidly after pre-exposure
318 to the parasite (Orr *et al.*, 1969). However, *S. solidus* plerocercoids were not able to survive in
319 *P. pungitius* longer than 14d (Orr *et al.* 1966). Detailed analysis of infections harboured by
320 three-spined sticklebacks that had been sequentially exposed to *S. solidus* showed that
321 plerocercoids from later exposures survived better and grew larger than ‘pioneering’ worms
322 (Jäger and Schjørring, 2006). These results include exposures where only the secondary *S.*
323 *solidus* survived and may be explained by the first invading worm paying higher costs of
324 immune manipulation / priming (Jäger and Schjørring, 2006).

325 These findings, together with the observation that priming of monocytes is detectable in
326 exposed sticklebacks that do not develop infections (Scharsack *et al.*, 2007) and the loss of
327 responsiveness of monocytes to (secondary) *in vitro* exposure to *S. solidus* antigens

(Scharsack *et al.*, 2004), suggest that *S. solidus* has a strong impact on the stickleback immune system. Immune priming, initiated to protect the invading parasite from host immune attack, seems to be so efficient that it persists even if the first invader is cleared, facilitating the establishment of subsequent infections.

To what extent immune priming by *S. solidus* can influence susceptibility of sticklebacks to other parasites has not yet been experimentally investigated. In populations with endemic *S. solidus* infection, fish harbouring plerocercoids tend to be more heavily infected by *Gyrodactylus* sp. parasites than those free from *S. solidus* (M. Kalbe, personal communication). Experimental exposure of *S. solidus* infected sticklebacks to other parasites could reveal the extent to which *S. solidus* can influence susceptibility to additional parasites.

Adaptive immunity

Clearance of *S. solidus* infection seems to depend mainly on an early innate immune response, potentially facilitated by previous exposure of the parasite to the aggressive environment in the digestive tract. An adaptive immune response - including the presence of specific antibodies - would need about 2-3 weeks to be fully in place in fish maintained at 18°C (Rijkers *et al.*, 1980). Thus substantial involvement of antibody-mediated immunity in early defence against invading *S. solidus* is unlikely, and since clearance of infection at later stages (beyond 17d p-e) was not observed, antibody-mediated responses to *S. solidus* infection are not expected to make a significant contribution. However, due to the lack of specific tools such responses have not yet been fully investigated.

Lymphocyte activation Nevertheless, in a kinetics study of immune parameters following exposure to *S. solidus*, the proliferation of lymphocytes (B- and T-cells) was measured in head kidney isolates. The clonal expansion and proliferation of lymphocytes forms a

significant component of the adaptive immune response and is expected 1-4 weeks after infection. In *S. solidus* infected sticklebacks, significant changes in lymphocyte proliferation, compared to sham exposed controls, were only observed among exposed fish that did not develop infections. Among these fish, lymphocyte proliferation was elevated at 7d p-e, dropped below controls at 17d p-e before returning to control values from 27 to 67d p-e (Scharsack *et al.*, 2004). (A less prominent and statistically non-significant pattern was recorded among sticklebacks that developed infections). The pattern of lymphocyte proliferation among exposed sticklebacks that did not become infected suggests a possible role in defence against *S. solidus*. Since lymphocyte proliferation dipped below controls at 17d p-e, B-cell proliferation and production of antibodies (T-helper cell 2 - Th2 mediated humoral immunity) is unlikely. Early lymphocyte proliferation might alternatively be explained by the proliferation of T-cells, maintaining a Th1 response that activates cellular immunity. This corresponds to the observation that monocyte proliferation was regulated contemporarily (see Scharsack *et al.* 2007).

Potential role of the Th1-Th2 system? In mammals, helminth parasites are considered a classical inducer of Th2 responses, which have the potential to damage parasites and clear infections (Maizels and Yazdanbakhsh, 2003; Wang *et al.*, 2008). However, the nature of interactions between helminth parasites and the Th1-Th2 system remains controversial (Maizels and Yazdanbakhsh, 2003); for example, schistosomes appear to have evolved immune evasion strategies in which the Th1-Th2 system is driven towards a Th1 response, thereby avoiding humoral responses (Herve *et al.*, 2003).

Information on Th1-Th2 mediated immune function in teleost fish is scarce, but molecular studies indicate that the Th1-Th2 system is at least present (Takizawa *et al.*, 2008a; Takizawa *et al.*, 2008b). The available information does not point towards a typical Th2 response in *S.*

solidus infected stickleback, as lymphocyte proliferation after an initial weak increase remained unaffected (Scharsack *et al.*, 2007) and degenerative changes at the surface of procercoids (as a result of a Th2 induced humoral response) were not detected by means of electron microscopy (Orr *et al.*, 1969). The extent to which protection against *S. solidus* infections in stickleback hosts might involve a Th1 response (or an abrogated Th2 response) therefore requires further investigation.

The MHC and influences on plerocercoid growth Overall, adaptive immunity seems unlikely to protect three-spined sticklebacks from *S. solidus* infection, but there is evidence that it can restrict parasite growth during ongoing infection. Proteins of the major histocompatibility complex (MHC) play a central role in presenting antigens to the adaptive immune system. Using three-spined sticklebacks that varied in their individual MHC class IIB allelic diversity, Kurtz *et al.* (2004) observed that *S. solidus* grew larger in sticklebacks with low and high MHC diversity compared with those having an intermediate number of MHC alleles. The underlying molecular mechanism is unknown, but these results support observations that sticklebacks with intermediate (optimal) MHC IIB diversity suffered less from parasite infections compared to fish with high and low (suboptimal) MHC IIB diversity (Wegner *et al.*, 2003).

Summary: innate and adaptive immunity

Clearance of *S. solidus* by the immune system of its specific second intermediate host, the three-spined stickleback, appears to be the exception rather than the rule. Damage to the parasite by the aggressive gut environment might reduce the infection success at least as prominently as attack by the immune system. *S. solidus* does not appear to be very vulnerable to immune attack, but rather appears to be capable of substantial immune evasion and

manipulation. The typically rapid death of plerocercoids following experimental transfer to fish species other than three-spined sticklebacks (Bråten, 1966; Orr *et al.*, 1969) strongly suggests that fish immune systems can, in principle, clear *S. solidus* infections. It therefore seems most likely that specific adaptation of *S. solidus* to the immune system of the three-spined stickleback permits its invulnerability to host immune responses. From an evolutionary perspective, adaptation to a host immune system is costly and balancing selection on *S. solidus* has resulted in an extremely high degree of specialisation towards a three-spined stickleback host.

The results on immune responses of stickleback against *S. solidus* described here are mainly derived from laboratory experiments. In the wild, *S. solidus* infection success and development in sticklebacks might be constrained by factors acting on the host immune system, such as activation of the immune system by pre-exposure to other parasites, and by other environmental stressors, both natural and anthropogenic.

IMPACTS OF EXPERIMENTAL *S. SOLIDUS* INFECTIONS ON HOST ENERGETICS AND BEHAVIOUR

Schistocephalus solidus infections are expected to impact host energetics and behaviour of host sticklebacks for two main reasons. First, plerocercoids grow to a large size and – because the nutrients to fuel this growth are entirely host-derived – this incurs a considerable energetic burden on host fish (Walkey and Meakins, 1970; Lester, 1971). Second, the parasite relies on the ingestion of the stickleback host to complete its life cycle, facilitating the evolution of parasite adaptations that increase the predation risk of host sticklebacks. A number of studies have quantified infection-associated variation in stickleback energetics and behaviour among fish from naturally infected populations. Experimental infection studies allow a number of fitness correlates to be measured under standardised conditions.

428

429 *Effects of experimental infections on host energetics, growth and sexual development*

430 Laboratory investigations of the impact of *S. solidus* on the growth and development of
431 stickleback hosts have been the subject of a recent review (Barber *et al.*, 2008), so here we
432 provide an overview of the major findings of these studies and outline future approaches and
433 potentially research questions.

434 In naturally infected populations, fish infected with *S. solidus* typically exhibit low growth
435 and poor body condition, and as a result – in most studied populations at least – they suffer
436 reduced sexual development and are unlikely to participate successfully in spawning (Arme
437 and Owen, 1967; Pennycuick, 1971; Tierney *et al.*, 1996; Bagamian *et al.*, 2004; Heins and
438 Baker, 2008). However, when naturally or experimentally-infected fish are maintained under
439 laboratory conditions such effects are less frequently observed, often because ethical guidance
440 on animal husbandry requires fish to experience benign environments, with access to
441 abundant, high quality food. The growth and energetic condition of infected fish can even
442 exceed that of non-infected individuals under certain types of lab housing.

443

444 *Experimental studies of fish held under benign conditions* Barber & Svensson (2003)

445 experimentally exposed lab-bred juvenile sticklebacks to single infective procercoids and held
446 them under a constant host ration of 8% body mass per day. Following, the length of infected
447 sticklebacks followed approximately the same trajectory of non-exposed, control fish over the
448 16-week p-e period. The mass of infected fish (including plerocercoid mass) also followed a
449 similar trajectory to controls, with infected fish actually showing elevated growth rates during
450 weeks 5-7 p-e. However, when the mass contributed by developing plerocercoids was
451 removed, the trajectory of mass increase of infected fish clearly differed from that of controls,
452 and infected fish weighed significantly less at the end of the experiment. On dissection,

infected females were found to have equivalent liver mass, but lower perivisceral fat reserves and, surprisingly, larger ovaries than non-exposed control fish. One explanation for the counterintuitive investment in gonad development is that it may reflect a life history change that could compensate for the likely reduction in survival associated with infection (Minchella, 1985).

Other studies under similarly benign conditions have recorded a similar lack of detectable impact of the parasite on host growth. In a recent infection experiment sticklebacks were fed *ad libitum* with frozen chironomids 3 times a week. Here, the mass of infected stickleback including parasite mass was significantly higher at 57 and 67d p-e compared to controls, but equally high with parasite mass subtracted (Koch, Scharsack, Hammerschmidt, unpublished data). In a study by Arnott *et al.* (2000), experimentally infected fish were held individually and fed *ad libitum* to excess each day. Under these conditions, infected sticklebacks outgrew non-infected fish, weighing significantly more than the latter at the end of the study even when correcting for plerocercoid mass. Infected female fish (though not males) held under these conditions also developed significantly larger livers relative to their body size, and they had an equivalent amount of perivisceral fat to fish that did not develop infections after exposure.

Experimental studies of fish held under more naturalistic conditions The results described above suggest that the feeding regime experienced by hosts has considerable influence on the energetic costs of infection experience by host fish, and hence the phenotype exhibited by *S. solidus* infection. Synthesising the results from a number of lab studies, Barber *et al.* (2008) showed that infection phenotypes more closely reflecting those found in natural populations, were more commonly found when experimentally infected fish are reared under less benign conditions in the lab. For example, when housing exposed and non-exposed sticklebacks

together in groups, effectively forcing competition between infection classes, Barber (2005) found the relative liver mass (hepatosomatic index, HSI) to be significantly reduced among experimentally-infected, compared to sham-exposed, females. Wright *et al* (2007) examined the effect of temporary food restriction on the growth and energetics of sticklebacks experimentally infected with *S. solidus*. In contrast to sham-exposed sticklebacks – which undertook rapid compensatory growth on commencement of *ad libitum* feeding to catch up to the mass of continually-fed fish after only three weeks of re-feeding – experimentally infected sticklebacks showed only partial compensation, reaching just 80% of the mass of continually-fed infected fish after six weeks of re-feeding. Infected fish reared under the compensatory regime also developed smaller livers than sham-exposed ‘compensatory’ fish, whereas infection status did not affect liver size among fish held under a continual feeding regime. Analysis of the food intake of individual fish revealed that the likely cause of the inability of infected fish to compensate was their failure to mount significant hyperphagic responses post-deprivation (Wright *et al.*, 2007). A subsequent study confirmed that the maximum voluntary meal size of infected fish was reduced in infected sticklebacks (Wright *et al.*, 2006). Because fish in natural environments, with temporally unpredictable food availability, are expected to rely heavily on compensatory growth responses, the inability to undertake such responses may exacerbate the growth effects of *S. solidus* and represent a hitherto ‘hidden cost’ of infection.

The goal of laboratory investigations of the stickleback-*S. solidus* system is generally to better understand the selective role that parasites play in nature, so it is becoming increasingly clear that investigating the growth and development effects of *S. solidus* in lab studies presents certain challenges. However, at the same time there is an urgent need to better understand how parasites and hosts interact under altered environments. A possible way in which laboratory studies of the stickleback-*S. solidus* system could contribute considerably to our understanding of host-parasite interactions in nature is to investigate the role of variation in

the rearing environment experienced. To date few studies have systematically investigated such effects, but the effects of factors such as temperature, food availability and other environmental stressors (including pollutants) could readily be examined in an experimental framework.

Behavioural effects of infection

Infection associated behavioural variation among wild-caught fish A number of authors have compared the behaviour of wild caught sticklebacks naturally infected with *S. solidus* with non-infected fish from the same population. These studies have identified a wide range of behaviours in which individual variation is associated with infection status, including shoaling behaviour (Barber and Huntingford, 1995; Barber *et al.*, 1995; Barber *et al.*, 1998) antipredator and risk-taking behaviour (Giles, 1983; Milinski, 1985; Giles, 1987b; Giles, 1987a; Godin and Sproul, 1988; Tierney *et al.*, 1993; Ness and Foster, 1999), prey choice (Milinski, 1984; Ranta, 1995) and competitive ability (Barber and Ruxton, 1998). In many cases behaviour studies are carried out to investigate hypotheses about the basis of altered behaviour, and specifically whether they may constitute examples of host ‘manipulation’ by parasites (Poulin, 1994). However although the results of these studies often suggest adaptive manipulation by parasites, such an approach can only ever produce correlational data, as infection status is not imposed experimentally. Alternative explanations, including the possibility that pre-existing behavioural variation influences exposure to infections, or underlying ‘quality’ factors that impact both susceptibility to infection and behaviour, mean that experimental infection studies are needed to unambiguously assign causality.

Behaviour change in experimentally infected fish In contrast to studies of wild-caught, naturally infected sticklebacks, relatively few have examined the behaviour in experimentally

infected sticklebacks. Aeschlimann *et al.* (2000) tested the risk taking behaviour of experimentally infected sticklebacks under threat of predation by pike *Esox lucius* during the early phase of infection before the parasite was infective to the definitive host. The aim of the study was to examine whether experimentally infected fish increased their risk-taking behaviour in order to maximise food intake, to reach sexual maturity early in an attempt to reduce the fitness impacts of infection. The results showed that during these early stages of infection, when host behaviour was predicted to reflect host responses to infection rather than being influenced by ‘manipulative’ parasites, there was no effect of infection status on the propensity to taking risks whilst foraging. This suggests that infected fish do not respond to infection by exploiting risky yet available prey, and is consistent with the finding that increased food intake actually appears to benefit parasites as well as hosts (Barber, 2005).

Studies of naturally and experimentally infected sticklebacks suggest that reduced predator avoidance behaviour is typically shown when fish harbour either a high burden (parasite index > 25%) of *S. solidus* (Milinski, 1984; Milinski, 1985) or when parasite mass exceeds 50mg (Tierney *et al.*, 1993), but not during early stages of infection (Aeschlimann *et al.*, 2000). Given the fact that 50mg appears to be the threshold mass for successful production of fertile eggs in the avian host (Tierney and Crompton, 1992), these observations are consistent with adaptive manipulation of behaviour. To examine this more closely, Barber *et al.* (2004) experimentally infected juvenile sticklebacks and used image analysis to track parasite growth alongside behavioural analysis of host escape responses over a 16 week p-e period. Reduced antipredator behaviour responses to a heron model were only observed in experimentally infected fish once plerocercoids had reached an estimated mass of 50mg, corroborating observational studies of the behaviour of naturally infected stickleback. This was the first study to use experimentally infected sticklebacks to demonstrate that *S. solidus* was responsible for the observed changes in behaviour.

554

555 *Potential for laboratory artefacts in behavioural studies* However, extrapolating results from
556 experimental lab studies to the field situation may again be difficult, as benign laboratory
557 conditions can also affect the behaviours exhibited by *S. solidus* infected sticklebacks.
558 Candolin and Voigt (2001) captured nest-holding males from a population in which 26% of
559 males harboured *S. solidus* and showed that nest holders were almost exclusively (33/35) non-
560 infected. They then transferred naturally infected fish to the lab, and found that after a 7d
561 period of *ad libitum* feeding, with access to nesting territory and materials, most infected fish
562 readily built nests and courted females. *Schistocephalus solidus* therefore appears to have
563 influenced reproductive performance of males in this population indirectly, by reducing the
564 ability of host sticklebacks to gain access to resources (food, territory, nesting material etc)
565 essential for successful reproduction. There also appears to be population variation in the
566 capacity to reverse the effects of infection under lab housing, with naturally infected males
567 from different populations being differentially capable of reproductive behaviour following a
568 period of benign housing (Rushbrook and Barber, 2006; Macnab *et al.*, in press).

569

570 *Manipulation of host behaviour: potential role of the immune system*

571 Reduced predator avoidance behaviour in the stickleback-*S. solidus* system is thought to be
572 caused by increased concentrations of monoamine neurotransmitters in neuronal tissues of the
573 brain in *S. solidus* infected sticklebacks (Øverli *et al.*, 2001). Whether the neuronal changes
574 are a consequence of changes in energy or endocrine status of the fish or are induced directly
575 by the parasite, for instance by the release of a neuroactive substance is unclear. However,
576 changes in neuroendocrine status are consistent with a chronic stress response in infected fish,
577 which could be, among other stressors, the result of an immune response (Øverli *et al.*, 2001).

Investigations of immune kinetics in *S. solidus* infected sticklebacks, detailed in section 2.3 above, reveal distinct changes in immune parameters during the period of parasite growth that corresponds with host behaviour change, when the parasites had passed the 50mg threshold weight between 40-60d p-e. Nonetheless, effects on parasite survival and fitness were not observed and plerocercoids kept on growing to attain 150mg at 67d p-e (Scharsack *et al.*, 2007) and about 200mg by 98d p-e (Scharsack *et al.*, 2004). As immune responses are presumably very costly for the host, but do not appear to have any effects on the parasite survival at this late stage of infection, one possible explanation for the observed pattern is that *S. solidus* – on attaining an infective size – triggers the stickleback immune system (in a ‘controlled’ manner that is not harmful for the parasite) to interfere with the crosstalk between neuro-endocrine system and immune system to induce reduced predation avoidance behaviour of its stickleback host (Scharsack *et al.*, 2007). A second explanation could be that *S. solidus* interacts directly with the neuro-endocrine system and that the observed changes in immune parameters are side effects of a host stress response. However, stress responses generally result in a broad (ubiquitous) activation of immunity, to which *S. solidus* is apparently vulnerable (Wedekind and Little, 2004; see above). The survival and growth of *S. solidus* plerocercoids relies on their ability to control the stickleback immune system, which would be more costly to maintain in a stress-induced activation of several immune traits, instead of single immune traits manipulated specifically by the parasite.

Due to the complex interactions between immunity and nervous system, it is difficult to distinguish whether a parasite directly or indirectly manipulates its host behaviour (Milinski, 1990; Adamo, 2002; Thomas *et al.*, 2005). In mammals the crosstalk between the immune response and brain is known as acute sickness behaviour, where behavioural changes that are associated with acute infections are typically immunologically mediated (Vollmer-Conna, 2001). For the *S. solidus* infection of sticklebacks, the exact mechanism responsible for translating the immune signal into a neural signal is still unclear, but it is well established in

teleosts and mammals that activation of innate immunity interferes with the neuro-endocrine system (Engelsma *et al.*, 2002; Dantzer, 2004). The observed changes in leukocyte responses during late infection with *S. solidus* could thus lead to the neuronal changes, which induce behavioural modifications of the stickleback and so elegantly enhance parasite transmission to the final host (Scharsack *et al.*, 2007)

STUDIES OF *IN VIVO* PARASITE GROWTH AND FITNESS

The large size of *S. solidus* plerocercoids relative to their stickleback hosts, and the subsequent distension of the host body cavity that is associated with infection, provides a useful tool to examine the growth of plerocercoids *in vivo*. The degree of distension caused by such parasites can be accurately measured using digital photography and image analysis software and converted into an estimate of plerocercoid mass (Barber, 1997; Loot *et al.*, 2002; Barber and Svensson, 2003), enabling plerocercoid growth to be examined non-invasively over the post-infection period. If fish are fed singly-infected copepods then the mass of individual plerocercoids can be tracked. This approach enabled Barber & Svensson (2003) to construct the growth curve for plerocercoids infecting stickleback hosts fed on a fixed ration of 8%bw.d⁻¹ and could be of considerable value in future studies designed to establish the impacts of host environmental factors on plerocercoid performance.

Furthermore, the ability to sexually mature the worms recovered from sticklebacks and collect data on adult fecundity makes it possible to examine the effects of host rearing environments and plerocercoid growth history on the fecundity of adult parasites (Dörücü *et al.*, 2007) as well as providing useful models for investigating egg production strategies, mate choice and the ‘hermaphrodite’s dilemma’ (Luscher and Wedekind, 2002).

CONCLUSIONS

In many ways *S. solidus* plerocercoids are unusual parasites; their typically extreme body size relative to the stickleback host is a dominant feature of infection that is not often observed in other host-parasite systems. It is likely that some of the effects of the parasites on stickleback hosts will be rather specific to this and a limited number of other systems that involve large bodied parasites, such as *Ligula intestinalis* infections of cyprinid fish, *Spirometra mansonoides* infections of mammals and some invertebrate-parasitoid systems. On the other hand, *S. solidus* exhibits features common to many ecologically important parasites, such as an indirect life cycle with trophic transmission and the potential to affect host growth, reproduction and survival. The great utility of the system lies in three key attributes; the typically important consequences that infections have for host performance, the ready availability of experimental infection techniques and the fact that the host fish is an extremely well characterised model organism. These attributes combine to facilitate experimental investigations into the role of parasites as agents of selection in host populations.

Acknowledgements

We are grateful to the editor and two anonymous referees for constructive, insightful comments on the manuscript.

647 **Table I.** Selected field studies documenting traits associated with *Schistocephalus solidus* infection in natural stickleback populations

Aspect of stickleback		
biology studied	Infection-associated trait	Reference
Nutritional condition	Infected fish had reduced body condition in the spring and autumn	(Tierney <i>et al.</i> , 1996)
	Infected fish had reduced stomach fullness and fed on smaller prey	(Tierney <i>et al.</i> , 1996)
	Seasonal differences in stomach fullness and diet composition of infected and non-infected fish	(Bergersen, 1996)
Sexual development	Infected fish had reduced body condition and liver energy reserves	(Tierney, 1994)
	Infected females showed reduced gonad development	(Arme and Owen, 1967)
	Infected females were less likely to be gravid	(Heins <i>et al.</i> , 1999)
Behaviour	Infected males developed less red nuptial coloration	(McPhail and Peacock, 1983)
	Infected males were less likely to hold nests	(Folstad <i>et al.</i> , 1994)
	Infected fish found further from cover in autumn	(Candolin and Voigt, 2001)
Morphology	Infected fish had less symmetric lateral plate counts	(Jakobsen <i>et al.</i> , 1988)
	Adults with asymmetric pelvis had increased incidence of infection (pattern reversed in 0+ fish).	(Reimchen and Nosil, 2001)
	Skin of infected fish was demelanised	(Reimchen, 1997)

References

- Adamo, S. A.** (2002). Modulating the modulators: parasites, neuromodulators and host behavioural change. *Brain, Behavior and Evolution* **60**, 370-377.
- Aeschlimann, P., Häberli, M. and Milinski, M.** (2000). Threat-sensitive feeding strategy of immature sticklebacks (*Gasterosteus aculeatus*) in response to recent experimental infection with the cestode *Schistocephalus solidus*. *Behavioral Ecology and Sociobiology* **49**, 1-7.
- Andersson, M.** (1994). *Sexual selection*, Princeton University Press, Princeton.
- Arme, C. and Owen, R. W.** (1967). Infections of 3-spined stickleback *Gasterosteus aculeatus* L with plerocercoid larvae of *Schistocephalus solidus* (Müller 1776) with special reference to pathological effects. *Parasitology* **57**, 301-304.
- Arnott, S. A., Barber, I. and Huntingford, F. A.** (2000). Parasite-associated growth enhancement in a fish-cestode system. *Proceedings of the Royal Society of London Series B-Biological Sciences* **267**, 657-663.
- Bagamian, K. H., Heins, D. C. and Baker, J. A.** (2004). Body condition and reproductive capacity of three-spined stickleback infected with the cestode *Schistocephalus solidus*. *Journal of Fish Biology* **64**, 1568-1576.
- Bakker, T. C. M. and Milinski, M.** (1991). The advantages of being red - sexual selection in the stickleback. In *Conference on Behavioural Ecology of Fishes* pp. 287-300. Erice, Italy.
- Barber, I.** (1997). A non-invasive morphometric technique for estimating cestode plerocercoid burden in small freshwater fish. *Journal of Fish Biology* **51**, 654-658.
- Barber, I.** (2005). Parasites grow larger in faster growing fish hosts. *International Journal for Parasitology* **35**, 137-143.
- Barber, I.** (2007). Host-parasite interactions of the three-spined stickleback. In *Biology of the Three-Spined Stickleback* (eds. Östlund-Nilsson, S., Mayer, I. & Huntingford, F. A.), pp. 271-318. CRC Press, Boca Raton, FL.
- Barber, I. and Arnott, S. A.** (2000). Split-clutch IVF: A technique to examine indirect fitness consequences of mate preferences in sticklebacks. *Behaviour* **137**, 1129-1140.
- Barber, I., Arnott, S. A., Braithwaite, V. A., Andrew, J. and Huntingford, F. A.** (2001). Indirect fitness consequences of mate choice in sticklebacks: offspring of brighter males grow slowly but resist

678 parasitic infections. *Proceedings of the Royal Society of London Series B-Biological Sciences* **268**, 71-
679 76.

680 **Barber, I., Downey, L. C. and Braithwaite, V. A.** (1998). Parasitism, oddity and the mechanism of shoal
681 choice. *Journal of Fish Biology* **53**, 1365-1368.

682 **Barber, I. and Huntingford, F. A.** (1995). The effect of *Schistocephalus solidus* (Cestoda: Pseudophyllidea) on
683 the foraging and shoaling behaviour of three-spined sticklebacks, *Gasterosteus aculeatus*. *Behaviour*
684 **132**, 1223-1240.

685 **Barber, I., Huntingford, F. A. and Crompton, D. W. T.** (1995). The effect of hunger and cestode parasitism
686 on the shoaling decisions of small fresh-water fish. *Journal of Fish Biology* **47**, 524-536.

687 **Barber, I. and Ruxton, G. D.** (1998). Temporal prey distribution affects the competitive ability of parasitized
688 sticklebacks. *Animal Behaviour* **56**, 1477-1483.

689 **Barber, I. and Svensson, P. A.** (2003). Effects of experimental *Schistocephalus solidus* infections on growth,
690 morphology and sexual development of female three-spined sticklebacks, *Gasterosteus aculeatus*.
691 *Parasitology* **126**, 359-367.

692 **Barber, I., Walker, P. and Svensson, P. A.** (2004). Behavioural responses to simulated avian predation in
693 female three spined sticklebacks: The effect of experimental *Schistocephalus solidus* infections.
694 *Behaviour* **141**, 1425-1440.

695 **Barber, I., Wright, H., Arnott, S. and Wootton, R.** (2008). Growth and energetics in the stickleback–
696 *Schistocephalus* host–parasite system: a review of experimental infection studies. *Behaviour* **145**, 647-
697 668.

698 **Bell, A. M.** (2005). Behavioural differences between individuals and two populations of stickleback
699 (*Gasterosteus aculeatus*). *Journal of Evolutionary Biology* **18**, 464-473.

700 **Bell, A. M. and Stamps, J. A.** (2004). Development of behavioural differences between individuals and
701 populations of sticklebacks, *Gasterosteus aculeatus*. *Animal Behaviour* **68**, 1339-1348.

702 **Bell, M. A. and Foster, S. A.** (Eds.) (1994). *The Evolutionary Biology of the Threespine Stickleback*, Oxford
703 University Press, Oxford, UK.

704 **Bergersen, R.** (1996). Sticklebacks from Greenland. *Journal of Fish Biology* **48**, 799-801.

705 **Bråten, T.** (1966). Host specificity in *Schistocephalus solidus*. *Parasitology* **56**, 657-&.

706 **Bunnajirakul, S., Steinhagen, D., Hetzel, U., Korting, W. and Drommer, W.** (2000). A study of sequential
707 histopathology of *Trypanoplasma borreli* (Protozoa : Kinetoplastida) in susceptible common carp
708 *Cyprinus carpio*. *Diseases of Aquatic Organisms* **39**, 221-229.

709 **Candolin, U. and Voigt, H. R.** (2001). No effect of a parasite on reproduction in stickleback males: a laboratory
710 artefact? *Parasitology* **122**, 457-464.

711 **Chappell, L. H.** (1969). Parasites of three-spined stickleback *Gasterosteus aculeatus* L from a Yorkshire Pond
712 .1. Seasonal variation of parasite fauna. *Journal of Fish Biology* **1**, 137-152.

713 **Chubb, J. C., Seppala, O., Luscher, A., Milinski, M. and Valtonen, E. T.** (2006). *Schistocephalus cotti* n. sp
714 (Cestoda : Pseudophyllidea) plerocercoids from bullheads *Cottus gobio* L. in an Arctic river in Finland,
715 with a key to the plerocercoids of the Palaearctic species of the genus. *Systematic Parasitology* **65**,
716 161-170.

717 **Clarke, A. S.** (1954). Studies on the life cycle of the pseudophyllidean cestode *Schistocephalus solidus*.
718 *Proceedings of the Zoological Society of London* **124**, 257–302.

719 **Colosimo, P. F., Hosemann, K. E., Balabhadra, S., Villarreal, G., Dickson, M., Grimwood, J., Schmutz, J.,**
720 **Myers, R. M., Schluter, D. and Kingsley, D. M.** (2005). Widespread parallel evolution in sticklebacks
721 by repeated fixation of ectodysplasin alleles. *Science* **307**, 1928-1933.

722 **Coombs, I. and Crompton, D. W. T.** (1991). *A guide to human helminths* Taylor & Francis Ltd, London.

723 **Dantzer, R.** (2004). Cytokine-induced sickness behaviour: A neuroimmune response to activation of innate
724 immunity. *European Journal of Pharmacology* **500**, 399-411.

725 **Dick, T. L.** (1816). Account of the worm with which the stickleback is infested. *Annals of Philosophy* **7**, 106-
726 109.

727 **Dörücü, M., Wilson, D. and Barber, I.** (2007). Differences in adult egg output of *Schistocephalus solidus* from
728 singly- and multiply-infected sticklebacks. *Journal of Parasitology* **93**, 1521-1523.

729 **Engelsma, M. Y., Huising, M. O., Van Muiswinkel, W. B., Flik, G., Kwang, J., Savelkoul, H. F. and**
730 **Verburg-Van Kemenade, B. M.** (2002). Neuroendocrine-immune interactions in fish: a role for
731 interleukin-1. *Veterinary Immunology and Immunopathology* **87**, 467-479.

732 **Folstad, I., Hope, A. M., Karter, A. and Skorping, A.** (1994). Sexually selected color in male sticklebacks - a
733 signal of both parasite exposure and parasite resistance. *Oikos* **69**, 511-515.

734 **French, J. R. P. and Muzzall, P. M.** (2008). First report of *Schistocephalus* sp (Cestoda : pseudophyllidea) in
735 slimy sculpin, *Cottus cognatus* Richardson, 1836, from Lake Michigan, USA. *Comparative*
736 *Parasitology* **75**, 132-134.

737 **Gibson, G.** (2005). The synthesis and evolution of a supermodel. *Science* **307**, 1890-1891.

738 **Giles, N.** (1983). Behavioral effects of the parasite *Schistocephalus solidus* (Cestoda) on an intermediate host,
739 the 3-spined stickleback, *Gasterosteus aculeatus* L. *Animal Behaviour* **31**, 1192-1194.

- Giles, N.** (1987a). A comparison of the behavioral responses of parasitized and non-parasitized 3-spined sticklebacks, *Gasterosteus aculeatus* L, to progressive hypoxia. *Journal of Fish Biology* **30**, 631-638.
- Giles, N.** (1987b). Predation risk and reduced foraging activity in fish - Experiments with parasitized and non-parasitized 3-spined sticklebacks, *Gasterosteus aculeatus* L. *Journal of Fish Biology* **31**, 37-44.
- Godin, J. G. J. and Sproul, C. D.** (1988). Risk-taking in parasitized sticklebacks under threat of predation - Effects of energetic need and food availability. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* **66**, 2360-2367.
- Hammerschmidt, K., Koch, K., Milinski, M., Chubb, J. C. and Parker, G. A.** (2009). When to go: optimization of host switching in parasites with complex life cycles. *Evolution* [Epub ahead of print].
- Hammerschmidt, K. and Kurtz, J.** (2005). Surface carbohydrate composition of a tapeworm in its consecutive intermediate hosts: Individual variation and fitness consequences. *International Journal for Parasitology* **35**, 1499-1507.
- Hammerschmidt, K. and Kurtz, J.** (2007). *Schistocephalus solidus*: Establishment of tapeworms in sticklebacks - fast food or fast lane? *Experimental Parasitology* **116**, 142-149.
- Hammerschmidt, K. and Kurtz, J.** (2009). Ecological immunology of a tapeworms' interaction with its two consecutive hosts. *Advances in Parasitology* **68**, 111-137.
- Hart, B. L.** (1990). Behavioral adaptations to pathogens and parasites - 5 strategies. *Neuroscience and Biobehavioral Reviews* **14**, 273-294.
- Hart, B. L.** (1992). Behavioral adaptations to parasites - an ethological approach. *Journal of Parasitology* **78**, 256-265.
- Hart, B. L.** (1997). Behavioural defence. In *Host-parasite evolution: general principle and avian models* (eds. Clayton, D. H. & Moore, J.), pp. 59-77. Oxford University Press, Oxford, UK.
- Heins, D. C. and Baker, J. A.** (2008). The stickleback-*Schistocephalus* host-parasite system as a model for understanding the effect of a macroparasite on host reproduction. *Behaviour* **145**, 625-645.
- Heins, D. C., Singer, S. S. and Baker, J. A.** (1999). Virulence of the cestode *Schistocephalus solidus* and reproduction in infected threespine stickleback, *Gasterosteus aculeatus*. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* **77**, 1967-1974.
- Herve, M., Angeli, V., Pinzar, E., Wintjens, R., Faveeuw, C., Narumiya, S., Capron, A., Urade, Y., Capron, M., Riveau, G. and Trottein, F.** (2003). Pivotal roles of the parasite PGD(2) synthase and of the host D prostanoid receptor 1 in schistosome immune evasion. *European Journal of Immunology* **33**, 2764-2772.

771 **Hoberg, E. P., Henny, C. J., Hedstrom, O. R. and Grove, R. A.** (1997). Intestinal helminths of river otters
772 (*Lutra canadensis*) from the Pacific Northwest. *Journal of Parasitology* **83**, 105-110.

773 **Hopkins, C. A. and Smyth, J. D.** (1951). Notes on the Morphology and Life History of *Schistocephalus*-
774 *Solidus* (Cestoda, Diphyllbothriidae). *Parasitology* **41**, 283-&.

775 **Hyslop, E. J. and Chubb, J. C.** (1983). *Schistocephalus pungitii* Dubinina, 1959 (Cestoda) in Britain.
776 *Parasitology* **87**, R30-R30.

777 **Jäger, I. and Schjørring, S.** (2006). Multiple infections: Relatedness and time between infections affect the
778 establishment and growth of the cestode *Schistocephalus solidus* in its stickleback host. *Evolution* **60**,
779 616-622.

780 **Jakobsen, P. J., Johnsen, G. H. and Larsson, P.** (1988). Effects of predation risk and parasitism on the feeding
781 ecology, habitat use, and abundance of lacustrine threespine stickleback (*Gasterosteus aculeatus*).
782 *Canadian Journal of Fisheries and Aquatic Sciences* **45**, 426-431.

783 **Jakobsen, P. J. and Wedekind, C.** (1998). Copepod reaction to odor stimuli influenced by cestode infection.
784 *Behavioral Ecology* **9**, 414-418.

785 **Kennedy, C. R.** (1974). Checklist of British and Irish freshwater fish parasites with notes on their distribution.
786 *Journal of Fish Biology* **6**, 613-644.

787 **Keymer, A. E. and Read, A. F.** (1991). Behavioural ecology: the impact of parasitism. In *Parasite-Host*
788 *Associations, Coexistence or Conflict* (eds. Toft, C. A., Aeschlimann, A. & Bolis, L.), Oxford Scientific
789 Publications, Oxford, UK.

790 **Kingsley, D. M.** (2003). Sequencing the genome of threespine sticklebacks (*Gasterosteus aculeatus*). In
791 *National Human Genome Research Institute White Paper*

792 **Kingsley, D. M., Zhu, B. L., Osoegawa, K., De Jong, P. J., Schein, J., Marra, M., Peichel, C., Amamiya, C.,**
793 **Schluter, D., Balabhadra, S., Friedlander, B., Cha, Y. M., Dickson, M., Grimwood, J., Schmutz,**
794 **J., Talbot, W. S. and Myers, R.** (2004). New genomic tools for molecular studies of evolutionary
795 change in threespine sticklebacks. *Behaviour* **141**, 1331-1344.

796 **Kurtz, J., Kalbe, M., Aeschlimann, P. B., Haberli, M. A., Wegner, K. M., Reusch, T. B. H. and Milinski,**
797 **M.** (2004). Major histocompatibility complex diversity influences parasite resistance and innate
798 immunity in sticklebacks. *Proceedings of the Royal Society of London Series B-Biological Sciences*
799 **271**, 197-204.

800 **Lafferty, K. D.** (1992). Foraging on prey that are modified by parasites. *American Naturalist* **140**, 854-867.

801 **Lester, R. J. G.** (1971). Influence of *Schistocephalus* plerocercoids on respiration of *Gasterosteus* and a possible
802 resulting effect on behavior of fish. *Canadian Journal of Zoology* **49**, 361-366.

803 **Lobue, C. P. and Bell, M. A.** (1993). Phenotypic manipulation by the cestode parasite *Schistocephalus solidus*
804 of its intermediate host, *Gasterosteus aculeatus*, the threespine stickleback. *American Naturalist* **142**,
805 725-735.

806 **Loot, G., Giraudel, J. L. and Lek, S.** (2002). A non-destructive morphometric technique to predict *Ligula*
807 *intestinalis* L. plerocercoid load in roach (*Rutilus rutilus* L.) abdominal cavity. *Ecological Modelling*
808 **156**, 1-11.

809 **Luscher, A. and Wedekind, C.** (2002). Size-dependent discrimination of mating partners in the simultaneous
810 hermaphroditic cestode *Schistocephalus solidus*. *Behavioral Ecology* **13**, 254-259.

811 **Maccoll, A. D. C.** (2009). Parasite burdens differ between sympatric three-spined stickleback species.
812 *Ecography* **32**, 153-160.

813 **Macnab, V. L., Katsiadaki, I. and Barber, I.** (in press). Reproductive potential of *Schistocephalus solidus*
814 infected male three-spined stickleback (*Gasterosteus aculeatus*) from two UK populations. *Journal of*
815 *Fish Biology*.

816 **Maizels, R. M. and Yazdanbakhsh, M.** (2003). Immune regulation by helminth parasites: cellular and
817 molecular mechanisms. *Nature Reviews Immunology* **3**, 733-744.

818 **Manning, M. J.** (1994). Fishes. In *Immunology: a comparative approach* (ed. Turner, R. J.), John Willey &
819 Sons, Chichester, New York, USA.

820 **Mckinnon, J. S., Mori, S., Blackman, B. K., David, L., Kingsley, D. M., Jamieson, L., Chou, J. and**
821 **Schluter, D.** (2004). Evidence for ecology's role in speciation. *Nature* **429**, 294-298.

822 **Mcphail, J. D. and Peacock, S. D.** (1983). Some effects of the cestode (*Schistocephalus solidus*) on
823 reproduction in the threespine stickleback (*Gasterosteus aculeatus*): Evolutionary aspects of a host-
824 parasite interaction. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* **61**, 901-908.

825 **Milinski, M.** (1984). Parasites determine a predator's optimal feeding strategy. *Behavioral Ecology and*
826 *Sociobiology* **15**, 35-37.

827 **Milinski, M.** (1985). Risk of predation of parasitized sticklebacks (*Gasterosteus aculeatus* L) under competition
828 for food. *Behaviour* **93**, 203-215.

829 **Milinski, M.** (1990). Parasites and host decision-making. In *Parasitism and host behaviour* (eds. Barnard, C. J.
830 & Behnke, J. M.), pp. 95-116. Taylor and Francis, London.

831 **Milinski, M. and Bakker, T. C. M.** (1990). Female sticklebacks use male coloration in mate choice and hence
832 avoid parasitized males. *Nature* **344**, 330-333.

833 **Minchella, D. J.** (1985). Host life-history variation in response to parasitism. *Parasitology* **90**, 205-216.

834 **Ness, J. H. and Foster, S. A.** (1999). Parasite-associated phenotype modifications in threespine stickleback.
835 *Oikos* **85**, 127-134.

836 **Orr, T. S. C., Hopkins, C. A. and Charles, G. H.** (1969). Host specificity and rejection of *Schistocephalus*
837 *solidus*. *Parasitology* **59**, 683-&.

838 **Östlund -Nilsson, S., Mayer, I. and Huntingford, F. A.** (2006). *Biology of the three-spined stickleback*, CRC
839 Press, Boca Raton. FL.

840 **Øverli, O., Páll, M., Borg, B., Jobling, M. and Winberg, S.** (2001). Effects of *Schistocephalus solidus*
841 infection on brain monoaminergic activity in female three-spined sticklebacks *Gasterosteus aculeatus*.
842 *Proceedings of the Royal Society of London Series B-Biological Sciences* **268**, 1411-1415.

843 **Parker, G. A., Ball, M. A., Chubb, J. C., Hammerschmidt, K. and Milinski, M.** (2009). When should a
844 trophically transmitted parasite manipulate Its host? *Evolution* **63**, 448-458.

845 **Pasternak, A. F., Huntingford, F. A. and Crompton, D. W. T.** (1995). Changes in metabolism and behavior
846 of the fresh-water copepod *Cyclops strenuus abyssorum* Infected with *Diphyllbothrium* spp.
847 *Parasitology* **110**, 395-399.

848 **Peichel, C. L., Nereng, K. S., Ohgi, K. A., Cole, B. L. E., Colosimo, P. F., Buerkle, C. A., Schluter, D. and**
849 **Kingsley, D. M.** (2001). The genetic architecture of divergence between threespine stickleback species.
850 *Nature* **414**, 901-905.

851 **Pennycuik, L.** (1971). Quantitative effects of three species of parasites on a population of three-spined
852 sticklebacks, *Gasterosteus aculeatus*. *Journal of Zoology* **165**, 143-162.

853 **Poulin, R.** (1994). The evolution of parasite manipulation of host behavior - a theoretical analysis. *Parasitology*
854 **109**, S109-S118.

855 **Ranta, E.** (1995). *Schistocephalus* infestation improves prey-size selection by 3-spined sticklebacks,
856 *Gasterosteus aculeatus*. *Journal of Fish Biology* **46**, 156-158.

857 **Reimchen, T. E.** (1997). Parasitism of asymmetrical pelvic phenotypes in stickleback. *Canadian Journal of*
858 *Zoology-Revue Canadienne De Zoologie* **75**, 2084-2094.

859 **Reimchen, T. E. and Nosil, P.** (2001). Lateral plate asymmetry, diet and parasitism in threespine stickleback.
860 *Journal of Evolutionary Biology* **14**, 632-645.

861 **Rijkers, G. T., Frederixwolters, E. M. H. and Vanmuiswinkel, W. B.** (1980). The immune system of cyprinid
862 fish - kinetics and temperature dependence of antibody producing cells in carp (*Cyprinus carpio*).
863 *Immunology* **41**, 91-97.

864 **Rushbrook, B. J. and Barber, I.** (2006). Nesting, courtship and kidney hypertrophy in *Schistocephalus*-infected
865 male three-spined stickleback from an upland lake. *Journal of Fish Biology* **69**, 870-882.

866 **Scharsack, J. P., Kalbe, M., Derner, R., Kurtz, J. and Milinski, M.** (2004). Modulation of granulocyte
867 responses in three-spined sticklebacks *Gasterosteus aculeatus* infected with the tapeworm
868 *Schistocephalus solidus*. *Diseases of Aquatic Organisms* **59**, 141-150.

869 **Scharsack, J. P., Koch, K. and Hammerschmidt, K.** (2007). Who is in control of the stickleback immune
870 system: interactions between *Schistocephalus solidus* and its specific vertebrate host. *Proceedings of the*
871 *Royal Society B-Biological Sciences* **274**, 3151-3158.

872 **Seppala, T., Chubb, J. C., Niemela, E. and Valtonen, E. T.** (2007). Introduced bullheads *Cottus gobio* and
873 infection with plerocercoids of *Schistocephalus cotti* in the Utsjoki, an Arctic river in Finland. *Journal*
874 *of Fish Biology* **70**, 1865-1876.

875 **Shapiro, M. D., Bell, M. A. and Kingsley, D. M.** (2006). Parallel genetic origins of pelvic reduction in
876 vertebrates. *Proceedings of the National Academy of Sciences of the United States of America* **103**,
877 13753-13758.

878 **Shapiro, M. D., Marks, M. E., Peichel, C. L., Blackman, B. K., Nereng, K. S., Jonsson, B., Schluter, D. and**
879 **Kingsley, D. M.** (2004). Genetic and developmental basis of evolutionary pelvic reduction in threespine
880 sticklebacks. *Nature* **428**, 717-723.

881 **Smyth, J. D.** (1946). Studies on tapeworm physiology. I. The cultivation of *Schistocephalus solidus* in vitro.
882 *Journal of Experimental Biology* **23**, 47-70.

883 **Smyth, J. D.** (1947). The physiology of tapeworms. *Biological Reviews* **22**, 214-238.

884 **Smyth, J. D.** (1990). *In vitro cultivation of parasitic helminths* CRC Press, Boca Raton, FL.

885 **Smyth, J. D. and Mcmanus, D. P.** (1989). *The physiology and biochemistry of cestodes*, Cambridge University
886 Press, Cambridge.

887 **Takizawa, F., Araki, K., Kobayashi, I., Moritomo, T., Ototake, M. and Nakanishi, T.** (2008a). Molecular
888 cloning and expression analysis of T-bet in ginbuna crucian carp (*Carassius auratus langsdorfii*).
889 *Molecular Immunology* **45**, 127-136.

890 **Takizawa, F., Mizunaga, Y., Araki, K., Moritomo, T., Ototake, M. and Nakanishi, T.** (2008b). GATA3
891 mRNA in ginbuna crucian carp (*Carassius auratus langsdorfii*): cDNA cloning, splice variants and
892 expression analysis. *Developmental and Comparative Immunology* **32**, 898-907.

893 **Thomas, F., Adamo, S. and Moore, J.** (2005). Parasitic manipulation: where are we and where should we go?
894 *Behavioural Processes* **68**, 185-199.

895 **Tierney, J. F.** (1994). Effects of *Schistocephalus solidus* (Cestoda) on the food intake and diet of the three-
896 spined stickleback, *Gasterosteus aculeatus*. *Journal of Fish Biology* **44**, 731-735.

897 **Tierney, J. F. and Crompton, D. W. T.** (1992). Infectivity of plerocercoids of *Schistocephalus solidus*
898 (Cestoda, Ligulidae) and fecundity of the adults in an experimental definitive host, *Gallus gallus*.
899 *Journal of Parasitology* **78**, 1049-1054.

900 **Tierney, J. F., Huntingford, F. A. and Crompton, D. W. T.** (1993). The relationship between infectivity of
901 *Schistocephalus solidus* (Cestoda) and antipredator behavior of its intermediate host, the three-spined
902 stickleback, *Gasterosteus aculeatus*. *Animal Behaviour* **46**, 603-605.

903 **Tierney, J. F., Huntingford, F. A. and Crompton, D. W. T.** (1996). Body condition and reproductive status in
904 sticklebacks exposed to a single wave of *Schistocephalus solidus* infection. *Journal of Fish Biology* **49**,
905 483-493.

906 **Urdal, K., Tierney, J. F. and Jakobsen, P. J.** (1995). The tapeworm *Schistocephalus solidus* alters the activity
907 and response, but not the predation susceptibility of infected copepods. *Journal of Parasitology* **81**,
908 330-333.

909 **Vanmuiswinkel, W. B.** (1995). The piscine immune system: innate and acquired immunity. In *Fish Diseases*
910 *and Disorders. Volume 1, Protozoan and Metazoan Infections* (ed. Woo, P. T. K.), pp. 729-750. CAB
911 International, Wallingford, UK.

912 **Vollmer-Conna, U.** (2001). Acute sickness behaviour: an immune system-to-brain communication?
913 *Psychological Medicine* **31**, 761-767.

914 **Walkey, M. and Meakins, R. H.** (1970). An attempt to balance energy budget of a host-parasite system.
915 *Journal of Fish Biology* **2**, 361-372.

916 **Wang, L. J., Cao, Y. and Shi, H. N.** (2008). Helminth infections and intestinal inflammation. *World Journal of*
917 *Gastroenterology* **14**, 5125-5132.

918 **Wedekind, C.** (1997). The infectivity, growth, and virulence of the cestode *Schistocephalus solidus* in its first
919 intermediate host, the copepod *Macrocyclops albidus*. *Parasitology* **115**, 317-324.

- Wedekind, C. and Little, T. J.** (2004). The clearance of hidden cestode infection triggered by an independent activation of host defense in a teleost fish. *Journal of Parasitology* **90**, 1329-1331.
- Wedekind, C. and Milinski, M.** (1996). Do three-spined sticklebacks avoid consuming copepods, the first intermediate host of *Schistocephalus solidus*? An experimental analysis of behavioural resistance. *Parasitology* **112**, 371-383.
- Wegner, K. M., Kalbe, M., Kurtz, J., Reusch, T. B. H. and Milinski, M.** (2003). Parasite selection for immunogenetic optimality. *Science* **301**, 1343-1343.
- Wootton, R. J.** (1976). *The biology of the sticklebacks* Academic Press, London
- Wootton, R. J.** (1984). *A functional biology of sticklebacks*, University of California Press, Berkeley, CA.
- Wright, H. A., Wootton, R. J. and Barber, I.** (2006). The effect of *Schistocephalus solidus* infection on meal size of three-spined stickleback. *Journal of Fish Biology* **68**, 801-809.
- Wright, H. A., Wootton, R. J. and Barber, I.** (2007). Compensatory growth in threespine sticklebacks (*Gasterosteus aculeatus*) inhibited by experimental *Schistocephalus* infection. *Canadian Journal of Fisheries and Aquatic Sciences* **64**, 819-826.