

Characteristics of north Jovian aurora from STIS FUV spectral images

5

J. Gustin¹, D. Grodent¹, L.C. Ray², B. Bonfond¹, E.J. Bunce³, J. D. Nichols³ and N.
Ozak⁴

10

15 ¹ Laboratoire de Physique Atmosphérique et Planétaire, Université de Liège, B-4000 Liège, Belgium

² Department of Physics and Astronomy, University College London, London, UK

³ Department of Physics and Astronomy, University of Leicester, Leicester LE1 7RH, UK

⁴ Department of Earth and Planetary Sciences, Weizmann Institute of Science, 76100 Rehovot, Israel

20

25 Revised version, October 2015

Abstract

30 We analyzed two observations obtained in Jan. 2013, consisting of spatial scans of the Jovian
north ultraviolet aurora with the HST Space Telescope Imaging Spectrograph (STIS) in the
spectroscopic mode. The color ratio (CR) method, which relates the wavelength-dependent absorption
of the FUV spectra to the mean energy of the precipitating electrons, allowed us to determine important
characteristics of the entire auroral region. The results show that the spatial distribution of the
35 precipitating electron energy is far from uniform. The morning main emission arc is associated with
mean energies of around 265 keV, the afternoon main emission (kink region) has energies near 105
keV, while the 'flare' emissions poleward of the main oval are characterized by electrons in the 50-85
keV range. A small scale structure observed in the discontinuity region is related to electrons of 232
keV and the Ganymede footprint shows energies of 157 keV. Interestingly, each specific region shows
40 very similar behavior for the two separate observations.

The Io footprint shows a weak but undeniable hydrocarbon absorption, which is not consistent
with altitudes of the Io emission profiles (~900 km relative to the 1 bar level) determined from HST-
ACS observations. An upward shift of the hydrocarbon homopause of at least 100 km is required to
reconcile the high altitude of the emission and hydrocarbon absorption.

45 The relationship between the energy fluxes and the electron energies has been compared to
curves obtained from Knight's theory of field-aligned currents. Assuming a fixed electron temperature
of 2.5 keV, an electron source population density of $\sim 800 \text{ m}^{-3}$ and $\sim 2400 \text{ m}^{-3}$ is obtained for the
morning main emission and kink regions, respectively. Magnetospheric electron densities are lowered
for the morning main emission ($\sim 600 \text{ m}^{-3}$) if the relativistic version of Knight's theory is applied.

50 Lyman and Werner H_2 emission profiles resulting from secondary electrons, produced by
precipitation of heavy ions in the 1-2 MeV/u range, have been applied to our model. The low CR

obtained from this emission suggests that heavy ions, presumably the main source of the X-ray aurora, do not significantly contribute to typical UV polar emission.

1. Introduction

55

1.1. Background

The ultraviolet Jovian aurora is mainly produced by the interaction between the H₂ atmosphere and precipitating magnetospheric electrons. In the far ultraviolet (FUV, between 1200 and 1700 Å), the emission is dominated by the Lyman- α line from atomic hydrogen resulting from H₂ dissociation and
60 H₂ vibronic lines from the Lyman ($B^1\Sigma_g^+ \rightarrow X^1\Sigma_g^+$) and Werner ($C^1\Pi_u^+ \rightarrow X^1\Sigma_g^+$) system bands. The auroral emission is known to interact with the atmosphere through absorption by the main hydrocarbons. Methane (CH₄) attenuates the emission at wavelengths below 1400 Å, ethane (C₂H₆), which has a continuous absorption cross-section shortward of 1550 Å, has a typical signature between
1400 and 1480 Å in the case of strongly attenuated spectra, and acetylene (C₂H₂) which has a more
65 peaked cross-section and has a significant effect at 1480 Å and 1520 Å. This very wavelength-dependent absorption is measured by the color ratio $CR = I(1550-1620\text{\AA}) / I(1230-1300\text{\AA})$ with I the brightness in photons unit, originally introduced in a slightly different form by Yung et al. (1982). The CR is directly correlated with the amount of hydrocarbon at or above the H₂ auroral emission and is related to the altitude of the aurora relative to the hydrocarbons. It is thus an important tool to
70 determine the energy of the primary electrons precipitating into the atmosphere. While broadband images are commonly used to characterize spatial and temporal effects of the aurora, FUV spectra allow the determination of the mean energy $\langle E \rangle$ of the precipitating electrons and improve our knowledge of the processes leading to the different auroral regions.

UV images obtained with HST cameras (STIS in the imaging mode and the Advanced Camera
75 for Survey (ACS)) have shown a complex and dynamical auroral structure and revealed that the auroral morphology is characterized by several distinct components of emission.

The first component is the main oval, which maps magnetically to the magnetosphere between

~20 and 30 Jovian radius (R_J) in the equatorial plane, and results from the magnetosphere–ionosphere coupling current system associated with the breakdown of corotation in the middle magnetosphere, specifically with the region of upward field-aligned currents (e.g. Cowley and Bunce (2001)). This emission has an apparent brightness generally in the 35-450 kR range of H_2 in the Lyman and Werner bands ($1 \text{ kR} = 10^9 \text{ photons per cm}^2 \text{ per sec emitted in } 4 \pi \text{ steradians}$). Occasionally, emissions of a few MR have been observed in the morning sector. The main emission is characterized by a narrow bright arc extending from dawn to noon or further. This main emission generally becomes more diffuse and weaker in the afternoon sector than in the morning sector, with a 'kink' region in the 110-150° System III sector of the northern aurora, associated with a localized magnetic anomaly (Grodent et al. 2008). Time-tagged observations show that the main oval energy flux usually varies steadily over several minute intervals and is somewhat correlated with the mean energy of the precipitating electrons, such that the current density (i.e. the electron flux) remains relatively constant (Gustin et al., 2004a). Assuming an emission exclusively due to electrons, the auroral mean electron energy lies within the range 35-200 keV. Two cases with harder electrons (~280 keV and ~460 keV) have also been revealed from a STIS observation of an exceptionally bright morning aurora (Gustin et al., 2006). Overall, Gustin et al. (2004a) showed that the main oval properties are consistent with Knight's theory of auroral electron acceleration associated with field-aligned current flow (Knight, 1973), involving a magnetospheric source of electrons with a thermal energy of 2.5 keV and with a density varying between 1000 and 10000 m^{-3} , consistent with "warm" magnetospheric electrons observed by Voyager (Scudder et al., 1981). A discontinuity region in the main emission, characterized by a significant drop of the brightness and fixed between 08:00 and 13:00 magnetic local time (MLT) has also been identified in HST images (Radioti et al. 2008). Work by Chané et al. (2013) suggests that this discontinuity is the result of an asymmetry in the thermal pressure distribution, decreasing the strength of the corotation current system in that sector. Finally, a transient small-scale structure in the main

emission, close to magnetic noon, has recently been identified by Palmaerts et al. (2014). This structure is present in both hemispheres, located between 09:00 and 15:30 MLT in the northern hemisphere and between 10:00 and 15:00 MLT in the southern hemisphere. Intermittent plasma flow in the equatorial plane near the noon sector is advanced as a possible explanation of this emission.

The second component, located poleward of the main oval, consists of scattered diffuse emissions, generally weaker than the main oval ($\sim 15 - \sim 70$ kR). These “polar cap” emissions exhibit large time variations on timescales of minutes or less (“flares”). Occasionally, very dynamical, intense and sometimes quasi-periodic brightenings of several MR have been observed, possibly related to abrupt solar wind variations (Waite et al. (2001), Bonfond et al. (2011)). The latter bright polar flares have never been directly observed with UV spectrographs, which makes it problematic to determine their characteristics, but STIS spectra obtained from “typical” polar emissions are generally associated with mean electron energies in the 30-100 keV range. Values up to ~ 180 keV have been observed but are exceptional, considering the few observations available. Furthermore, the energy fluxes and current densities associated with polar emissions determined by Gustin et al. (2004 a) are significantly lower than main oval values (≤ 10 mWm $^{-2}$ and ≤ 0.12 μ Am $^{-2}$ for polar regions compared to values up to 30 mWm $^{-2}$ and 0.3 μ Am $^{-2}$ for the energy flux and current density, respectively, of the main oval). At the higher end of the energy flux range in this region, the auroral acceleration mechanism may be similar to that in the main oval, while the weaker emissions are not clearly associated with field-aligned currents and could result from precipitation from a quasi-isotropic hot magnetospheric electron source (Gustin et al. 2006). The time evolution of $\langle E \rangle$ generally shows little correlation (positive or negative) with the energy flux precipitated during these transient events. This behavior suggests that the mechanism responsible for rapid transient brightenings does not increase the energy of the precipitated electrons, but it enhances their number flux (Gérard et al. 2003).

The third type of auroral emission consists of spots at the magnetic footprints of Io, Ganymede

and Europa, observed both in UV and in infrared wavelengths (Prangé et al. (1996, 1998), Grodent et al., 2006, Clarke et al., 1996, 1998, 2002). The Io footprint (IFP) consists of several spots observed in both hemispheres (Bonfond et al. 2009), followed by a trailing tail that sometimes extends for more than 180° in longitude (Clarke et al., 1998, Gérard et al., 2002). The IFP brightness in the Lyman and
130 Werner H₂ bands may be very variable with time and location, as spectra and images of the IFP shows observed values from 35 kR (e.g. Bonfond et al., 2009), with peaks of up to 670 kR (e.g. Gérard et al., 2002) and all intermediate values (Wannawichian et al., 2010). The vertical brightness of the IFP spots can go up to 6000 kR and depend on both the S_{III} longitude of the satellite and the hemisphere (Bonfond et al. 2013). The IFP spectral observations are characterized by remarkably stable CR from
135 1.4 to 2.3. Compared to the other auroral regions, where the CR can vary from 1.4 to more than 10, these IFP color ratios (CRs) correspond to precipitating electrons of mean energy between 40 keV and 70 keV. These values were obtained from the methane vertical profile and the pressure-altitude relationship in the North Equatorial Belt (NEB) atmosphere from Gladstone et al. (1996), adapted to gravity at the polar region. Study of limb profiles by Bonfond et al. (2009) shows that the peak of IFP
140 emission profile is situated between 550 and 1300 km. The peak altitude, at ~900 km on average, is found to be relatively constant with distance from the main spot. By using Monte Carlo simulations, these authors showed that an observed emission peaking at 900 km is compatible with precipitation of electrons distributed as a kappa function with a mean energy of ~1.1 keV, much lower than the values obtained by Gérard et al. (2002). This discrepancy, still not elucidated, is usually interpreted as being
145 the result of our poor knowledge of the hydrocarbon density profiles in polar regions, which is thought to be very different from observed and modeled profiles determined at low latitudes because of the energy deposition induced by the aurora.

A fourth component consist of emissions equatorward of the main emission: transient emissions possibly due to magnetospheric injections (Dumont et al. 2014) and diffuse emissions,

150 presumably associated with electron scattering by whistler mode waves (Radioti et al. 2009).

1.2. Objectives of the study

Contrary to the auroral brightness, easily monitored by the thousands of images obtained with HST instruments, the factors which influence the primary electron mean energy variations are difficult to interpret for two reasons. First, the limited number of spectral observations available has been obtained through the long 1997-2009 period and thus reflects various solar and magnetospheric-ionospheric conditions, which make it difficult to directly compare different measurements. Second, the observed spectra either cover a limited portion of the aurora due to the small size of the aperture projected onto the planet (e.g. Gladstone and Skinner (1989) with the International Ultraviolet Explorer, Dols et al. (2000) with the Goddard High resolution spectrograph onboard the Hubble Space Telescope (HST), Gustin et al. (2002, 2004a, 2006), Gérard et al. (2002, 2003) with the HST Space Telescope Imaging Spectrograph (STIS) in the spectroscopic mode), or integrated the whole polar region without spatial resolution (e.g. observations with the Far Ultraviolet Spectroscopic Explorer by Gustin et al. 2004b). Such former spectroscopic observations thus do not allow us to draw a global picture of the processes responsible for polar auroras.

Spectral images of the Jovian aurora obtained recently with STIS allow a quasi-instantaneous diagnostic of the energy of the precipitating electrons. First, CR maps derived from the two spectral observations of the Northern pole obtained on 16 and 24 Jan. 2013 through the GO12883 HST program revealed that the spatial distribution of the mean energy is far from uniform (Gustin et al. 2013, 2014). Several regions show little absorption (e.g. the Io footprint or the discontinuity region), while others are significantly attenuated (e.g. the main oval morning arc and flare emissions). A confined emission situated near the noon sector, also shows strong hydrocarbon attenuation. This emission, associated with the Ganymede footprint in both observations presented by Gustin et al. (2013, 2014) is now

identified as the small scale structure described by Palmaerts et al. (2014) in the 16 Jan. 2013
175 observation. Four more STIS spectral images obtained in 2014 (three of the Northern and one of the
Southern hemisphere) also revealed such spatial inhomogeneity of the energy distribution (Gérard et
al., 2014). These authors used mono-energetic precipitating electrons to derive the mean energy of the
auroral electrons. Depending on the region probed, the electron energy varies from a few keV up to 700
keV.

180 The two sets of observations used in the present study consist of a continuous time-tagged
spatial scan of the north Jovian polar region with the STIS 52"x0.5" slit in spectroscopic mode. This
procedure allows us to obtain reconstructed images of the aurora, where each pixel contains spectral
information and makes it possible to derive the CR and estimate the energy of the precipitating
electrons for each pixel. The present data thus exhibit a quasi-simultaneous picture of the whole
185 northern aurora, from which several regions may be extracted and compared. In particular, this dataset
allows for the first time the analysis of the 'small scale structure' and the Ganymede footprint auroral
emissions. The results described in this paper will bring new insight on auroral characteristics and will
be discussed within the frame of the current models describing the mechanisms responsible for the
auroral phenomenon at Jupiter. Section 2 presents the dataset. In section 3 and appendix A, we describe
190 the model that links the distribution of electrons precipitating at the top of the atmosphere to the CR. In
section 4, several representative auroral zones of the aurora are inspected, from which the energy-flux
relationship and the energy and brightness distributions are examined. In addition to the common
interpretation that assumes that UV aurora is primarily created by electrons, the hypothesis of UV
emission created by precipitation of oxygen ions has been explored. Section 5 summarizes our results
195 and provides the conclusions of this study.

2. Observations and data reduction.

This study is based on two STIS observations obtained on 16. Jan. and 24 Jan., 2013, operated in the long slit spectral mode, as part of the GO12883 HST program. The G140L grating was used along with the 52''x0.5'' aperture and the FUV-MAMA detector, providing spatially resolved spectra in the range 1150-1750 Å with ~12 Å spectral resolution. For MAMA first-order modes, only 25'' of the slit's length projects on the detector. The spectral range includes both absorbed and unabsorbed FUV H₂ emission, which is ideal to derive color ratios. The HST performed a continuous slew to move the slit projection from above the north limb to the equatorward edge of the aurora. The observations were designed to optimize the angle between the slit and the planetary equator in order to maximize the chances of intercepting the whole aurora during the scan. The two observations have an exposure time of 2528.2 seconds and were obtained in the time-tag mode, where the position and detection time of every photon is recorded in an event list with a 125 microsecond precision. In order to correctly identify each auroral feature and compute the χ angle (angle between the local normal and the observer) for each point, we reconstructed orthonormed images corresponding to chosen wavelength ranges based on the time-tagged event list. The orientation of the slit relative to the sky remained constant during each exposure. However, the motion of the center of slit relative to Jupiter involves a translation along a rotating axis as well as a compensation for the motion of Jupiter relative to the sky. In order to recompose orthonormed images with the Y-axis corresponding to the slit axis, it is necessary 1) to adjust the integration time corresponding to each row so that the angle covered by the motion of slit perpendicular to it equals the STIS platescale (i.e. 0.024'') and 2) shift this row to account for the motion of the center of the slit along an axis parallel to it. To do so, the motion of the slit relative to the sky is retrieved from the observational log file (the "jitter" file) and the motion of Jupiter is derived from the ephemeris. For each time step, a calibrated 2D spectrum is built out of the time-tagged event list using the standard CALSTIS pipeline. Then the brightness is integrated over the appropriate spectral window in order to produce a row on the orthonormed image. Once such an image is built, we

manually localize the X and Y position of IFP main spot. Since each row of the image corresponds to a different time, we compute the System III longitude of Io corresponding to this time and then the longitude and latitude of the IFP main spot, using the reference oval from Bonfond et al. (2009). Based on this information, it is then possible to infer the X and Y location of the center of Jupiter on the image. Such a location is crucial to build the polar projection map corresponding to each observation, as it allows us to provide a latitude and a System III longitude for each point on the image, acknowledging that the CML is different for each row since it varies with time. Unlike a true image where each pixel is simultaneously illuminated by the source, images reconstructed by this method should be called “pseudo-images”, as 42 minutes were necessary to perform the scan of the North pole. Figure 1 shows maps of the unabsorbed H₂ brightness of the aurora in the whole UV bandwidth (700-1700 Å) for the 16 Jan 2013 observation (Obs. #1, Fig. 1a & 1b) and 24 Jan 2013 (Obs. #2, Fig. 1c & 1d). These images were obtained by integrating the spectrum corresponding to each spatial pixel in the 1550-1620 Å bandwidth, unaffected by hydrocarbon absorption, and multiplied by 8.1 to extend the result over the whole UV. The latter factor has been determined from a synthetic H₂ spectrum as described by Gustin et al. (2004b). All brightnesses mentioned hereafter correspond to unabsorbed emission for the whole UV bandwidth. The two observations display very comparable features, which correspond to the different regions of the aurora described in the Introduction. These regions were manually selected from the images (red contours in Fig. 1) and are discussed in details throughout the paper. First, a very bright main arc in the morning sector (ME1), with a distribution of emission in the 50-600 kR range (centered at ~275 kR), evolving to a discontinuity between 150° and 185° S_{III} longitude, characterized by a very faint emission (mostly around 25 kR and < 60 kR). As the morning arc, the main emission in the afternoon sector (ME 2) is also very bright (in the 25-800 kR range centered around 240 kR), but is more spread out. Emissions inside the main oval are also present (FL1, FL2), taking the form of diffuse emissions, mostly in the range 30-300 kR. The Io footprint (IFP),

clearly visible in both images, is characterized by pixels of brightness between 40 and 300 kR, centered at ~ 165 kR. The small scale structure (SSS) is present on Obs. #1. It is situated within the discontinuity region (thus unaffected by the bright main emission) and exhibits very scattered brightnesses between ~ 20 and ~ 170 kR. The Ganymede footprint is visible on Obs. #2 and also show a wide range of brightnesses, from 40 to 240 kR. It should be stressed that these brightnesses correspond to observed values, and thus do not account for the effect of viewing angle. Signal from the pixels corresponding to these different regions has been extracted from the images as highlighted in Fig.1.

3. Mean energy of the primary electrons

3.1. Overview of the auroral model

The ratio between the images integrated in the 1550-1620 Å and the 1230-1300 Å ranges directly provides a map of the CR, which in turn can be converted to mean energy $\langle E \rangle$ of the precipitating electrons. The necessary prerequisite to build a map of $\langle E \rangle$ is to set up a model that relates CR to $\langle E \rangle$ and accounts for the emission angle χ . Indeed, at first order, the thickness of the hydrocarbon layer crossed by the auroral photons varies with $\cos(\chi)$ and influences the observed CR. The emission angle thus affects the amount of hydrocarbon absorption perceived by the observer, and hence the determination of $\langle E \rangle$.

Another aspect that is usually overlooked in such studies is the potential effect of the width D of the auroral curtain on the brightness and CR retrieved from the observations. These features are accounted for in our auroral model and thoroughly described in appendix A1. In brief, our model uses the atmosphere described in Grodent et al. (2001), which accounts for the higher gravity near the poles, the influence of the electron precipitation on the thermal structure, as well as several observational constraints such as the altitude of the H_2 emission peak, the temperature profile deduced from UV and

infrared emissions, and UV emission rates deduced from observed images and spectra. To relate $\langle E \rangle$ to CR, we used single Maxwellian distributions with mean energies from 2 to 1000 keV, representing a wide range of initial energy distributions of the precipitating electrons. An auroral volume emission rate vertical profile (VER: number of auroral photons emitted per second per cm^3) is then produced from each Maxwellian distribution with the two-stream Grodent et al. (2001) model. For each primary electron distribution, two synthetic emergent spectra are then calculated for emission angles χ from 0° to 80° and auroral widths of 150 km (representing the IFP, GFP and SSS), 600 km (representing the main emissions) and 1500 km (representing the flare emission in the polar cap region). A first synthetic spectrum, unaffected by the hydrocarbon attenuation is used to determine the influence of $\langle E \rangle$, D , and χ on the intrinsic auroral brightness while a second synthetic spectrum, absorbed by the hydrocarbons, is used to determine the influence of these parameters on the CR. Results from this modeling work are presented in Figure 11 in Appendix A1. It is seen that 1) for a given $\langle E \rangle$, the CR noticeably increases with χ , 2) for a given χ , the auroral width D has a minor effect on the CR, and 3) the unabsorbed brightness depends strongly on both χ and D .

Additional outputs from this model, described in appendix A2, focus on the characteristics of the VER profiles resulting from the different Maxwellian electron distributions. Results, summarized in Figure 12, show that 1) the altitude of the auroral peak decreases with increasing $\langle E \rangle$ and reaches a minimum value at ~ 150 km, 2) the FWHM of the VER profile decreases with increasing $\langle E \rangle$ (FWHM = 440 km for $\langle E \rangle = 2$ keV and 53 km for 500 keV), and 3) for a constant energy input of 1 mW m^{-2} , the emergent brightness strongly depends on $\langle E \rangle$, with a maximum value of 14.6 kR at $\langle E \rangle = 20$ keV and a decrease towards higher energies, reaching 3.2 kR at $\langle E \rangle = 500$ keV.

3.2. Important remarks regarding the energy – CR relation

To date, there is no direct observation that strongly constrains:

1. The temperature profile in the polar regions (i.e. the pressure-altitude relationship). The
295 knowledge of the H_2 density profile is crucial, as it provides the penetration depth of the precipitating
electrons for a given mean energy. For different temperature profiles, a given pressure level will
correspond to different altitudes and thus to different levels of absorption. Consequently, the use of, for
example, the Grodent et al. (2001) or the Moses et al. (2005) temperature profile does not yield the
same $\langle E \rangle$ -CR relationship. Also, the temperature profile affects the scale height of the emission and
300 thus the shape of the auroral vertical profile, which influences the portion of the aurora affected by
hydrocarbons.

2. The altitude of the hydrocarbon homopause, which influences the level of absorption for a
given altitude of auroral emission. In particular, the eddy diffusion coefficient K_h controls the
hydrocarbon mixing strength in the homosphere and thus plays an important role in the relative
305 hydrocarbon absorption of the auroral emission for a given energy distribution of electrons.

3. The energy distribution of the precipitating electrons, which strongly affects the shape and
altitude of the VER profile of the auroral emission. For example, a monokinetic beam of 100 keV
electrons produces a vertical H_2 emission with $CR=3.9$ while a Maxwellian distribution with $\langle E \rangle=100$
keV produces a vertical emission with $CR=3.0$. The type of electron energy distributions thus
310 influences the mean energy inferred from a given CR. In the case of kappa distributions, which are
controlled by two parameters (the most probable value E_0 and the spectra index k that controls the
amplitude of the distribution towards high energies), the $\langle E \rangle$ -CR relationship is not unique, as two
distributions with same $\langle E \rangle$ may lead to VER of different shape and thus of different color ratios.

As the CR - $\langle E \rangle$ relationship strongly depends on unconstrained auroral characteristics, it is
315 strongly model-dependent. In this work we chose the auroral atmosphere described by Grodent et al.
(2001), which is the most appropriate for auroral studies, and used single Maxwellian distributions to
create a $\langle E \rangle$ -CR relationship. The nominal eddy diffusion coefficient in Grodent et al. (2001) is

$K_h = 1.4 \times 10^6 \text{ cm}^2 \text{ s}^{-1}$ at the homopause. Assuming a doubling of the K_h coefficient, we estimate the error on the energy determinations as approximately 30 % of the nominal value. This value (30 %) should be
320 seen as a lower estimate of the total error, as other sources of uncertainty occur. These include errors propagated by data manipulations (such as background subtraction,...), or uncertainties on the calibration curves and various cross-section measurements needed in this study.

Constraints on the parameters just described could be provided by 1) solar or stellar occultations (for the temperature profile and pressure-altitude relationship), 2) direct observation of the potential
325 drop in the acceleration region and the electron energy distribution, and 3) direct observation of the altitudinal UV profile from images, from which the electron energy distribution could be inferred. The latter point is important, as VER profiles of the IFP have been observed from ACS images and analyzed by Bonfond et al. (2009). This special case will be discussed in section 4.3.4. The observations that should be made with the different instruments onboard the Juno spacecraft, from mid-2016 to late 2017
330 are expected to significantly contribute to these aspects of our understanding of the Jovian aurora.

In the following analysis, three different topics have been addressed. First, the auroral model briefly described in the preceding lines is directly applied to the spectral images to provide spatial maps of the electron energy. Several auroral regions have been delimited and results are discussed in section 4.1. Second, the energy - energy flux relationship for the different auroral regions is examined in order
335 to infer the processes leading to these emissions (section 4.2.). Third, the signal in the pixels forming the different auroral regions have been summed to provide corresponding average spectra. Similarly, new electron energy distributions have been determined for each region. This analysis is provided in section 4.3.

340 **4. Auroral characteristics inferred from observed spectral images**

4.1. Map of the auroral electron energy

The first application of the model described in section 3.1. and appendix A is the determination of the mean energy $\langle E \rangle$ of the precipitating electrons from the two STIS reconstructed images, whose spatial pixels contain spectral information. First, a map of the CR is obtained from the ratio between
 345 the images integrated in the (1550-1620) Å and (1230-1300) Å bandwidths. Second, a map of angle χ is calculated for each spatial pixel from latitude and longitude grids as described in section 2. Values of CR and χ are thus associated with each spatial pixel, from which $\langle E \rangle$ is determined, based on interpolation of the CR- $\langle E \rangle$ relationship obtained at various χ angles (see Fig. 11e). Figure 2 shows the map of the precipitating electron mean energies $\langle E \rangle$ for the two observations. The different auroral
 350 regions defined in Fig.1 are also outlined in Fig. 2, and each region defined in these figures is discussed in more detail in the following sections. First, a clearer view of the dispersion of the brightness and mean energies within each region is obtained by grouping the brightness and $\langle E \rangle$ of each pixel by bins of 5 or 10 kR (or keV), forming histograms of the vertical brightness and mean energy, as shown in Fig 3 and Table 1. Each point included in these histograms corresponds to the brightness and mean energy
 355 of a spatial pixel of the auroral regions considered. In order to remove the effects of viewing angle, usually not considered in auroral models, the observed brightnesses have been converted to vertical brightness for each pixel, following the curve plotted in Fig. 11c. The moderate influence of $\langle E \rangle$ on the conversion has also been taken into account in the calculations. For the main emissions (ME region 1 and ME region 2) and discontinuity (DC, region 5), we assume an auroral width of 600 km. For the
 360 flares (FL, region 3 and 4) we assume a width of 1500 km. A width of 150 km is then assumed for the Io footprint (IFP, region 6), the small scale structure (SSS, region 7 in Obs #1) and the Ganymede footprint (GFP, region 7 in Obs #2).

In order to empirically determine the brightness and mean energy distribution associated with each auroral region, both the vertical brightness and $\langle E \rangle$ histograms have been fitted with two distributions
 365 often used in this field of study. First, a Maxwellian distribution,

$$D = C \frac{E}{E_0} e^{-E/E_0} \quad (1)$$

and a kappa distribution

$$D = C \frac{E}{E_0} \left(1 + E \frac{E_0}{k} \right) e^{(-1-k)} \quad (2)$$

A third distribution, allowing a greater flexibility in terms of shape, has also been used, namely the
 370 'generalized' Maxwellian distribution

$$D = C E^{n/2} E_0^{-(\frac{n}{2}+1)} e^{-E/E_0} \quad (3)$$

In these equations, C is a constant factor, E is the electron energy (or the brightness), E_0 is the most probable value for the Maxwellian and kappa distributions (i.e. the characteristic energy), while the most probable value is $nE_0/2$ for the generalized Maxwellian. Parameter k is the spectral index of the
 375 kappa distribution and n is a free parameter of the generalized Maxwellian distribution. This distribution is Maxwellian for $n=1$ and becomes strongly non-Maxwellian for high values of n . More details on this distribution may be found in Dzifcakova (1997) and references therein. Figure 3 shows that the kappa and Maxwellian distributions do not provide good fits to either the brightness or energy distributions. Instead, the generalized Maxwellian provides the best agreement with the observed
 380 profiles, usually with strong non-Maxwellian characteristics ($n \gg 1$). Interestingly, the brightness and energy distribution of each auroral zone share very similar characteristics for both observations. The E_0 and n parameters of the best generalized Maxwellian are provided in Table 1.

As seen in Fig. 3 a,b,o,p, the ME 1 region is very bright and characterized by the most energetic primary electrons. The vertical brightness distributions are relatively symmetric, around ~260 kR. The
 385 electron mean energies $\langle E \rangle$ determined from the CR method are mostly between 100 and 700 keV, with several individual pixels at higher or lower values. The average $\langle E \rangle$ value is quite high (~300 and ~355 keV for observation 1 and 2, respectively). It should be noted that mean energies found in the literature are generally determined from spectra obtained from numerous pixels, which averages the energies so obtained, while the distributions shown here reflect the range of values reached by the
 17

390 individual pixels of the auroral region, which expand the range of inferred values. The Maxwellian and
 kappa distributions do not satisfactorily fit the brightness and $\langle E \rangle$ distributions. A very good fit is
 obtained with the generalized Maxwellian distributions. Figure 3o, p clearly show that several pixels
 have $\langle E \rangle$ larger than 400 keV, well above values generally observed. As seen in Fig. 11a, for $\langle E \rangle$
 larger than 400 keV, the CR is larger than 20 for a viewing angle of 70° . For a pixel with a total
 395 unabsorbed UV brightness of 200 kR and CR=20, the (1550-1620 Å) unabsorbed intensity is 25 kR,
 while the absorbed (1230-1300 Å) region is 20 times less, i.e. 1.25 kR. In other words, pixels
 associated with very high electron energy undergo a very low S/N in the absorbed spectral bandwidth,
 which induces a very high uncertainty on the energies inferred for high energy pixels of the
 distributions. As a result of this large uncertainty, pixels with $\langle E \rangle$ larger than 400 keV should be
 400 considered with caution. A way to overcome this problem is to bin the pixels in groups or sum all the
 pixels of a given region, to increase the S/N ratio, as proposed in the sections 4.2.1. and 4.3.1. The
 mean energy map (Fig. 2) and distributions (Fig. 3) should then be seen as first order estimations.

The bright winding afternoon main emission (ME2 known as the “kink” region, see Grodent et al.
 (2003) and Ballester et al. (1996)) shows a very asymmetrical vertical brightness distribution, with an
 405 average value around 160 kR, reaching values above 400 kR for several pixels. Although the mean
 energies are large (~ 130 keV), $\langle E \rangle$ in ME2 are roughly 2.5 times lower than ME1 values. The
 distributions are still best fitted by the generalized Maxwellian function.

The vertical brightness distributions from the four polar cap emissions selected in region 3 and
 4 (FL1 and FL2) are very comparable in shape and value, with peaks between 40 and 70 kR. The mean
 410 energies are significantly lower than in the main emission cases, with averaged values between ~ 65 and
 ~ 100 keV. Again, the departure from a Maxwellian distributions is clear. The $\langle E \rangle$ distributions are
 relatively narrow with FWHM ~ 80 keV, and with virtually no electrons with energies > 200 keV.

As expected, the discontinuity region exhibits very weak emission, with an average at ~ 30 kR.

The three tested distributions provide a reasonable fit to the brightness histogram, again with a best
415 model provided by the generalized Maxwellian. In terms of mean energies, a substantial number of
individual pixels (~40%) have a total UV intensity lower than 25 kR (thus low S/N), which does not
allow a safe determination of $\langle E \rangle$. We thus estimate that the energy distributions from individual pixels
in the discontinuity region are not trustworthy and are disregarded from this analysis. A precise
determination of $\langle E \rangle$ will be discussed in the next section, based on the spectra obtained from the
420 summed pixels of this region.

The number of pixels belonging to the Io footprint is relatively small (198 for observation #1
and 138 for observation #2), which makes it difficult to derive statistically significant distributions. In
addition, the CR method is probably not appropriate to determine $\langle E \rangle$ for each pixel, because 1) the
CR from our data is always close to the unabsorbed value, which makes $\langle E \rangle$ more sensitive to S/N
425 ratio, and 2) the Maxwellian distributions used in the CR method to link $\langle E \rangle$ to CR are probably not
appropriate in the IFP case. Indeed, from several IFP emission profiles determined from ACS images,
Bonfond et al. (2009) determined an average VER consistent with precipitating electrons in a kappa
distribution instead of a Maxwellian, which modifies the CR- $\langle E \rangle$ relationship (see paragraph 3.2.).
This point will be discussed in more details in the spectral analysis (section 4.3.4.). Consequently, the
430 distribution of the pixels per energy bin cannot be determined for the IFP. Our sample shows observed
brightnesses from 15 to 420 kR. Due to the effect of viewing angle and auroral width (assumed to be
150 km for footprints in the model), these observed values should be converted to vertical values.
Again, this conversion is not possible because of the unknown value of each pixel's mean energy. We
thus decided to show in Figure 3 the observed IFP brightness distribution instead of the vertical value.
435 Because of the very narrow range of viewing angles associated with the IFP spot, the shape of the
vertical and observed brightness distributions are virtually identical. The only difference is the absolute
value of the brightness, which thus shifts the whole brightness distribution accordingly. The best

Maxwellian electron distribution used by Bonfond et al (2009) to fit the average IFP profile has a mean energy of 2 keV. Although this Maxwellian distribution do not provide the best fit the IFP profile, it
440 allows us to estimate the observed-to-vertical conversion factor from our model. (With such energy and a viewing angle of 80° , the observed brightness must be multiplied by 7 to represent the vertical value, which is consistent with vertical conversion factors determined by Bonfond et al. (2013) from 3D simulations. This quite big conversion factor is explained by the very limited extent of the intrinsic auroral emission, which thus needs to be very intense to provide the observed value. The vertical
445 brightnesses of our sample thus lie between ~ 100 and ~ 3000 kR.

The small scale structure (SSS) is also clearly seen in the 16 Jan. 2013 observation (region 7 in Fig.1 a & b). Since it is situated inside the discontinuity region, we specifically used the discontinuity as background for the SSS in order to withdraw the potential influences of the DC on the SSS emission. Because of the relatively low number of pixels in the SSS region (~ 100 , to be compared with values
450 between 850 and 2400 for the main emission and flare regions), the brightness distributions, spread over the 30 - 400 kR range, do not reveal a clear shape. We assumed an auroral curtain of width 150 km to convert observed to vertical brightnesses, which corresponds to a factor of ~ 2 for $\langle E \rangle = 200$ keV. As for the IFP, the low number of pixels considered does not allow us to present a reliable distribution of mean energies. Still, a first order examination of the mean energy distributions shows that the average
455 $\langle E \rangle$ is surprisingly large, at ~ 260 keV. Again, a more precise examination of SSS characteristics, based on the spectra obtained from the sum of all SSS pixels, is discussed in section 4.3.5. Characteristics of the GFP emission (region 7 in Fig. 1 c&d) are very close to those of the SSS. The low number of pixels defining the GFP makes it difficult to infer a mean energy from the distribution, but the average value is around 120 keV.

460

4.2. Mean electron energy – energy flux relationship

4.2.1. Determination of the data points.

After the first-order overview based on the STIS maps (Fig. 2) and individual pixel distributions (Fig. 3), a more accurate examination of each region is given in this section. As already mentioned, $\langle E \rangle$ may be loosely determined when the individual pixels exhibit a low count rate (hence low S/N), which consequently increases the uncertainty of the $\langle E \rangle$ retrievals. To improve the accuracy of the mean energy and energy flux values (and thus avoid large scatter of the data points in a mean electron energy – energy flux diagram), we sorted the pixels belonging to each zone from lowest to highest brightness, and binned them by groups of eight pixels. We thus implicitly assume that there is an unequivocal relationship between the energy flux and the mean energy. A higher S/N spectrum is derived from each new point, from which a new CR hence $\langle E \rangle$ is determined using the CR method described in the previous section. The energy flux is determined for each new point following the relationship illustrated in Fig. 12d, where the energy flux - brightness conversion is energy-dependent. This method is much better than a simple smoothing or binning of the original data points, as it removes potential unreliable values from original low S/N pixels. As expected, the mean energy – energy flux diagrams obtained from the binned data points conserve the trend seen from the original points, but with much less dispersion. Very high localized mean energies seen in some distributions in the previous section (for example, $\langle E \rangle$ above 400 keV for ME1 in Fig. 3a) disappear in the newly binned energy data points.

The main remaining source of uncertainty in the mean electron energy determination with the CR method stems from the hydrocarbon density profiles, not constrained by observations in the polar regions. As described by Gustin et al. (2004a), we determined the uncertainty in $\langle E \rangle$ by considering the effect of doubling the eddy diffusion coefficient K_h , which increases the homopause altitude. This coefficient is $1.4 \times 10^6 \text{ cm}^2\text{s}^{-1}$ in the nominal atmospheric model. This variation of the homopause altitude has a non-linear effect on the CR and is thus energy dependent, which implies a ~12% to 30%

uncertainty in $\langle E \rangle$. The uncertainty on the energy flux comes from the S/N ratio (error between ~5 and ~25 %) and the systematic uncertainty due to the kR – mW m⁻² conversion factor, which is model dependent. At 100 keV, the latter factor may vary from 8.4 to 10.6 kR per injected mW m⁻² depending on the atmospheric model and method chosen, which represents a constant 22% uncertainty. The error
 490 in the energy flux thus varies from ~23 to ~33%.

4.2.2. Theory

The relationship between the energy fluxes and precipitation energies inferred from the data can be compared to theoretical predictions for static auroral acceleration. Based on Knight's (1973) kinetic
 495 theory of field-aligned currents, Lundin & Sandahl (1978) first derived the relationship between the incident energy flux at the atmosphere and the energy of electrons precipitating through a steady-state, monotonic field-aligned potential drop. The location and extent of the field-aligned potential was specified by the mirror ratio, $R_x = B_i/B_x$, between the top of the acceleration region, denoted by the subscript x , and the atmosphere, denoted by the subscript i . An isotropic, Maxwellian distribution of
 500 electrons was assumed at the top of the acceleration region. The precipitating electron energy flux is then described by

$$E_{fi} = n_x \left(\frac{W_x}{2\pi m_e} \right)^{1/2} \left[R_x (2W_x + e\Phi_{||}) - (R_x e\Phi_{||} + 2(R_x - 1)W_x) \exp \left(-\frac{e\Phi_{||}}{W_x(R_x - 1)} \right) \right], \quad (4)$$

505 where W_x is the electron temperature (in energy units) at the top of the acceleration region, $\Phi_{||}$ is the magnitude of the potential drop, m_e is the mass of the electron, n_x is the electron density at the top of the acceleration region, and e is the fundamental charge of the electron. Lundin & Sandahl (1978) also showed that in the limit where $e\Phi_{||} \gg W_x$ and $B_i \gg B_x$, the relationship can be simplified to

$$E_{fi} = n_x \sqrt{\frac{1}{2\pi m_e}} W_x^{-1/2} (e\Phi_{||})^2. \quad (5)$$

Physically, this limit describes an accelerating potential in which the energy gained by the electron is far greater than its initial energy, and the extent of the potential drop is such that an infinite reservoir of electrons can be assumed at the top of the acceleration region.

Jupiter's main auroral emission is, to first order, a steady-state phenomenon that is driven by the outward transport of Iogenic plasma within the magnetosphere and fixed in the frame rotating with the planet. Therefore, the steady-state relationship between the energy flux and field-aligned potential strength described by equation (4) is applicable and, when combined with the mean energy and energy flux of the precipitating electrons inferred from STIS observations, can offer insight into the properties of the auroral acceleration region and the source population of the precipitating electrons. The location of the acceleration region, and hence whether or not equation (5) is applicable for the Jovian system has not yet been determined. This has been discussed in detail (e.g. Cowley & Bunce (2003), Su et al. (2003), Nichols and Cowley (2005), Ray et al. (2009, 2010)) and will be explored further in a companion paper (Ray et al. in preparation).

The Lundin & Sandahl (1978) analysis was for the terrestrial system, where the precipitating electron energies are on the order of 10s of keV. At Jupiter, the inferred precipitating energies are much larger, sometimes reaching 500 keV. At these magnitudes, the precipitating electrons are relativistic. Cowley [2006] derived current-voltage and energy flux relations for an isotropic Maxwellian distribution of electrons accelerated by field-aligned potentials to relativistic speeds. The kinetic energy flux deposited in the planetary atmosphere by relativistic electrons can exceed that estimated by the non-relativistic relation by a factor of 2 (for $e\Phi_{||}/kW_x = 20$, $W_x = 25$ keV) to 5 (for $e\Phi_{||}/kW_x = 200$, $W_x = 25$ keV). Therefore, the estimates of the precipitating electron density into the atmosphere will also be modified for a given electron energy flux and precipitating electron energy.

4.2.3. Analysis

535 Figure 4a shows the energy flux – precipitation energy relationships from the main emission and flare regions of the two STIS observations. What is evident is that the range of energy fluxes and mean energies for the flare (black and orange stars) and main emission 2 (green stars) regions are coincident, while the mean energies estimated for the main emission 1 regions (light blue stars) are a factor of 2 – 3 larger. First, we use equation (5) to estimate the density of the ‘source’ magnetospheric electron
540 population, as in Gustin et al. (2004a). Figure 4b – g show the individual ME and FL auroral regions, with the dashed and dashed-dot lines corresponding to the energy flux – mean energy curves for $W_x = 2.5$ keV and densities of $n_x = 1000 \text{ m}^{-3}$ and $n_x = 10000 \text{ m}^{-3}$, respectively. The latter values correspond to the hot electron density range measured near the current sheet by Voyager 1 and 2 (Scudder et al. 1981).

545 We first estimate the density of the electrons in the middle magnetosphere plasma sheet by determining the best fit to the data using equation (5) and $W_x = 2.5$ keV (red and green lines in Fig. 4). The inferred precipitating electron densities for the ME2 (kink) region are $\sim 2400 \text{ m}^{-3}$, consistent with Voyager values. The density estimates from the narrow ME1 emission are a factor of 3 lower ($\sim 800 \text{ m}^{-3}$). However, these emissions map to the dawn or possibly midnight sector of the magnetosphere, which
550 Voyager did not sample. For ME1, the relativistic energy flux – mean energy relationship has also been tested, as the electron’s acceleration is a significant fraction of the electron rest energy for this region (panel 4 d & e). Compared to the non-relativistic case, electron densities are $\sim 45\%$ lower, with values of 619 and 521 m^{-3} . These values correspond well with densities obtained from the study of a remarkably bright auroral morning arc observed by STIS on 21 September 1999, which also exhibit
555 very high electron energies (Gustin et al. 2006). It should be noted that the 2.5 keV electron temperature we use is the center energy of the narrow 2-3 keV range quoted by Scudder et al. (1981).

Globally, an equivalent Knight curve is obtained from an increase (decrease) of 0.5 keV of the electron temperature associated with an increase (decrease) of the density by ~10 %.

Unlike the relatively well-understood main aurora emission, it is less clear what acceleration processes drive the variable flare emissions, which map to the outer magnetosphere, or possibly to the boundary region or beyond. Therefore, the energy – energy flux relationship of the flare regions shown in Fig. 4a are not further discussed in the present paper but will be examined in Ray et al., in preparation

4.3. Spectral analysis

4.3.1. Mean energy from average spectra

In order to maximize the S/N of the spectral information at each auroral region defined in Fig. 1, an average spectrum has been constructed for each region by summing the spectra associated with the spatial pixels defining each auroral region. Since each spectrum may correspond to a region that covers substantial latitude and longitude ranges, an average viewing angle has been determined in each case. In contrast to the average background used for the $\langle E \rangle$ map production shown in Fig. 2, a specific background, chosen close to the auroral region considered, has been used and subtracted in each case. The mean energy of the extracted spectra, determined from the CR method, and the unabsorbed UV brightness in the 700-1700 Å range are summarized in Table 1. For a given region, values from the average spectrum and the distributions described in paragraph 4.1. are slightly different. This is explained by several factors, such as the different background subtraction, the average viewing angle associated with each spectrum, the lower S/N ratio of the individual pixels used to determine the mean energy of each pixel, etc.

4.3.2. Can the energy distribution be inferred from spectral analysis?

As mentioned before, since no observation provides sufficient constraints on the energy distribution of the precipitating electrons and on the auroral VER profiles, the $\langle E \rangle$ -CR relationship is not unequivocal: different electron energy distributions with the same $\langle E \rangle$ will produce emergent spectra with different color ratios. Our CR method includes the contributions of CH_4 , C_2H_2 and C_2H_6 to
585 attenuate the auroral signal, but only accounts for the absorbed 1230-1300 Å spectral range. As we reach deeper atmospheric layers (~250 km), the density profiles of acetylene and ethane become very important and affect the shape of the auroral spectra, especially in the 1330-1530 Å spectral regions, not used in the CR definition. Indeed, unlike CH_4 , density profiles of C_2H_2 and C_2H_6 do not monotonically increase with decreasing altitude, instead presenting density peaks (see Fig. 9c). Since
590 the amount of absorption depends on the altitude of the emission but also on the shape of the VER profile, specific absorption of C_2H_2 and C_2H_6 may bring new constraints on the VER. Here we thus test the possibility of determining a VER profile that would provide a fit to a given FUV spectrum that is significantly better than any other VER. This preferred VER could then be used to infer a ‘best’ electron energy distribution.

595 We use synthetic spectra produced by the method described in appendix A1 to fit the observed spectra. At ~12 Å spectral resolution, the specific signature of each hydrocarbon can be easily identified. We used Chapman profiles to simulate the VER, using the parameters described by equation (10) in Appendix A2 as free parameters. The unabsorbed H_2 spectrum generated at each altitude bin of the synthetic VER is absorbed by the overlying hydrocarbon layer, and a total emergent (absorbed)
600 spectrum is calculated using the atmospheric model described in section 3.1. and appendix A1, taking into account the finite width of the auroral curtain and the mean viewing angle of the auroral region considered. The synthetic emergent spectrum is compared to the observed spectrum and the VER parameters varied until the difference between the two spectra reaches a fit minimum chi-square. This method thus presumes that a unique VER profile provides a spectral fit that is significantly better than

605 any other synthetic VER profiles. This method has been tested on the spectra of all auroral regions and
 is shown here for the STIS ME 1 spectrum of obs. #1, which is strongly attenuated by hydrocarbons
 (CR=13.6). This strong attenuation implies that the auroral emission reaches altitudes where C_2H_2 and
 C_2H_6 number densities are important enough to show clear signatures of absorption. Results from this
 spectral analysis demonstrate that several synthetic VER provide synthetic spectra that satisfactorily fit
 610 the ME 1 spectrum, which makes it impossible to unequivocally determine electron characteristics
 from spectral analysis. Figure 5 shows two simulations (plain red and green curves) where the synthetic
 VERs, although quite different, lead to very similar fits to ME 1. As seen in Fig. 5a, the VER profile
 from fit 1 (red) is more intense, peaks at lower altitude, and has a smaller FWHM than VER from fit 2
 (green). The corresponding emergent spectra (red and green curves in Fig. 5b and c), are difficult to
 615 distinguish from the data and do not allow one to select a preferred model. The light blue dashed curve
 in Fig. 5 b & c show the unabsorbed emergent spectrum and gives an idea of the hydrocarbon
 attenuation. It should be noted that a one-layer model (an emitted layer overlaid by an absorbing layer)
 was tested too and never provided a satisfactory fit to the data, thus supporting a more complex
 approach such as the one described here. We used Fig. 12a to determine $\langle E \rangle$ from the peak altitude of
 620 the synthetic VER. The best VER 1 peaks at 165 km, which corresponds to 375 keV. By contrast, VER
 2 peaks at 195 km and corresponds to $\langle E \rangle = 188$ keV. These two energies thus lie either side of the 278
 keV value obtained from the CR method. This example demonstrates that the $\langle E \rangle$ value and thus the
 electron distribution cannot be determined from a spectral analysis without additional constraints. The
 CR method thus provides a fair estimate of the energy of the electrons responsible for the auroral
 625 emission, under the limits provided by the hypotheses made to define it.

4.3.3. Electron energy distribution deduced from observations

We thus adopted another strategy in order to estimate the primary electron energy distributions

from the observations. As described in section 4.2.1., the energy flux E_f and mean energy $\langle E \rangle$ have
630 been determined from the brightness and CR of each binned pixel of several auroral regions of the
STIS observations. Panel a of Fig. 4 clearly demonstrates that the E_f and $\langle E \rangle$ are 1) very similar for a
given auroral region and 2) very different for different auroral regions. This characteristic can be used
as a constraint to empirically determine the energy distribution of the primary electrons of each auroral
region. Assuming that the emission at each binned pixel i of a given region is due to electrons in a
635 Maxwellian distribution of mean energy $\langle E(i) \rangle$ and energy flux $E_f(i)$, the primary electron distribution
that best describes on average this region is thus the sum of the $\langle E(i) \rangle$ Maxwellians, weighted by the
 $E_f(i)$.

The average primary electron energy distributions obtained from this procedure are shown in
Figure 6 for ME 1, ME 2, and FL regions for the two STIS observations. Distributions are not provided
640 for the DC (because of very weak brightness, $\langle E \rangle$ cannot be safely determined for each pixel), IFP
(because absorption when present, is at the limit of detection, which does not permit an accurate
estimate of $\langle E \rangle$), and SSS (because of the low number of scattered points in the distribution) regions.

To provide an analytical form of these constrained distributions, they have been fitted with
Maxwellian, kappa and generalized Maxwellian distributions as described by equations 1-3. As seen in
645 Fig. 6, each auroral region in both STIS observations has very similar electron energy distribution –
and thus shares specific characteristics, as already mentioned. At the exception of the ME1 region
whose constrained electron distribution can be well fitted by a Maxwellian distribution, kappa
distributions with a spectral index between 3 and 9 usually provide better fits to the constrained
distribution than Maxwellians do for ME2 and FL regions. Indeed, the high energy 'tail' observed in the
650 constrained distributions is better reproduced by the slower decrease of the kappas with energy,
compared to the Maxwellians. Overall, the generalized Maxwellian functions provide the best fit in
each case. The main characteristics of these distributions are provided in Table 2. As expected, the

mean energy of the distributions is close to the mean energy determined from the spectra with the CR method shown in Table 1. The VER associated with the constrained distribution corresponding to region ME 1 obs. #1 (shown in dashed dark blue in Fig. 5a) has been used to generate an emergent synthetic H_2 spectrum. This modeled spectrum compares very well to the observed spectrum. In particular, all the spectral features influenced by the CH_4 , C_2H_2 and C_2H_6 absorptions are nicely reproduced by the model, which validates the method performed to derive these constrained primary electron distributions and shows that the CH_4 , C_2H_2 and C_2H_6 density profiles adopted in our model are appropriate to describe the auroral atmosphere.

4.3.4. The Io footprint: a special case

In contrast to the main oval and polar cap emissions, a determination of $\langle E \rangle$ from hydrocarbon absorption is difficult for the IFP because of its weak attenuation. In theory, at a viewing angle of 70° , the CR is 1.2 for $\langle E \rangle = 20$ keV and 1.5 for $\langle E \rangle = 30$ keV. A 50% increase of the mean energy thus only changes the CR by 0.3. This small CR increase is below the limit of accuracy reached by spectra associated with the single IFP pixels of the spectral images. Deriving an average electron energy distribution from these individual pixels is thus not possible in this case. However, several comments can be addressed from the analysis of the IFP spectra obtained from the sum of all the IFP pixels.

1. Based on the CR method, the CR of the summed spectra pixels suggest an electron mean energy of ~ 20 keV for both IFP observations. Figure 7a shows an emission profile (VER 1, orange line) determined from a Chapman profile with free parameters that fits well the IFP spectrum of the 16 Jan. 2013 STIS observation (quasi-identical to the 24 Jan. observation). Its characteristics correspond well to VERs obtained from a Maxwellian electron distribution. It has a narrow shape (FWHM=125 km), peaks at 330 km and has a CR of 1.24, compared to 1.28 for the observation. As seen in Fig. 12a, this altitude peak corresponds to a mean energy of 17 keV, in agreement with the CR method, accounting

for the uncertainties.

2. Although a constraint on the electron distribution cannot be derived, several other parameters may be constrained. First, observed emission profiles of the IFP has been observed from ACS images
680 by Bonfond et al. (2009). These authors selected 38 observed VERs from the IFP where the auroral curtain is quasi perpendicular to the observer in order to generate a 'typical' emission profile with good S/N ratio. This emission was best fitted with a modeled VER profile obtained from a kappa distribution of primary electrons with a spectral index $k=2.3$, $E_0=70$ eV and mean energy = 1.1 keV. This modeled VER peaks at ~900 km and has a FWHM of ~1200 km, much larger than the values attained by VER
685 due to Maxwellian distributions (the maximum FWHM of the VER is ~440 km, from a 2 keV Maxwellian electron distributions, as seen Fig. 12c). The VER profile best fitting the ACS observations is plotted in blue in Fig. 7a and is used to constrain the shape of the VER used in our model. Second, examination of the full ACS database reveals that the peak altitude of the IFP emission may be quite variable, with values from ~ 550 to ~1300 km, which constrains the range of altitudes allowed for the
690 modeled emission peak. A third constraint arises from the minimum altitude reached by the precipitating electrons. As shown in Fig.12a, the peak altitude of the VER is ~ 164 km for a Maxwellian distribution of 500 keV electrons, with a corresponding FWHM of 53 km. If we assume mono-energetic electrons of 500 keV, these electrons are stopped by a total H_2 column of $2.3 \times 10^{22} \text{ cm}^{-2}$, which corresponds to an altitude of 148 km in the atmospheric model used here. Consequently, there
695 is no significant emission at altitudes below ~150 km, stemming from primary electrons precipitating in the atmosphere. The shape of the synthetic VER must then be adapted in order to eliminate all emission below ~150 km. From these constraints, we built a synthetic VER from a Chapman profile, whose shape corresponds to the modeled VER determined by Bonfond et al. (2009) that is allowed to peak in the 500-1300 km range. The modeled emission profile is forced to decrease at low altitude in
700 order to eliminate all emissions below 150 km. The peak altitude and maximum value of the VER

profile are the only free parameters of the model from which synthetic spectra are produced until a synthetic spectrum best fitting the observation is obtained. A best emission profile, VER 2, that provides a good fit to the spectrum is shown in green in Fig 7a. The shape of this VER is significantly thinner than the Bonfond et al. (2009) observations because of the limited emission allowed to emerge at low altitudes, as just explained. The maximum emission arises at an altitude ~ 300 km, significantly lower than the minimum peak altitude of ~ 550 km observed by Bonfond et al. (2009). The peak altitude and shape of VER 2 thus make this solution very unlikely, as they do not meet the constraints provided by the observed emission profiles.

3. The only way to derive an absorbed spectrum with a “thick” VER peaking at higher altitude is to change the altitude of the homopause level relative to the auroral emission and allow an upwelling/vertical transport of the hydrocarbon layer. In addition to the peak altitude and maximum value of the synthetic VER emission, we thus added a third free parameter in the model, i.e. an upward shift of the hydrocarbon density profiles with respect to the H_2 density profile in the atmospheric model. Several fits with different shifts of the hydrocarbon homopause provided comparably good fits to the observed IFP spectrum as shown in Fig. 7c (stars). It is seen that the peak altitude of the best VERs varies linearly with the upwelling, which is expected as the absorption depends on the relative position of the VER with the hydrocarbon profiles. Interestingly, a minimum upwelling of 70 km is required to obtain a VER peaking at 550 km. Assuming that a redistribution of hydrocarbons at higher altitude is exclusively due to a change of the eddy diffusion coefficient K_h , a 70 km upward shift of the homopause corresponds to $K_h = 1.4 \times 10^7 \text{ cm}^2 \text{ s}^{-1}$, i.e. 10 times the nominal value used in the Grodent et al. (2001) atmosphere. However, the association of an auroral peak at 550 km with a CR of 1.3 is very unlikely, as 550 km is already the lowest altitude observed from the ACS images, and 1.3 is in the lower range of CR values determined so far from IFP spectra (CR from 1.44 to 2.28 where obtained by Gérard et al. (2002) from STIS spectra). Assuming that the relative uplift of hydrocarbon is exclusively

725 due to a variation of K_h , a maximum homopause shift can be estimated. Results from the Grodent et al. (2001) model show that the homopause altitude varies linearly with K_h in log scale (stars in Fig. 7d). On the other hand, Parkinson et al. (2006) analyzed UVIS observation of the He 584 Å emission during the Cassini flyby of Jupiter and determined that K_h at the homopause in the auroral region is 'possibly greater' than $4 \times 10^7 \text{ cm}^2 \text{ s}^{-1}$. Extrapolation of the relationship presented in Fig 7d to $K_h = 4 \times 10^7 \text{ cm}^2 \text{ s}^{-1}$ 730 leads to a homopause shift of 100 km. By fixing this value in our fitting procedure, we found a VER best fitting the IFP spectrum peaking at 580 km (red curve in Fig 7 a and b, diamond in Fig. 7 c), which is still in the very low range of altitudes determined by Bonfond et al. (2009).

4. To obtain an IFP emission peaking at a 'standard' altitude of ~ 900 km that generates an absorbed emission, an homopause upward shift of 280 km is necessary. Based on the extrapolation shown in Fig. 7d, this corresponds to an increase of K_h to $2 \times 10^{11} \text{ cm}^2 \text{ s}^{-1}$, i.e. five orders of magnitude above the 735 nominal value in the Grodent et al. (2001) model, which seems very unlikely as well. However, an increase of turbulent diffusion is not the only way to change the hydrocarbon vertical distribution. Vertical wind transport generated by the auroral precipitation may carry hydrocarbon molecules to altitudes higher than the expected homopause level, thus affecting the absorption of the emergent 740 emission (see discussion in Bonfond et al. (2009)).

5. It is quite obvious that the emission profiles presented in Fig. 7a cannot result from Maxwellian electron distributions. Except for the first case discussed (VER 1 compatible with a Maxwellian distribution, orange curve in Fig. 7a), the other emission profiles show a much larger scale height. Thanks to the two free parameters of the kappa distribution, both the shape and altitude of the profile 745 can be adjusted by varying E_0 and the spectral index k . Since several (VER profile, homopause shift) configurations may simulate the IFP emission in the framework of the broad constraints just explained, the electron characteristics determined by Bonfond et al. (2009) cannot be updated here. Still, as a rule of thumb, one can consider that an increase of E_0 with constant k will favor emission at lower altitude

stemming from a shift of the electron energy toward larger values, and thus shift the whole emission
750 profile towards lower altitudes. Keeping E_0 constant and decreasing k (i.e. increasing the contribution
of high energy electrons in the 'tail' of the distribution) drastically changes the shape of the distribution.
The low energy electrons conserved by keeping the value of E_0 constant will keep producing
significant emission at high altitudes, while the energetic electrons in the tail of the distribution will
increase emissions at low altitude and provide a profile with a shape like VER 2 in Fig. 7a.

755 In summary:

- 1) With the hydrocarbon distribution in the nominal atmospheric model of Grodent et al. (2001), no
emission profile can provide at the same time an absorbed emergent spectrum (as constrained by STIS
spectra) and VER characteristics consistent with the ACS constraints.
- 2) The only way to reconcile the constraints of absorption and the observed profiles is to significantly
760 shift upward the altitude of the homopause level, by at least 100 km.
- 3) The CR method used so far to determine $\langle E \rangle$ is not appropriate for the IFP emission, because 1) the
FWHM of the VER profiles obtained from monokinetic or Maxwellian electron distribution is not
consistent with the much 'thicker' observed profiles of the IFP emission (for a given $\langle E \rangle$, different
energy distributions will provide different VER profiles and thus different absorption levels), and 2) the
765 altitude of the emission peak obtained from the CR method are significantly lower than the range of
values observed by ACS observations. Consequently, $\langle E \rangle$ determined by the CR method (40 to 70 keV
in Gérard et al. (2002) and up to 100 keV in Gérard et al. (2014) are significantly overestimated.
Simulations have been performed to determine the $\langle E \rangle$ - CR relationship with the procedure and
atmosphere described in appendix A1, assuming a 100 km upward shift of the homopause level. With
770 such a modified atmosphere, mean energies are, roughly, lowered by a factor of 3 for a given CR. In
particular, the energies between 40 and 70 keV determined from IFP spectra in Gérard et al. (2002)
now range between 13 and 23 keV.

4) Processes that could lead to such an important upward shift of the hydrocarbon layer with respect to the auroral emission are not well understood. Contrary to the main emission, which is the result of precipitating electrons accelerated by a quasi-static potential, acceleration of electrons by Alfvén waves are involved in the IFP emission (Bonfond et al. (2008), Hess et al. (2010)). This difference could be a clue implying that this uplift of the hydrocarbon layer applies to the IFP region only. In terms of the characteristic of the precipitating electrons, the main difference between the Io and main oval emissions is the current density. The energy input is very similar in both cases, but the mean energy of the electrons lies in the range 50-250 keV for the polar emission and around 1 keV for the IFP emission. This difference directly induces a current density that is ~20 to ~250 times higher in the case of the IFP. For example, an energy input of 10 mW m^{-2} leads to a current density of $0.1 \text{ } \mu\text{A m}^{-2}$ for a beam of 100 keV electrons and $10 \text{ } \mu\text{A m}^{-2}$ for a beam of 1 keV electrons.

4.3.5. Small scale structure and Ganymede footprint

As for the discontinuity and IFP, a distribution of primary electrons based on the data (section 4.3.3.) has not been calculated for the small scale structure emission seen in the Jan. 16 STIS observation and the Ganymede footprint seen in the Jan. 24 observation, because of the low number of pixels involved and thus low statistics. Interestingly, the extracted spectra show very important hydrocarbon attenuation for both emissions, with a CR of 11 and 6.7 for the SSS and the GFP, respectively. By using the CR method, these values correspond to $\langle E \rangle$ of 232 and 157 keV, respectively. Since the primary electron distribution and the shape of the emission profile are unknown for these emissions, the spectral analysis is identical in both cases. We present in Figure 8 the analysis of the GFP emission, first tested with a ‘thin’ emission profile to fit the observed spectrum. A best fit of the GFP spectrum associated with a thin Maxwellian-type VER profile is presented in Fig. 8 a & b. The emission peaks at 200 km, which corresponds to energies of 140 keV, relatively close to the 157

keV value obtained from the CR method. The difference is explained by the fact that our regression method tries to minimize differences between the modeled and observed spectrum over the full 1250-1700 Å spectral region and not only the bandwidths associated with the CR. Several spectral features of the GFP spectrum are not well defined because of the low S/N, due to the low number of pixels used to define this region. Still, comparison with an unabsorbed H₂ spectrum (dashed blue line in Fig.8b) clearly demonstrates that the GFP emission is strongly affected by hydrocarbon absorption.

We also tested ‘thick’ synthetic emission profiles to fit the GFP spectrum, with a method similar to the one used for the IFP in the previous paragraph. It is found that ‘thick’ emission profiles could not be sufficiently absorbed to provide the observed high CR. If the VER is peaking at altitudes higher than 400 km, the attenuation is too weak to provide high CR, and if the peak is situated at lower altitudes, the shape of the original thick VER changes (decrease of the FWHM) as the emission below 150 km is strongly dampened in our procedure, in order to account for the altitude limit reached by electrons whatever their energy. Since the vertical thickness of the emission decreases as the emission moves to lower altitude, the shape of the VER becomes similar to the ‘thin-Maxwellian type VER’, as used in the CR method.

4.4. Alternative interpretation of UV emission: ion precipitation

Strong X-ray emissions have been observed over the past 20 years from the polar caps of Jupiter (e.g. Branduardi-Raymont et al., (2008), Ozak et al., (2013) and references therein), with a power on the order of 1 GW for each hemisphere (Elsner et al., 2005). This emission is thought to result from precipitation of oxygen and sulfur ions of several MeV. A model describing the ion precipitation process has been developed by Ozak et al. (2010, 2013). Outputs from this model include secondary electron production rates, field-aligned currents, and auroral emission rates in the UV bandwidth resulting from secondary electrons produced by ion precipitation. Interestingly, this model

predicts an unabsorbed UV auroral emission of 60-80 kR in the north and 10-20 kR in the south in the Lyman and Werner bands of H₂, assuming a total X-ray emitted power of 1 GW. This corresponds to a 'UV efficiency' of ~3 kR per incident mW m⁻² from ion precipitation, i.e. on average a third of the electron precipitation efficiency. These authors tested precipitation of ions with energies 1, 1.5 and 2 MeV/u, the latter considered as an upper limit of plausible values. The vertical emergent UV brightness of 60-80 kR produced by ion precipitation in the northern polar cap corresponds well to the vertical brightnesses obtained from the FL1 and FL2 STIS north observations studied here (mean vertical value from 47 to 85 kR, Fig. 3 g-i). In addition to the emergent brightness, the UV profiles resulting from ion precipitation can be used in our model to derive the CR associated with ion precipitation which can be compared with the STIS observations.

The Jovian atmosphere adopted by Ozak et al. (2013), first introduced by Maurellis and Cravens (2001), is based on Galileo probe data (Seiff et al. 1996) and remote observations (Sada et al. 1998). In particular, the temperature profile (hence the H₂ density profile) used by Ozak et al. (2013) differs from the Grodent et al. (2001) profiles used in the present study. The ion precipitation model by Ozak et al. (2013) has thus been performed with the Grodent et al. (2001) atmosphere in order to derive H₂ emission profiles directly compatible with the other outputs from the present work.

The VER profiles of H₂ emission so obtained are found to peak between 325 and 290 km for 1 and 2 MeV/u ions, respectively. They are used in our auroral model as described in appendix A1 and the related emergent spectra are determined for $\chi=70^\circ$, i.e. the typical viewing angle corresponding to the 'flare' emissions seen in our observations. It is seen that the CR due to the UV emission resulting from ion precipitation varies from 1.10 to 1.34 for 1 MeV/u to 2 MeV/u ions, respectively (black curves in Fig. 11b). These values are significantly lower than the CR from 2.0 to 3.2 obtained from the 'flare' emissions observed in the present study. The latter values correspond well to previous determinations of the CR obtained so far. Fifteen polar cap spectra observed by STIS have been examined by Gustin et

845 al. (2004a). They were characterized by CR values from 1.7 to 7.5: one CR at 1.7, five CRs between
2.1 and 3, and nine CRs above 3. Several observations from this data set were obtained in the time tag
mode, allowing inspection of transient events (Gérard et al., 2003). These data reveal that only two
flaring events were associated with a CR of 1.7, nine with a CR between 2 and 3 and two with a CR
higher than 3. These two studies thus reveal that 89% of the observed flare emissions (93% for Gustin
850 et al. (2004) and 85 % for Gérard et al. (2003)) are associated with CRs higher than 2, leaving ~10 % of
the cases with a CR of 1.7. Referring to these observed CRs compared to the CR simulated from the
Ozak et al. (2013) outputs, polar emission exclusively due to ion precipitation is thus very unlikely.
Still, a partial contribution of ions to the total UV emission is possible. The effect of the ion
precipitation, combined with electron precipitation, would be to lower the observed CR and thus
855 underestimate the mean energy of the precipitating electrons. The 60-80 kR UV brightnesses resulting
from this modeled ion precipitation are relatively low and correspond to 'typical polar cap emissions.
Much brighter emissions have been observed at high latitudes. For example, a spectacular event has
been observed simultaneously by Chandra in X-ray and HST/STIS in FUV, on 26 February 2003
(Elsner et al. 2005). During this event, the FUV brightness reached ~5 MR and the emitted X-ray
860 power attained 8 GW and both emissions were somewhat correlated. In this exceptional event, a
significant contribution from ion precipitation to the UV emission is possible, as these observed
characteristics were not considered by Ozak et al (2013) (X-ray power peak at ~8 GW compared to the
'standard' 1 GW used here).

865 **5. Summary – Conclusions.**

Several important results have been obtained from this study, both from a theoretical and observational
point of view.

1. On the theoretical side, we set up a model of auroral H₂ emission sensitive to the 700-1700 Å
870 bandwidth. This model adopts the auroral atmosphere described in by Grodent et al. (2001). Several
Maxwellian distributions of primary electrons at mean energies from 2 to 1000 keV have been used to
determine the influence of the energy input on the characteristics of the auroral emissions. Results
show that:

- the altitude of the emission peak monotonically decreases as $\langle E \rangle$ increases. Assuming that 1
875 MeV corresponds to the upper limit possibly reached by primary electrons, virtually no auroral photons
can be emitted at altitudes below ~130 km, which corresponds to an H₂ column of $\sim 5.5 \times 10^{22} \text{ cm}^{-2}$.

- the efficiency of the atmosphere to produce UV aurora strongly depends on the energy of the
precipitating electrons. For a constant energy flux input of 1 mW m^{-2} , the brightness in the Lyman and
Werner bands is maximum at $\langle E \rangle = 20 \text{ keV}$ (14.6 kR) and then continuously decreases for higher
880 energies to reach 3.2 kR at 500 keV.

- in an altitude scale, the FWHM of the emission profile due to Maxwellian distributions
decreases with increasing mean energy with values from 440 km (2 keV) to 53 km (500 keV).

The observed color ratio (CR) has been related to the mean energy of the precipitating electrons
885 $\langle E \rangle$ with a model that includes effects of the emission angle χ and the width D of the auroral curtain. It
is found that

- a) for a given $\langle E \rangle$, the perceived CR increases with χ ,
- b) for a given χ , the auroral width D has a minor effect on the CR
- c) the unabsorbed (intrinsic) brightness strongly depends on χ and the auroral width considered (Fig.
890 11).

It should be kept in mind that the CR- $\langle E \rangle$ relationship is strongly model-dependent. It is influenced by

- 1) the temperature profile, 2) the altitude of the hydrocarbon homopause level and 3) the energy

distribution of the primary electrons. This modelling has been applied to two sets of STIS FUV spectral images of the north Jovian aurora. Several auroral regions have been selected in both images: morning
895 main emission 1 (ME1), main emission 2 (kink region, ME2), discontinuity (DC) flare emissions (FL1 and FL2), Io footprint (IFP), Ganymede footprint (GFP) and small scale structure emission (SSS). Interestingly each region exhibit very specific characteristics, as described hereafter.

2. Mean Energy

900 1. The maps of the electron mean energy show that the spatial distribution of the precipitating electron energy is clearly not uniform (Fig. 2 & Fig. 3). Because of low S/N of several individual pixels, such maps only provide a first order estimate of the mean energy of the precipitating electrons.

2. The spectra obtained from the sum of the pixels defining each auroral region do not bring direct information on the electron energy distribution, but provide the most accurate average mean energy for
905 each region. The morning ME1 region is the most energetic (~260 keV), followed by the ‘kink’ ME2 region (~110 keV) and the polar cap flare emissions (~75 keV). The higher S/N of the spectra allows determining mean energies for the DC, IFP, GFP and SSS regions, defined from low number of pixels and/or weak signal. Mean electron energies are ~60 keV for the DC, ~155 keV for the GFP, 230 keV for the SSS emissions. The CR method, providing energies around 20 keV for the IFP, was found
910 inappropriate for this emission (see point 4.3.4.). The spatial distribution of mean energies corresponds well with the energy map from the 8 Jan 2014 STIS observation presented in Gérard et al. (2014), which was obtained with comparable CML, although our $\langle E \rangle$ values are lower.

3 Mean energy – energy flux relationship

915 The pixels in each auroral region have been sorted by ascending brightness and binned by groups of height data points to increase the S/N ratio and determine reliable mean energy and energy

flux values. The electron energies of the main emission and flare regions are positively correlated with the energy fluxes and follow well Knight's (1973) theory of field-aligned electric field acceleration associated with field-aligned currents. Magnetospheric electron sources have densities around 2400 m^{-3} for ME2 with a thermal energy of 2.5 keV, which is consistent with a density between 1000 m^{-3} and 10000 m^{-3} measured by Voyager in the Jovian magnetospheric equatorial plane (Scudder et al. 1981). The relativistic generalization of Knight's theory developed by Cowley (2006) has been used for ME1, as the electron's acceleration is a significant fraction of the electron rest energy for this region. Lower electron source densities are obtained ($520\text{-}620 \text{ m}^{-3}$) and correspond well with values obtained from the study of a remarkably bright auroral morning arc observed by STIS on 21 September 1999 (Gustin et al. 2006).

4 Spectral analysis

Strong hypotheses on the atmospheric structure (hydrocarbon density profiles, temperature profile), and on the energy distribution of the primary electrons must be postulated to infer the CR- $\langle E \rangle$ relationship. Since these characteristics are not strongly constrained by observations, results obtained from the CR method are highly model-dependent. The spectral resolution of the STIS G140L observations examined here is too low ($> 1 \text{ \AA}$) to resolve the H_2 rotational lines, which hence does not allow us to derive information on the temperature profile of the auroral atmosphere. Still, the 500 \AA bandwidth STIS FUV spectra at 12 \AA resolution provide more information than the CR, which is limited to the ratio of two integrated 70 \AA bands. We tested the hypothesis that comparison between synthetic and observed spectra provide information on the energy distribution of the precipitating electrons and on the appropriateness of the atmospheric model used. We use Chapman functions defined with 4 free parameters to simulate the VER profile and tested the hypothesis that a specific VER profile (thus a unique electron energy distribution) provides spectral fits significantly better than

any other VER, which would bring additional constraints on primary electron characteristics. Chapman functions were found to be an excellent tool to fit the VER resulting from the two-stream energy degradation model of Grodent et al. (2001) and are thus appropriate for simulations. It is found that without constraints, several significantly different VER provide equivalent fits to the observed spectra, which does not allow to provide additional knowledge on the primary electron distributions. Still, simulations show that C_2H_6 is required to fit strongly absorbed spectra and that CH_4 , C_2H_2 and C_2H_6 mixing ratio profiles of the model are appropriate to fit the different bandpasses influenced by these hydrocarbons.

In general, the peak of the intrinsic VER emission is situated at lower altitude than the observed peak emission because of the attenuation at and below the hydrocarbon layer. For example, for a CR of 3.2, the difference of perceived altitude is ~45 km.

Since the energy-flux relationship is very specific to each auroral regions, we used this property to determine energy distributions of the primary electrons, constrained by the $\langle E \rangle$ and E_f values associated to each pixel of a given region (section 4.3.3. & Fig. 6). It is seen that distributions for ME1 are Maxwellian, distributions for the ME2 and flares region exhibit more electrons at high energy (compared to a Maxwellian distribution). These electron distributions provide ad hoc solutions able to represent the average emission in each region.

Unlike the other regions, ACS images provide observed constraints on altitude and shape of the VER profile of the IFP emission (Bonfond et al. (2009)). The spectral analysis shows that a synthetic VER profile consistent with a Maxwellian electron energy distribution and the CR method provides a good fit to the IFP spectra, but is incompatible with the observed constraints. Indeed, the Maxwellian-type VER profile is much thinner (~150 km) and peaks at much lower altitude (~330 km) than the FWHM of ~1200 km and peak altitude between 550 and 1300 km provided by the limb analysis of the ACS images. The CR method is thus not applicable to the IFP emission. Several tests reveal that the

965 only way to reconcile constraints on the VER profile and hydrocarbon absorption altogether is to allow
an upward shift the hydrocarbon layer with respect to the fixed H₂ density profile.. Several fits with
varying levels of upwelling show that the homopause altitude shift and the peak altitude of the best
VER profile are linearly correlated. An minimum upwelling of 100 km is needed to satisfy all the
constraints, but an homopause upward shift of 280 km is needed to be consistent with an IFP emission
970 peaking at a typical altitude of ~ 900 km. This corresponds to an increase of K_h to $2 \times 10^{11} \text{ cm}^2 \text{ s}^{-1}$, i.e. 6
orders of magnitude above the nominal value used in the Grodent et al. (2001) model, which seems
very unlikely. Other phenomena like vertical winds could account for such hydrocarbon upward shift,
but our present data cannot strongly constrain the cause of such a shift.

975 5 Ion precipitation

The auroral model described in this paper has been applied to FUV emission profiles resulting
from the precipitation of heavy ions in the 1-2 MeV/u range (Ozak et al. 2013), in order to determine if
such precipitation could explain, at least partly, polar cap auroral emissions. Results show that the CR
of the UV emission due to ions (<1.4) is lower than the CR observed poleward of the main oval (≥ 2). A
980 significant contribution of ion precipitation to flare emissions typically observed (brightness <100 kR)
is thus very unlikely. In case of brighter flare emission, such as the 5 MR of FUV emission recorded by
STIS on 26 February 2003 concurrently with Chandra X-ray observation (Elsner et al. 2005), an
important contribution of ion precipitation cannot be ruled out.

985

Appendix A: Determination of the mean energy of the primary electrons

A1. Auroral model

Our auroral atmospheric model uses the auroral thermal profile and H_2 , CH_4 and C_2H_2 profiles described in Grodent et al (2001). It couples a two-stream electron transport model of energy deposition with a 1-D thermal conduction model. From the initial North Equatorial Belt (NEB) atmospheric structure deduced from Galileo observations (Gladstone et al., 1996) adapted to the larger gravitational acceleration near the poles, the Grodent et al. (2001) model self-consistently adapts to the effects of the incident primary electron distribution. At each layer of the atmosphere, the energy distribution of the primary electrons changes as they interact with the atmosphere through ionizations, collisions and creation of secondary electrons. The thermal profile and the CH_4 and C_2H_2 profiles automatically adapt to the auroral conditions. This model is constrained by observations such as the altitude of the H_2 emission peak, the temperature profile deduced from UV and infrared emissions, and UV emission rates deduced from observed images and spectra. All density profiles from this atmosphere, defined from 117 to 3900 km above the 1 bar level, are used in our model. To extend these profiles to lower altitudes and cover the widest possible altitude range, we merged the density profiles from -65 to 117 km defined in the low-latitude atmospheric model of Moses et al. (2005), and scaled them to obtain continuous densities in our final auroral atmosphere. Ethane is not included in the Grodent et al. (2001) atmosphere. Since the signature of ethane is clearly observed in the case of strong absorption ($\text{CR} > 2$), it is included here by using the C_2H_6 density profile described by Moses et al. (2005). The vertical density profile in our model is obtained by multiplying the C_2H_6 mixing ratio relative to H_2 found in Moses et al. (2005) by the H_2 density profiles outlined in Grodent et al. (2001). This auroral atmospheric model, along with the Moses model are shown in Fig. 9. The 'hotter' temperature profile of the Grodent et al. (2001) model in Fig. 9a demonstrates the consequence of the electron precipitation on the atmosphere, which also has an effect on the H_2 scale height shown in Fig.

9b. The most abundant absorber is methane, followed by ethane and acetylene (Fig. 9c).

To relate $\langle E \rangle$ to CR, we use single Maxwellian distributions with mean energies from 2 to 1000 keV, representing the initial energy distribution of the precipitating electrons and covering the widest possible range of mean energies. An auroral volume emission rate vertical profile (VER: number of
1015 auroral photons emitted per second per cm^3) is then produced from each Maxwellian distribution from the two-stream Grodent et al. (2001) model. The atmosphere is divided into altitude bins of 10 km, from -65 to 3900 km and a normalized unabsorbed synthetic H_2 spectrum, weighted by the local VER, is produced at each layer. At altitude bin z , the local unabsorbed H_2 emission $H_2^{unabs}(z)$ is

$$H_2^{unabs}(z) = H_2^{synth} * VER(z) * dz * Chap(z, \chi) \quad (6)$$

1020 where H_2^{synth} is the unabsorbed H_2 spectrum in 700-1700 Å, $VER(z)$ is the auroral emission rate at altitude z , dz is the thickness of a layer and $Chap(z, \chi)$ is the Chapman function that converts vertical to slant thickness in curved atmosphere (Smith and Smith, 1972). This function substantially differs from the simple $\sec(\chi)$ factor for high χ . We use the gravity and radius at 75° latitude to compute $Chap(z, \chi)$, which brings values applicable to the whole polar region.

1025 Accounting for hydrocarbon absorption, the attenuated H_2 emission at altitude z is

$$H_2^{abs}(z, \lambda) = H_2^{unabs}(z, \lambda) * e^{(-\tau(z, \lambda))} \quad , \quad (7)$$

where the optical depth τ at wavelength λ at altitude z is

$$\tau_{z, \lambda} = \sigma_{\lambda}(\text{CH}_4) \int_z^{\infty} n_{\text{CH}_4} ds + \sigma_{\lambda}(\text{C}_2\text{H}_2) \int_z^{\infty} n_{\text{C}_2\text{H}_2} ds + \sigma_{\lambda}(\text{C}_2\text{H}_6) \int_z^{\infty} n_{\text{C}_2\text{H}_6} ds \quad , (8)$$

1030 where $\sigma_{\lambda}(\text{CH}_4)$, $\sigma_{\lambda}(\text{C}_2\text{H}_2)$, $\sigma_{\lambda}(\text{C}_2\text{H}_6)$ are the methane, acetylene and ethane absorption cross sections at wavelength λ , and n_{CH_4} , $n_{\text{C}_2\text{H}_2}$ and $n_{\text{C}_2\text{H}_6}$ are the local number densities. The slant path length ds corresponds to the thickness of a layer dz multiplied by the Chapman function $Ch(z, \chi)$, at emission angle χ .

A point that is often overlooked in auroral study is the finite size of the auroral curtain. To

1035 account for this aspect, we assume that the auroral emission has a rectangular section of width D that extends from the lowest to the highest limits of the atmospheric model (Figure 10). Although the whole altitude range is taken into account for ease of computing, only a limited vertical extension contributes to the emergent emission (represented by the shaded region in Fig.10). The emergent unabsorbed spectrum $H_2^{unabs_tot}$ (i.e. the intrinsic auroral emission), in kR, at angle χ is

1040
$$H_2^{unabs_tot} = \sum_{i=0}^{n-1} R_i / ne \quad (9)$$

where each emergent beam ray R_i is the sum of the local emissions H_2^{unabs} defined by (6) intercepted by R_i inside the rectangle of width D (dots in Figure 10), n is the total number of rays considered, and e is the minimum distance between two consecutive rays. Similarly, the emergent H_2 spectrum attenuated by the hydrocarbon layer at and above the emission is given by equation (9) where H_2^{unabs} is replaced
 1045 by H_2^{abs} defined by equation (7). The resulting synthetic emergent spectrum represents the observed, potentially absorbed, emission as seen by an observer, from which a CR can be determined.

It should be emphasized here that the kilo-Rayleigh is a 'brightness density', as it is defined 'per unit of emitted surface'. On the one hand, if one takes into account rays that do not cross substantial emission (i.e. far from the emission peak, e.g. R_j in Fig.10), this would significantly lower the total
 1050 brightness of the emergent emission. On the other hand, selection of the brightest rays close to the emission peak only artificially enhances the perceived emergent brightness of the whole emission. In a nadir view ($\chi=0^\circ$), the procedure illustrated in Fig.10 is not necessary, as the emergent brightness is simply the sum of the vertical VER at each layer multiplied by the vertical thickness dz of a layer. It is thus independent of the number of rays considered for an homogeneous emission. We use this vertical
 1055 emergent brightness to determine the number of R_i that provides a smooth transition between vertical and slant emission. We thus determined the emergent unabsorbed H_2 intensity and CR at $\chi=5^\circ$ for various number of rays at fixed distance e by varying n in equ. (9). We found that the number of R_i considered in the sum leading to 99% of the total emission provides smoothest transition of CR and

brightness between $\chi = 0^\circ$ and 5° . This number varies with the width of the VER profile and the
1060 viewing angle considered.

To simulate the effect of the auroral width D , the mean energy $\langle E \rangle$ and the viewing angle χ on
the emergent unabsorbed and absorbed H_2 emission, we used several single Maxwellian electron
distributions from $\langle E \rangle = 2$ to 1000 keV and determined the corresponding auroral VER by using the
two-stream model from Grodent et al. (2001). We applied these VER in our model for various auroral
1065 widths and viewing angles from 0 to 85° . Several outputs from this model are summarized in Figure 11.
Figure 11a and 11b show the CR as function of the incident electron energy $\langle E \rangle$, for an auroral curtain
of arbitrary width $D=1500$ km. It is seen that the effect of the viewing angle is far from negligible:
compared to vertical values, the CR increase at $\chi=70^\circ$ is less than 5% for $\langle E \rangle < 20\text{keV}$, 30% for $\langle E \rangle$
 $= 100$ keV and reaches 41% at $\langle E \rangle = 400$ keV.

1070 The effect of the auroral width on the emergent unabsorbed brightness is shown in Fig 11 c for
primary electrons of 80 keV (which can be seen as an 'average' energy in the case of the Jovian aurora).
Four cases are examined: $D=150$ km, which is used for footprint emissions, $D=600$ km, which
represents a narrow auroral emission comparable to the main oval, $D=1500$ km which can represent the
thicker, more diffuse emissions, as seen poleward of the main oval, and a case with $D=10000$ km,
1075 which depicts an homogeneous “infinite” auroral emission. The latter case simulates the effect of the
viewing angle on the H_2 brightness and CR when the auroral width is not taken into account. For a
viewing angle of 70° , the brightness increase is very important for the infinite case (increase by a factor
of 2.8), and then decreases as the aurora is getting thinner (factor of 1.66, 1.04 and 0.39 for $D=1500$,
600 and 150 km, respectively). Interestingly, for $D=150$ km, the emergent brightness decreases with the
1080 viewing angle. This counter intuitive geometric effect is due to the limited vertical extent of the auroral
emission. For example, simulations show that the VER profile obtained from a 80 keV Maxwellian
distribution peaks at 226 km and has a FWHM of 106 km. The portion of the emission centered on 226

km and of width 150 km corresponds to 85 % of the total emission. To first order, the transverse cut of an 80 keV auroral curtain of 150 km width can thus be seen as a square of 150 x 150 km width. The observed emission at $\chi=0^\circ$ and 90° should then be very similar, which is what we observe in the blue curve of Fig. 11c. For intermediatr viewing angles, although the total emitted brightness is unchanged, the total surface perceived by the observer increases with $\secant(\chi)$. Since the kilo-Rayleigh represents an observed brightness per emitting surface unit, the observed brightness decreases for increasing χ . This effect diminishes as the auroral width increases, as it is compensated by the increase of emission width. It should be noted that since the FWHM of the VER decreases with increasing $\langle E \rangle$ (see appendix A2), the effect of D on the emergent brightness is energy dependent. This characteristic is an important matter and is taken into account in the data analysis. As seen in Fig. 11d, the effect of the auroral width on the CR is quite minor, as the maximum variation due to the auroral width is 3% at 85° for $\langle E \rangle = 80$ keV, below the accuracy achieved by the observations. This means that the width affects the absorbed and unabsorbed emissions in a similar way. The CR - $\langle E \rangle$ relationship is presented in Fig. 11e for a standard viewing angle of 70° . Considering that 70° is a typical viewing angle of Jupiter's poles as seen from HST, Fig. 11e can be used to estimate the incident precipitating energy from an observed absorbed emission. The effect of the auroral width on the emergent brightness is presented in Fig. 11f, for an incident electron energy flux of 20 mWm^{-2} . As expected, the observed brightness increases with the width of the aurora. The point at 10000 km on the x-axis of Fig. 11f simulates the observed brightness that would be obtained without taking into account D . The red curve shows the emergent total H_2 brightness after attenuation by hydrocarbons. On average, the factor between the unabsorbed and absorbed emission in Fig. 11f is 2.12. Another way to interpret this plot is to say that in order to obtain a given emergent brightness, the input energy flux must increase as the auroral width decreases.

A.2. Characteristics of auroral emission produced from Maxwellian electron distributions

Before applying our model to observations, it is interesting to explore the characteristics of the auroral emission resulting from 2 to 1000 keV single Maxwellian distributions computed in the Grodent et al. (2001) model. All cases used a total energy flux input of 20 mW m^{-2} . We first tested the hypothesis that the VER profiles could be approximated by a Chapman profile

$$\nu = \nu_m e^{[f(1-\alpha-e^{-\alpha})]} \quad (10)$$

where $\alpha = (z-z_m)/H$ is the reduced height, ν_m is the maximum of ν at the altitude z_m and f is a form factor. Since f and the scale height H influence the shape and FWHM of the VER, H should be considered here as a simple free parameter, not necessary equal to the actual scale height of the emission. We found that the VER originating from these incident Maxwellians could be easily fitted with Chapman profiles. The influence of $\langle E \rangle$ on the emission profiles is summarized in Figure 12. As seen in Fig. 12a, the altitude of the maximum emission naturally decreases as the mean electron energy increases. The peak altitude slowly decreases and reaches $\sim 160 \text{ km}$ for electrons in a Maxwellian distribution of $\langle E \rangle = 500 \text{ keV}$. There is very little emission at lower altitudes, which is intuitively explained by the exponential increase of the number of H_2 molecules encountered by the electrons with decreasing altitude, which makes it difficult to reach deeper layers, whatever the primary electron energy. The emission peaks at 165 km for $\langle E \rangle = 400 \text{ keV}$ and at 163 km for $\langle E \rangle = 500 \text{ keV}$. This 2 km difference leads to a variation of 100 keV in the mean energy determination. The corresponding CR at $\chi = 70^\circ$ is 20 and 23 for 400 and 500 keV, respectively, i.e. a 15% difference. Since high CR values correspond to strongly attenuated emission in the $1230\text{-}1300 \text{ \AA}$ range (i.e. emission with low S/N ratio), one must be careful in the interpretation of data when high electron energies are involved. Figure 12a also shows that the altitude of the VER peak resulting from a Maxwellian distribution of electrons at a given $\langle E \rangle$ corresponds very well with the altitudes attained by monokinetic electrons of same energy. The altitude reached by monokinetic electron has been calculated following a continuous slowing down

stopping power approximation. The plot in Fig. 12b shows that the maximum value of the VER increases with $\langle E \rangle$, to reach a maximum at 200 keV, and then decreases for higher energies. The FWHM of the VER profiles also significantly changes with $\langle E \rangle$ and decreases with increasing energies (Fig. 12c). Although these FWHM remain fixed in pressure scale for different atmospheres, they
1135 depend on the pressure-altitude relationship in altitude scale. Values in Fig. 12c are thus only valid in the Grodent et al. (2001) atmosphere, as the temperature profile affects the scale height of the emission. As seen in Fig. 12d for a constant incident energy flux of 1 mW m^{-2} , the emergent vertical intensity strongly depends on the mean energy. The efficiency of the atmosphere to create UV auroral emission is maximum for $\langle E \rangle = 20 \text{ keV}$ ($14.6 \text{ kR per mW m}^{-2}$) and then sharply drops for higher energies (7.8 kR
1140 per mW m^{-2} at 200 keV to only $3.2 \text{ kR per mW m}^{-2}$ at 500 keV). In other words, the energy flux input necessary to create an aurora of given emergent brightness strongly depends on the energy of the precipitating electrons. The value of $10 \text{ kR per mW m}^{-2}$ based on several aeronomic models and proposed as a rule of thumb by Gustin et al. (2012) should then be reconsidered when the mean electron energy is not in the 70-140 keV range. Since the energies determined in our study exceed this
1145 range, values shown in Fig. 12d are used throughout this work. They are particularly important when setting up the energy – energy flux diagrams, as presented in section 4.2.

1150

1155 Highlights:

1. maps of the mean energy of the electrons creating the aurora
2. each auroral region presents very specific characteristics
3. mean energy and energy flux are correlated
4. an hydrocarbon upwelling of min. 100 km is necessary to reconcile STIS and ACS constraints
1160 on IFP emission
5. ion precipitation cannot explain ‘typical’ polar cap flare ultraviolet emissions

Acknowledgements.

1165 We thank two anonymous reviewers for their constructive comments and suggestions that allowed to
improve the manuscript. J.G. was supported by the PRODEX program managed by ESA in
collaboration with the Belgian Federal Science Policy Office. B.B. acknowledges support from the
Belgian Fund for Scientific Research (FNRS). This research is based on observations with the Hubble
Space Telescope, obtained at the Space Telescope Science Institute, which is operated by AURA for
1170 NASA.

References.

- 1175 Bonfond, B., D. Grodent, J.-C. Gérard, A. Radioti, J. Saur and S. Jacobsen, UV Io footprint leading spot: a key feature for understanding the UV Io footprint multiplicity? *Geoph. Res. Let.*, **35**, L05107, doi:10.1029/2007GL032418.
- Bonfond, B., Grodent, D., Gérard, J.C., Radioti, A. , Dols, V., Delamere, P.A., Clarke, J.T. The Io UV
1180 footprint: location, inter-spot distances and tail vertical extent. *J. Geophys. Res.* **114**, A07224, doi:10.1029/2009JA014312, 2009.
- Bonfond, B., M.F. Vogt, J-C. Gérard, D. Grodent, A. Radioti and V. Coumans, Quasi-periodic polar flares at Jupiter: a signature of pulsed dayside reconnection? *Geoph. Res. Let.*, **38**, L022104,
1185 doi:10.1029/2010GL045981, 2011.
- Bonfond, B., S. Hess, J-C. Gérard, D. Grodent A . Radioti, V. Chantry, J. Saur, S. Jacobsen and J. T. Clarke, Evolution of the Io footprint brightness I: far-UV observations, *Plan. And Space Science*, **88**, 64-75, doi:10.1016/j.pps.2013.05.023, 2013.
- 1190 Branduardi-Raymont, G., Elsner, R.F., Galand M., Grodent, D., Cravens, T.E., Ford, P., Gladstone, G.R., Waite, J.H., Spectral morphology of the X-ray emission from Jupiter's aurorae, *J. Geophys. Res.*, 113 (A2), doi: 10.1029/2007JA012600, 2008.
- 1195 Chané, E., Saur, J., Poedts, S., Modeling Jupiter's magnetosphere: Influence of the internal sources, *J. Geophys. Res.* **118**, 2157-2172, doi: 10.1002/jgra.50258

Clarke, J.T., Ajello, J., Luhmann, J., Schneider, N., Kanik, I., Hubble Space Telescope UV spectral observations of Io passing into eclipse, *J. Geophys. Res.*, 99(E4), 8387, doi:10.1029/93JE02547, 1994.

1200

Clarke, J. T., Ballester, G. E., Trauger, J., Evans, R., Connerney, J. E. P., Stapelfeldt, K., Crisp, D., Feldman, P. D., Burrows, C. J., Casertano, S., and 11 coauthors, Far-Ultraviolet Imaging of Jupiter's Aurora and the Io "Footprint", *Science*, 274, 404, doi :10.1126/science.274.5286.404, 1996

1205 Clarke, J.T., Ballester, G. Trauger, J. Ajello, J. Pryor, W., Tobiska, K., Connerney, J.E.P., Gladstone, G.R., Waite, J.H., Ben Jaffel, L. Gérard, J.C. Hubble Space Telescope imaging of Jupiter's UV aurora during the Galileo orbiter mission, *J. Geophys. Res.*, 103 (E9), 20217-20236, doi : 10.1029/98JE01130, 1998.

1210 Clarke, J.T., Ajello, J., Ballester, G., Ben Jaffel, L., Connerney, J., Gérard, J.-C., Gladstone, G.R., Grodent, D., Pryor, W., Trauger, J., Waite, J.H., Ultraviolet emissions from the magnetic footprints of Io, Ganymede and Europa on Jupiter, *Nature*, **415** (6875), p. 997-1000, 2002.

Cowley, S. W. H., and E. J. Bunce, Origin of the main auroral oval in Jupiter's coupled magnetosphere-ionosphere system, *Planet. Space Sci.*, **49**, 1067, doi: 10.1016/S0032-0633(00)00167-7, 2001,

1215 Cowley, S. W. H., and E. J. Bunce, Modulation of Jupiter's main auroral oval emissions by solar wind induced expansions and compressions of the magnetosphere, *Planet. Space Sci.*, **51**, 57-59, doi: 10.1016/S0032-0633(02)00118-6, 2003.

1220

- Cowley, S. W. H., Current-voltage and kinetic energy flux relations for relativistic field-aligned acceleration of auroral electrons, *Ann. Geophys.*, 24, 325 – 338, doi: 10.5194/angeo-24-325-2006, 2006.
- 1225 Dols, V., Gérard, J.-C., Clarke, J.T., Gustin, J., Grodent, D. Diagnostics of the jovian aurora deduced from ultraviolet spectroscopy: Model and HST/GHRS observation. *Icarus* 147, 251–266., doi :10.1006/icar.2000.6415, 2000.
- Dols, V., P. A. Delamere, and F. Bagenal. A multispecies chemistry model of Io's local interaction with the Plasma Torus, *J. Geophys. Res.*, 113, A09208, doi:10.1029/2007JA012805, 2008.
- 1230 Dumont, M., D. Grodent, A. Radioti, B. Bonfond, J.C. Gérard, Jupiter's equatorial features: possible signatures of magnetospheric injections, *J. Geophys. Res.* **119**, doi:10.1002/2014JA020527, 2014.
- 1235 Elsner, R. F., Lugaz, N. and 9 coauthors, Simultaneous Chandra X-ray, Hubble Space Telescope ultraviolet, and Ulysses radio observations of Jupiter's aurora, *J. Geophys. Res.*, 110, A01207, doi:10.1029/2004JA010717, 2005.
- Gérard, J.-C., J. Gustin, D. Grodent, P. Delamere, and J. T. Clarke. The excitation of the FUV Io tail on Jupiter: Characterization of the electron precipitation, *J. Geophys. Res.*, 107(A11), 1394, doi:10.1029/2002JA009410, 2002.
- 1240 Gérard, J.-C., J. Gustin, D. Grodent, and J. T. Clarke. Spectral observations of transient features in the FUV Jovian polar aurora, *J. Geophys. Res.*, 108(A8), 1319, doi:10.1029/2003JA009901, 2003.
- 1245 Gérard, J.-C., Gustin, J., Pryor, W.R., Grodent, D., Bonfond, B., Radioti, A., Gladstone, G. R., Clarke,

J.T., Nichols, J.D., Remote sensing of the energy of auroral electrons in Saturn's atmosphere: Hubble and Cassini spectral observations, *Icarus*, 223 (1), doi: 10.1016/j.icarus.2012.11.033.

- 1250 Gérard, J.C., Bonfond, B., Radioti, A., Clarke, J.T., Gladstone, G.R., Waite, J.H., Bisikalo, D., Shematovich, V.I. Mapping the electron energy in Jupiter's aurora: Hubble spectral observations, *J. Geophys. Res.*, 119 (11), doi : 10.1002/2014JA020514, 2014.

- Gladstone, G. R., and T. F. Skinner 1989. Spectral analysis of jovian auroral emissions. In Proceedings of the Workshop on Time-Variable Phenomena in the Jovian System (M. J. S. Belton, R. A. West, and J. Rahe, Eds), pp. 221–228. NASA SP-494.
- 1255

Gladstone, G.R., Allen, M., Yung, Y.L. Hydrocarbon photochemistry in the upper atmosphere of Jupiter. *Icarus* 119, 1–52, 1996.

1260

Grodent, D., Waite Jr., J.H., Gérard, J.C. A self-consistent model of the jovian auroral thermal structure. *J. Geophys. Res.* 106, 12933–12952, doi:10.1029/2000JA900129, 2001.

- Grodent, D., Gérard, J.C., Gustin, J., Mauk, B.H., Connerney, J.E.P, Clarke, J.T. Europa's FUV auroral tail on Jupiter. *Geophys. Res. Lett.*, 33, L06201, doi : 10.1029/2005GL025487, 2006.
- 1265

Grodent, D., J.-C. Gérard, J. Gustin, B.H. Mauk, J.E.P. Connerney, J.T. Clarke, Europa's FUV auroral tail on Jupiter. *Geophys. Res. Lett.* 33, L06201, doi:10.1029/2005GL025487, 2006.

- 1270 Grodent, D., B. Bonfond, J.-C. Gérard, A. Radioti, J. Gustin, J.T. Clarke, J. Nichols and J.E.P. Connerney, Auroral evidence of a localized magnetic anomaly in Jupiter's northern hemisphere, *J.*
- 54

Geophys. Res., 113, A09201, doi:10.1029/2008JA013185, 2008.

1275 Gustin J., D. Grodent, J. -C. Gérard, and J. T. Clarke, Spatially resolved far ultraviolet spectroscopy of
the Jovian aurora, *Icarus*, **157**, 91, 2002.

Gustin, J., J.-C. Gérard, D. Grodent, S. W. H. Cowley, J. T. Clarke, and A. Grard. Energy-flux
relationship in the FUV Jovian aurora deduced from HST-STIS spectral observations, *J. Geophys. Res.*,
109, A10205, doi:10.1029/2003JA010365, 2004a.

1280

Gustin, J., Feldman, P.D., Gérard, J.-C., Grodent, D., Vidal-Madjar, A., Ben Jaffel, L., Desert, J.-M.,
Moos, H.W., Sahnou, D.J., Weaver, H.A., Wolven, B.C., Ajello, J.M., Waite, J.H., Roueff, E., Abgrall,
H. Jovian auroral spectroscopy with FUSE: Analysis of self-absorption and implications for electron
precipitation. *Icarus* 171, 336–355, 2004b.

1285

Gustin, J., Cowley, S.W.H., Gérard, J.-C., Gladstone, G.R., Grodent, D., Clarke, J.T. Characteristics of
Jovian morning bright FUV aurora from Hubble Space Telescope/Space Telescope Imaging
Spectrograph imaging and spectral observations, *J. Geophys. Res.*, 111, A09220, 13 PP,
doi:10.1029/2006JA011730, 2006.

1290

Gustin, J., Gérard, J.C., Pryor, W., Feldman, P.D., Grodent, D., Holsclaw, G., Characteristics of
Saturn's polar atmosphere and auroral electrons derived from HST/STIS, FUSE and Cassini/UVIS
spectra. *Icarus* 200, 176–187, doi:10.1016/j.icarus.2008.11.013, 2009.

1295 Gustin, J., B. Bonfond, D. Grodent, and J.-C. Gérard, Conversion from HST ACS and STIS auroral
counts into brightness, precipitated power, and radiated power for H₂ giant planets, *J. Geophys. Res.*,
117, A07316, doi:10.1029/2012JA017607, 2012.

Gustin, J., Grodent, D., Gérard, J.-C., Radioti, A., Bunce, E. J., Nichols, J. D., Clarke, J. T., Mapping of
1300 the Jovian auroral electron energy with HST/STIS observations, European Planetary Science Congress
2013, EPSC Abstracts Vol.8, EPSC2013-596, 2013.

Gustin, J., Gérard, J.C., Grodent, D., Gladstone, G.R., Clarke, J.T., Pryor, W.R., Dols, V. Bonfond, B.,
Radioti, A., Lamy, L. ; Ajello, J.M. Effects of methane on giant planet's UV emissions and implications
1305 for the auroral characteristics ; *J. Mol. Spec.*, 291, 108-117, doi : 10.1016/j.jms.2013.03.010, 2013.

Gustin, J., Ray, L., Grodent, D., Clarke, J. T., Bonfond, B., Radioti, A., Nichols, J. D., Bunce, E. J.,
Mapping of the Jovian auroral electron energy with HST/STIS observations, American Geophysical
Union, Fall Meeting 2014, abstract #P23A-3978, 2014.

1310 Hess, S.L.G., P. Delamere, V. Dols, B. Bonfond and D. Swift, Power transmission and particle
acceleration along the Io flux tube. *J. Geophys. Res.*, **115** (A6), doi:10.1029/2009JA014928, 2010.

Knight, S. Parallel electric fields, *Planet. Space Sci.*, 21, 741, doi: 10.1016/0032-0633(73)90093-7,
1315 1973.

Lundin, R., Sandahl, I., Some characteristics of the parallel electric field acceleration of electrons over discrete auroral arcs as observed from two rocket flights, ESA European Sounding Rocket, Balloon and Related Res., 125-136, 1978.

1320

Mauk, B.H., J.T. Clarke, D. Grodent, J.H. Waite, C.P. Paranicas and D.J. Williams, Transient aurora on Jupiter from injections of magnetospheric electrons, *Nature*, **415**, 6875, pp 1003-1005, 2002.

Maurellis, A. N., and T. E. Cravens, Ionospheric effects of Comet Shoemaker-Levy 9 impacts with
1325 Jupiter, *Icarus*, 154, 350–371, doi:10.1006/icar.2001.6709., 2001.

Moses, J. I., Fouchet, T., Bézard, B., Gladstone, G. R., Lellouch, E., Feuchtgruber, H. Photochemistry and diffusion in Jupiter's stratosphere: Constraints from ISO observations and comparisons with other giant planets, *J. Geophys. Res.*, **110** (E8), doi: 10.1029/2005JE002411, 2005.

1330

Nichols, J. D., Cowley, S.H.W., Magnetosphere-ionosphere coupling currents in Jupiter's middle magnetosphere: effect of magnetosphere-ionosphere decoupling by field-aligned auroral voltages, *Annales Geoph.*, **23**, 799-808, doi: 10.5194/angeo-23-799-2005, 2005,

1335 Ozak, N., Schultz, D.R., Cravens, T.E., Kharchenko, V., Hui, Y.-W., Auroral X-ray emission at Jupiter: Depth effects. *J. Geophys. Res.: Space Phys.*, **115** (A11), doi: 10.1029/2010JA015635, 2010.

Ozak, N., Cravens, T.E., Schultz, D.R., Auroral ion precipitation at Jupiter: Predictions for Juno, *Geoph. Res. Let.*, **40**, 4144-4148, doi: 10.1002/grl.50812, 2013.

1340

Palmaerts, B., Radioti, A., Grodent, D., Chané, E., Bonfond, B., Transient small-scale structure in the main auroral emission at Jupiter, *J. Geophys. Res.*, **119**, 9931-9938, doi: 10.1002/2014JA020688

1345 Prangé, R., D. Rego, D. Southwood, P. Zarka, S. Miller and W. Ip, Prangé, Rapid energy dissipation and variability of the Io-Jupiter electrodynamic circuit, *Nature*, **379**, pp 323-325, doi:10.1038/379323a0, 1996.

Prangé, R., D. Rego, L. Pallier, J. Connerney, P. Zarka, J. Queinnec, Detailed study of FUV Jovian auroral features with the post-COSTAR HST faint object camera, *J. Geophys. Res.*, **103**, p. 20195-1350 20216, doi: 10.1029/98JE01128, 1998.

Radioti, A., Gérard, J.C., Grodent, D., Bonfond, B. Krupp, N., Woch, J., Discontinuity in Jupiter's main auroral oval, *J. Geophys. Res.*, **113**, (A1), doi: 10.1029/2007JA012610

1355 Radioti, A., A.T. Tomas, D. Grodent, J.-C. Gérard, J. Gustin, B. Bonfond, N. Krupp, J. Woch, J.D. Menietti, Equatorward diffuse emissions at Jupiter: Simultaneous HST and Galileo observations, *Geoph. Res. Let.*, **36**, doi:10.1029/2009GL037857, 2009.

1360 Ray, L. C., Y. Su, R. E. Ergun, P. A. Delamere, and F. Bagenal, Current-voltage relation of a centrifugally confined plasma, *J. Geophys. Res.*, **114**, A04214, doi:10.1029/2008JA013969, 2009.

Ray, L. C., Ergun, R.E., Delamere, P.A., Bagenal, F., Magnetosphere-ionosphere coupling at Jupiter: Effect of field-aligned potentials on angular momentum transport, *J. Geophys. Res.*, **115**, (A9), doi : 10.1029/2010JA015423, 2010

1365

Scudder, J. D., E. C. Sittler, Jr., H. S. Bridge, A survey of the plasma electron environment of Jupiter: A view from Voyager, *J. Geophys. Res.*, **86**, 8157, doi: 10.1029/JA086iA10p08157, 1981.

Seiff, A., and 10 coauthors, Structure of the atmosphere of Jupiter: Galileo probe measurements,
1370 *Science*, 272, 844–845, doi:10.1126/science.272.5263.844, 1996.

Smith, F. L., C. Smith, Numerical evaluation of Chapman's grazing incidence integral $Ch(X,\chi)$, *J. Geophys. Res.*, **77**, 3592-3597, doi:10.1029/JA077i019p03592, 1972.

1375 Su, Y.-J., R. E. Ergun, F. Bagenal, and P. A. Delamere, Io-related Jovian auroral arcs: Modeling parallel electric fields, *J. Geophys. Res.*, 108 (A2), 1094, doi:10.1029/2002JA009247, 2003.

Waite Jr., J. H., G. R. Gladstone, W. S. Lewis, R. Goldstein, D. J. McComas, P. Riley, R. J. Walker, P.
1380 Robertson, S. Desai, J. T. Clarke, and D. T. Young, An auroral flare at Jupiter. *Nature*, **410**, 787, 2001.

Wannawichian, S., Clarke, J. T., Nichols, J. D., Ten years of Hubble Space Telescope observations of the variation of the Jovian satellites' auroral footprint brightness, *J. Geophys. Res.*, **115** (A2), doi: 10.1029/2009JA014456, 2010.

1385

Yung, Y. L., G. R. Gladstone, K. M. Chang, J. M. Ajello, and S. K. Srivastava. H₂ fluorescence

spectrum from 1200 to 1700 Å by electron impact: Laboratory study and application to jovian aurora, *Astrophys. J.*, 254, L65–L69, doi:10.1086/183757, 1982.

Tables

Table 1: Brightness and mean energy of the different auroral regions shown in Fig. 1 and Fig. 2.

		Observation #1 (16 Jan. 2013)							Observation #2 (24 Jan. 2013)						
		ME 1	ME 2	FL 1	FL 2	DC	IFP	SSS	ME 1	ME 2	FL 1	FL 2	DC	IFP	GFP
# pixels		1801	988	865	2350	1319	198	112	1107	1381	905	1725	1314	138	95
Mean view angle (°)		69	66	72	69	59	80	60	66	67	68	69	58	76	58
Brightness															
Mean vertical brightness from images (kR) ⁽¹⁾		286	148	47	59	31	178 ⁽²⁾	194	273	180	91	75	28	145 ⁽²⁾	361
Best fit parameters ⁽³⁾	E ₀	30.5	48.1	10.5	10.6	10.0	109.0	113.1	78.5	21.7	21.4	12.2	8.3	55.7	102.7
	n	17.5	2.8	7.5	9.4	3.7	2.0	2.0	5.5	10.9	6.5	10.3	4.3	4.1	5.9
Vertical brightness from spectrum (kR) ⁽¹⁾⁽⁴⁾		219	132	48	58	36	-	91	215	166	88	73	17	-	242
Observed brightness from spectrum (kR) ⁽¹⁾		278	204	79	93	32	175	57	259	264	140	118	15	144	109
Mean energy															
Mean observed value from image (keV)		301	122	69	91	-	-	-	356	138	69	88	-	-	-
Best fit parameters ⁽³⁾	E ₀	38.6	27.4	15.7	16.1	-	-	-	61.5	38.2	20.6	10.7	-	-	-
	n	12.0	5.9	6.6	8.9	-	-	-	7.7	4.3	7.5	13.9	-	-	-
CR from spectrum		13.6	3.9	2.0	2.9	2.3	1.3	11.0	13.0	4.4	3.2	2.9	1.9	1.3	6.7
Value from spectrum (keV) ⁽⁵⁾		278	102	52	77	67	-	232	259	111	86	79	53	-	157
Value from constrained distribution (keV) ⁽⁶⁾		256	106	51	71	-	-	-	275	112	83	78	-	-	-

1395

(1) Unabsorbed brightness in 700-1700 Å derived from maps shown in Fig.1

(2) The conversion from observed to vertical brightness uses VER profiles resulting from Maxwellian distributions. Since such distribution is not consistent with the observed constraints in the case of the IFP, observed values are shown for the IFP emission.

(3) Best parameters of the generalized Maxwellian distribution for histograms based on all pixels included in each auroral region (see Fig. 3)

1400

(4) Corrected for the viewing angle and finite width of the auroral curtain. Width of 600 km is used for ME1, ME2 and DC, 1500 km for FL1 and FL2, and 150 km for the IFP, GFP and Small Scale Structure.

(5) Based on the CR method

(6) see section 4.3.3.

1405 **Table 2: Fit parameters of the electron distributions inferred from the binned pixels**

	Generalized Maxwellian					Kappa
	Mean energy (keV)	Energy flux (mW/m ²)	Constant factor C	E ₀	Parameter n	Spectral index
Main emission 1 a	256	30.7	7.69x10 ¹¹	132.4	1.87	8.04
Main emission 1 b	275	30.9	7.31x10 ¹¹	146.2	1.80	7.96
Main emission 2 a	106	13.9	9.06x10 ¹¹	71.3	0.79	3.17
Main emission 2 b	112	16.0	9.38x10 ¹¹	58.5	1.57	8.75
Flare 1 a	51	3.8	5.13x10 ¹¹	29.7	1.21	5.10
Flare1 b	83	7.6	6.38x10 ¹¹	50.8	1.09	4.48
Flare 2 a	71	4.5	4.50x10 ¹¹	51.1	0.78	3.81
Flare2 b	78	6.4	5.38x10 ¹¹	42.0	1.64	9.16

1410

Figure captions

Figure 1: Brightness map of the two STIS datasets examined in this study.

- 1415 a) Image obtained from the 16 Jan. 2013 observation. Several auroral regions have been selected by eye and are highlighted by red contours (see text for details). A grid of 15° -spaced planetocentric latitudes and longitudes is shown by the white dotted lines and a length of 5000 km is displayed by the white line on the disk.
- 1420 b) System III polar projection of the same data. The 180° longitude is indicated towards the bottom and 90° to the right. The main emission and Io footprint statistical locations are also overplotted (white solid lines).
- 1425 c) and d) respectively show the brightness map as seen from Earth and the corresponding polar projection, for the 24 Jan. 2013 observation. With the exception of region 7 (small scale structure for the Jan. 16 image and Ganymede footprint for the Jan 24. image), the other auroral regions are common to the two observations.

1430 Figure 2. Map of the precipitating electron mean energy for the two STIS observations, both in a Earth-orbit view (a and c panels) and in a polar projected view (b and d panels). All the auroral regions, labels and grids are identical to those of Figure 1. The ME1 and ME2 regions are the most energetic (~ 330 keV and ~ 130 keV respectively), followed by the poleward flare emissions (70-90 keV). The IFP shows very weak absorption while the SSS and GFP show significant hydrocarbon absorption. The images are smoothed over a 3 pixels boxcar for better legibility.

1435

Figure 3. a-n) Histogram of the vertical brightness due to the individual pixels forming the different auroral zones defined in Fig.1. Since the conversion to vertical brightness uses emission profiles resulting from Maxwellian electron distributions that are not appropriate for the IFP, the observed brightness is provided for the IFP. The distributions (black) are fitted with a Maxwellian (dashed blue), kappa (green), and generalized Maxwellian (orange) functions.

1440

o-v) Same as a-n) for the mean energy.

1445 Fig. 4. a) Energy – energy flux relationship for the main emissions (light blue and green stars) and flare emissions (black and yellow crosses) of the two STIS observations, along with data points obtained by Gustin et al. (2004) (dark blue stars and red diamonds).

b) and c) STIS points from ME1 with the classic Knight relation. Curves are shown for best fits (red) and electron source density $n=1000\text{ m}^{-3}$ (dashed) and $n=10000\text{ m}^{-3}$ (dot-dashed), with electron source temperature fixed to 2.5 keV.

1450

d) and e): same as b) and c) with the relativistic Knight relation

f) and g) STIS points for ME2 with the classic Knight relation.

Figure 5. a) For a given atmosphere, the VER must be strongly constrained to provide information on the primary electron distribution. The two parameter-free VERs (green and red lines) have a very different shape and peak altitudes and are associated with very different mean electron energies. The blue dashed line shows the VER derived from the constrained electron distribution plotted in Fig. 6a

1455

b) all VERs pictured in a) provide an equivalent fit to the spectrum ME1 a. This auroral

region is characterized by the most energetic primary electrons, as demonstrated by the huge hydrocarbons absorption when compared to the unabsorbed modeled spectrum (dashed light blue). c) same as b), with a log scale on the vertical axis.

d) optical depth due to the main hydrocarbons. For very absorbed spectra, absorption by C_2H_2 and C_2H_6 clearly influence the spectral shape, especially in the 1420-1470 Å (C_2H_6) and 1470-1530 Å (C_2H_6) bandwidths and thus cannot be neglected.

Figure 6. Energy distribution of the precipitating electrons determined from the brightness and mean energy of the pixels that define each auroral zone. In order to provide analytical formulation, these distributions have been fitted with Maxwellian (dashed blue), kappa (green) and generalized Maxwellian (orange) functions. In general, the generalized Maxwellian provides the best fit to the observed distributions (see text for details).

Figure 7 a) Several VER profiles provide a good fit to the IFP spectra. The synthetic VER 1 (orange) corresponds to a thin-Maxwellian-type profile of $\langle E \rangle \sim 20$ keV, as used for the CR method. The VER 2 (green) has a thicker FWHM, consistent with ACS observations (blue), but peaks at lower altitude than the emission observed by ACS. To reconcile the observational constraints ('thick' VER with altitude peak >550 km and CR ~ 1.2), an hydrocarbon upwelling of at least 100 km is required. The VER 3 (red) satisfies these constraints.

b) All VER profiles seen in a) provide equivalent synthetic spectra (red), slightly attenuated by hydrocarbons compared to the unabsorbed model (dashed light blue).

c) Hydrocarbon upwelling is necessary to meet all the observed constraints related to the IFP emission. Several VERs peaking at different altitudes provide equivalent spectral fits to the observed IFP spectra, depending on the strength of the homopause upward shift.

d) Relationship between the upwelling and the turbulent diffusion coefficient K_h . The very high values reached by K_h needed to meet observational constraint suggest combination with other vertical flow processes.

Figure 8. a) Synthetic VER profile leading to a good fit of the spectrum related to the Ganymede footprint seen on the 24 Jan. 2013 STIS observation.

b) Observed GFP spectrum (black line), along with the unabsorbed synthetic spectrum (dashed blue) and absorbed spectrum (red) associated to the VER profile shown in a).

Figure 9. Atmospheric model used in this study. From 3900 to 115 km, we use the auroral model from Grodent et al. (2001), and below 115 km, we used the low latitude model from Moses et al. (2005) whose density profiles have been scaled to provide a good merging with the Grodent et al. (2001) model. Panel a: temperature profile, b: H_2 density profile, c: density profiles of the main hydrocarbons

Figure 10. Inclusion of the width of the auroral curtain 'D' in the model. Rays ' R_i ' emitted at angle χ and separated by ' e ' km are summed to represent the total emergent emission. For each ' R_i ', a limited number of 'emitting points', due to different layers of thickness dz , contribute to the emission.

Figure 11. a) effect of the viewing angle on the observed color ratio, for different primary electron mean energies

b) zoomed version of a), with CR for UV emission due to secondary electrons from ion precipitation now included (black curves), showing that UV emission resulting from ion precipitation is weakly affected by hydrocarbon absorption.

c) effect of the width of the auroral curtain on the unabsorbed auroral H_2 emission, for

primary electrons of 80 keV. The y axis represents the ratio between the unabsorbed emission at angle χ to the emission at $\chi=0^\circ$. A width $D=150$ km is used to model the Io and Ganymede footprints and the small scale structure, $D=600$ km is used for the main emissions and $D=1500$ km is used for the polar cap emissions.

1510 d) effect of the auroral width on the CR for $\langle E \rangle = 80$ keV.
 e) effect of the electron mean energy on the CR. The influence of the auroral width on the CR is negligible.
 f) effect of the auroral width on the intrinsic (unabsorbed) and observed (absorbed)

1515 emergent auroral brightness for $\langle E \rangle = 80$ keV and a viewing angle of 70° . As expected, the brightness significantly increases with the auroral width.

Figure 12. Characteristics of the VER profile for different mean energies, assuming Maxwellian distributions of primary electrons. It is seen that

1520 a) The altitude of the VER decreases with increasing energy.
 b) The maximum of the VER is very energy-dependent
 c) In an altitude scale, the vertical thickness of the auroral emission decreases continuously with increasing $\langle E \rangle$.
 d) For a constant energy flux input, the total emitted brightness strongly depends on the electron

1525 mean energy. The vertical brightness in the Lyman and Werner bands is maximum at 20 keV (14.6 kR per incident mW m^{-2}) and then significantly decreases for higher energies.




Figure 1


















Figure 2















Figure 3










Figure 5




1560 Figure 6
74



1565 Figure 7



1570 Figure 8



1575 Figure 9




Figure 10

1580




Figure 11




Figure 12

1590