

Composite type-2 biomarker strategy versus a symptom-risk-based algorithm to adjust corticosteroid dose in patients with severe asthma: a multicentre, single-blind, parallel group, randomised controlled trial



Liam G Heaney, John Busby, Catherine E Hanratty, Ratko Djukanovic, Ashley Woodcock, Samantha M Walker, Timothy C Hardman, Joseph R Arron, David F Choy, Peter Bradding, Christopher E Brightling, Rekha Chaudhuri, Douglas C Cowan, Adel H Mansur, Stephen J Fowler, Robert M Niven, Peter H Howarth, James L Lordan, Andrew Menzies-Gow, Tim W Harrison, Douglas S Robinson, Cecile T J Holweg, John G Matthews, Ian D Pavord, on behalf of the investigators for the MRC Refractory Asthma Stratification Programme*



Summary

Background Asthma treatment guidelines recommend increasing corticosteroid dose to control symptoms and reduce exacerbations. This approach is potentially flawed because symptomatic asthma can occur without corticosteroid responsive type-2 (T2)-driven eosinophilic inflammation, and inappropriately high-dose corticosteroid treatment might have little therapeutic benefit with increased risk of side-effects. We compared a biomarker strategy to adjust corticosteroid dose using a composite score of T2 biomarkers (fractional exhaled nitric oxide [FENO], blood eosinophils, and serum periostin) with a standardised symptom-risk-based algorithm (control).

Methods We did a single-blind, parallel group, randomised controlled trial in adults (18–80 years of age) with severe asthma (at treatment steps 4 and 5 of the Global Initiative for Asthma) and FENO of less than 45 parts per billion at 12 specialist severe asthma centres across England, Scotland, and Northern Ireland. Patients were randomly assigned (4:1) to either the biomarker strategy group or the control group by an online electronic case-report form, in blocks of ten, stratified by asthma control and use of rescue systemic steroids in the previous year. Patients were masked to study group allocation throughout the entirety of the study. Patients attended clinic every 8 weeks, with treatment adjustment following automated treatment-group-specific algorithms: those in the biomarker strategy group received a default advisory to maintain treatment and those in the control group had their treatment adjusted according to the steps indicated by the trial algorithm. The primary outcome was the proportion of patients with corticosteroid dose reduction at week 48, in the intention-to-treat (ITT) population. Secondary outcomes were inhaled corticosteroid (ICS) dose at the end of the study; cumulative dose of ICS during the study; proportion of patients on maintenance oral corticosteroids (OCS) at study end; rate of protocol-defined severe exacerbations per patient year; time to first severe exacerbation; number of hospital admissions for asthma; changes in lung function, Asthma Control Questionnaire-7 score, Asthma Quality of Life Questionnaire score, and T2 biomarkers from baseline to week 48; and whether patients declined to progress to OCS. A secondary aim of our study was to establish the proportion of patients with severe asthma in whom T2 biomarkers remained low when corticosteroid therapy was decreased to a minimum ICS dose. This study is registered with ClinicalTrials.gov, NCT02717689 and has been completed.

Findings Patients were recruited from Jan 8, 2016, to July 12, 2018. Of 549 patients assessed, 301 patients were included in the ITT population and were randomly assigned to the biomarker strategy group (n=240) or to the control group (n=61). 28·4% of patients in the biomarker strategy group were on a lower corticosteroid dose at week 48 compared with 18·5% of patients in the control group (adjusted odds ratio [aOR] 1·71 [95% CI 0·80–3·63]; p=0·17). In the per-protocol (PP) population (n=121), a significantly greater proportion of patients were on a lower corticosteroid dose at week 48 in the biomarker strategy group (30·7% of patients) compared with the control group (5·0% of patients; aOR 11·48 [95% CI 1·35–97·83]; p=0·026). Patient choice to not follow treatment advice was the principle reason for loss to PP analysis. There was no difference in secondary outcomes between study groups and no loss of asthma control among patients in the biomarker strategy group who reduced their corticosteroid dose.

Interpretation Biomarker-based corticosteroid adjustment did not result in a greater proportion of patients reducing corticosteroid dose versus control. Understanding the reasons for patients not following treatment advice in both treatment strategies is an important area for future research. The prevalence of T2 biomarker-low severe asthma was low.

Funding This study was funded, in part, by the Medical Research Council UK.

Copyright © 2020 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Lancet Respir Med 2020

Published Online
September 8, 2020
[https://doi.org/10.1016/S2213-2600\(20\)30397-0](https://doi.org/10.1016/S2213-2600(20)30397-0)

*Members listed in the appendix p 3

Centre for Experimental Medicine, School of Medicine, Dentistry and Biomedical Sciences, Queen's University, Belfast, UK (Prof L G Heaney MD, J Busby PhD, C E Hanratty PhD); 23andMe, Sunnyvale, CA, USA (J G Matthews MD); Oxford Respiratory NIHR BRC, Nuffield Department of Medicine, University of Oxford, Oxford, UK (Prof I D Pavord FMedSci); Genentech, South San Francisco, CA, USA (JR Arron MD, CT J Holweg PhD, D F Choy BS); Department of Respiratory Sciences, Institute for Lung Health and Leicester NIHR Biomedical Research Centre, University of Leicester, Leicester, UK (Prof P Bradding DM, Prof C E Brightling PhD); NHS Greater Glasgow and Clyde Health Board, Gartnavel Hospital, and University of Glasgow, Glasgow, UK (Prof R Chaudhuri MD); NHS Greater Glasgow and Clyde, Stobhill Hospital, Glasgow, UK (D C Cowan MD); Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, University of Manchester and Manchester Academic Health Science Centre and NIHR Manchester Biomedical Research Centre, Manchester University Hospitals NHS Foundation Trust, Manchester, UK (Prof A Woodcock MD, SJ Fowler MD, R M Niven MD); Niche Science & Technology, Falstaff House, Richmond, UK

(T C Hardman PhD);
**Nottingham Respiratory NIHR
 Biomedical Research Centre,
 University of Nottingham,
 Nottingham, UK**
 (Prof T W Harrison MD); School
 of Clinical and Experimental
 Sciences, University of
 Southampton, NIHR
 Southampton Biomedical
 Research Centre,
 Southampton, UK
 (Prof R Djukanovic MD,
 Prof PH Howarth MD);
**The Newcastle upon Tyne NHS
 Foundation Trust, Newcastle
 upon Tyne, UK** (J L Lordan MD);
 University of Birmingham and
 Heartlands Hospital, University
 Hospitals Birmingham NHS
 Foundation Trust,
 Birmingham, UK
 (A H Mansur PhD); Royal
 Brompton & Harefield NHS
 Foundation Trust, London, UK
 (Prof A Menzies-Gow MG);
 University College Hospitals
 NHS Foundation Trust,
 London, UK
 (Prof D S Robinson MD); and
 Asthma UK & British Lung
 Foundation Partnership,
 London, UK (S M Walker PhD).

Correspondence to:
 Liam G Heaney, Centre for
 Experimental Medicine, School
 of Medicine, Dentistry and
 Biomedical Sciences, Queen's
 University, Belfast, UK
 l.heckney@qub.ac.uk
 See Online for appendix

Research in context

Evidence before this study

Treatment guidelines for asthma recommend a step-wise increase of corticosteroid treatment to control symptoms and reduce exacerbations. Evidence suggests minimal therapeutic benefit from corticosteroids in the absence of type 2 (T2)-driven eosinophilic inflammation, and in many symptomatic patients with asthma treated with corticosteroids, this type of inflammation is not present. Biomarker-based adjustment of corticosteroid treatment in severe asthma by means of sputum eosinophilia reduces exacerbations and allows safe down-titration of corticosteroid treatment, but is difficult to deliver in real-world settings. To our knowledge, no previous study has considered a biomarker-based strategy with the primary aim to reduce corticosteroid treatment in patients with severe asthma on high-dose treatment. Guidelines advocate corticosteroid dose reduction based on a combination of exacerbation risk and symptom control but evidence to support use of this strategy in severe asthma is scarce.

Added value of this study

We found no significant difference in the proportion of patients who had reduced corticosteroid dose at week 48

Introduction

Treatment guidelines for asthma recommend a step-wise increase in inhaled corticosteroid (ICS) treatment to control symptoms and reduce exacerbations.¹ However, as many as 50% of patients with symptomatic asthma might not have eosinophilic airways inflammation and respond poorly to corticosteroids.²⁻⁴ In this population, there is a clear potential for treatment to be increased to high intensity ICS and oral corticosteroids (OCS) without therapeutic benefit and with an increased potential for side-effects.⁵⁻⁹

Titration of corticosteroids with the goal of minimising sputum eosinophilia reduces severe exacerbations and allows safe down-titration of corticosteroid treatment in severe asthma.¹⁰⁻¹² However, sputum-guided management is difficult to deliver in a real-world setting. When used as a single biomarker of type-2 (T2) cytokine inflammation, fractional exhaled nitric oxide (FENO)-guided management in mild and moderate asthma results in reduced exacerbations with minimal effect on lung function or asthma symptoms.¹² However, whether a biomarker-based strategy can be used to primarily reduce corticosteroid exposure in patients with severe asthma on high-dose treatment is unclear. Asthma guidelines advocate corticosteroid reduction by assessment of exacerbation risk and symptom control but there is little evidence base to support this strategy in patients with severe asthma on high-dose corticosteroid treatment, particularly in patients receiving OCS.¹ Given that high doses of corticosteroid cause considerable morbidity, new algorithms that can be delivered in the

between those who were assigned to the T2 biomarker strategy compared with those assigned to standard care. A high proportion of patients chose not to follow study treatment advice but in those who did, a significant benefit was seen in patients in the biomarker-based strategy group. Importantly, patients who reduced treatment according to biomarker-directed therapy showed no evidence of clinical deterioration, despite being on lower corticosteroid doses.

Implications of all the available evidence

Factors driving patient choice not to follow advice to reduce corticosteroid treatment are an important area for future research. We suggest that before progression to high-dose corticosteroid treatment, predictive biomarkers of therapeutic response should be assessed to guide treatment decisions, because once a patient becomes established on high-dose corticosteroid treatment, biomarker-driven corticosteroid reduction can be difficult to achieve in symptomatic patients.

clinic for guiding corticosteroid treatment in patients with severe asthma are needed.

We have shown previously that a raised composite score of three T2-inflammation biomarkers (FENO, blood eosinophils, and serum periostin) is independently associated with increased risk of exacerbation in patients with severe asthma.¹³ This suggests that a low composite score could be used to identify a low-risk population in whom corticosteroid dose could be safely reduced.¹³ In this randomised, multicentre, parallel group trial in patients with severe asthma, we aimed to show whether such a scoring system could be used to guide corticosteroid treatment better than a standardised symptom-based or risk-based strategy, and would do so more effectively without increasing exacerbation rates, worsening asthma control, or precipitating a fall in lung function. An important secondary aim of our study was to establish the proportion of patients with severe asthma in whom T2 biomarkers remain low when corticosteroid therapy is decreased to a minimum ICS dose.

Methods

Study design and participants

This was a multicentre, single-blind (study participant), parallel group randomised controlled trial in patients with severe asthma. Details of the protocol have been published elsewhere.¹⁴ In brief, we compared a composite biomarker-based adjustment of corticosteroid therapy (using a composite index of blood eosinophil count, serum periostin, and FENO concentration; active intervention) with adjustments using an algorithm

based on asthma symptoms, lung function, and recent exacerbation history (control). The Medical Research Council Refractory Asthma Stratification Programme research consortium included a patient input platform recruited to provide direction with regard to patient needs and understanding; the group was recruited through Asthma UK.

The study was done in patients with severe asthma (Global Initiative for Asthma steps 4 and 5 classification of asthma severity) attending one of 12 specialist severe asthma centres across the UK (appendix pp 2–3). Patients eligible for enrolment were aged 18–80 years, had met well characterised diagnostic criteria for severe asthma, and had an FENO of less than 45 parts per billion (ppb) to enrich for a T2 biomarker-low population with greater opportunity to reduce corticosteroid treatment (appendix pp 5–6). Full eligibility criteria are included in the trial protocol (appendix pp 5–6).

The protocol was reviewed and approved by the Office for Research Ethics Northern Ireland (NI0158) and obtained local National Health Service Research and Development approval for individual sites. All patients provided written informed consent for study participation.

Randomisation and masking

Following a 2-week run-in period, patients were randomly assigned in a 4:1 ratio to one of the two treatment groups: biomarker-based adjustment strategy or control adjustment strategy. This randomisation ratio was employed to assist study recruitment to the biomarker-guided treatment strategy and to ensure maximal numbers of patients undergoing biomarker-based corticosteroid treatment adjustment (to establish the proportion of patients who remained T2 biomarker-low when corticosteroid doses were reduced). The online electronic case-report form (Dendrite Clinical Systems, Reading, UK) assigned patients following a random schedule to one of the treatment groups in blocks of ten, stratified by asthma control (Asthma Control Questionnaire [ACQ-7] ≥ 1.5) and use of rescue systemic steroids (≥ 2 courses in previous year). This was a single-blind study with patients masked to their allocated treatment group throughout the study.

Procedures

Following randomisation, patients attended the clinic every 8 weeks for review of their asthma control and treatment. Patient ACQ-7 scores, post-bronchodilator FEV₁, and FENO concentration data were entered into the electronic case-report form at all study visits; blood eosinophil count data were entered into the electronic case-report form within 24 h of the visit; and periostin values were entered automatically by a central laboratory within 3–5 days of sample collection (or when available). The electronic case-report form software processed individual patient data by means of the study algorithms (tables 1 and 2) and generated a treatment advisory

	0	1	2
FENO, ppb	<15	15–30	>30
Blood eosinophil count, n/ μ L	<150	150–300	>300
Periostin, ng/mL	<45	45–55	>55

FENO, blood eosinophil count, and serum periostin were measured at each study visit with each biomarker assigned a score of 0, 1, or 2. The composite biomarker score was calculated automatically by the eCRF software using the rounded average of the sum of all three biomarker scores. Treatment algorithms were generated automatically by the eCRF software; a composite biomarker score of 0 advised treatment reduction, a score of 1 advised maintenance of current treatment, and a score of 2 advised treatment increase. ppb=parts per billion. FENO=fractional exhaled nitric oxide. eCRF= electronic case-report form.

Table 1: Composite biomarker scoring system

	Score
ACQ-7 ≥ 1.5 and ≥ 1 change from baseline score or a severe exacerbation since last study visit (previous 8 weeks at baseline randomisation visit)	2
ACQ-7 is 1.0 to <1.5 or ACQ-7 ≥ 1.5 and <1 change from baseline score AND no severe exacerbation since last study visit (previous 8 weeks at baseline randomisation visit)	1
ACQ-7 <1.0 and no severe exacerbation since last study visit (previous 8 weeks at baseline randomisation visit)	0

ACQ-7 and recent exacerbation history were recorded at each study visit. To mirror usual clinical care, patients were not asked to withhold bronchodilator medication before study spirometry measurements. Treatment algorithms were generated automatically by the eCRF software; this was considered essential as prestudy observations in the UK Severe Asthma Registry had identified standard care and specifically corticosteroid treatment regimes differed substantially across clinical centres. A score of 0 advised treatment reduction, a score of 1 advised maintenance of current treatment and a score of 2 advised treatment increase. ACQ-7=Asthma Control Questionnaire-7. eCRF= electronic case-report form.

Table 2: Symptom-risk-based treatment adjustment

in both treatment groups (ie, recommendations for therapeutic adjustment as appropriate to decrease, maintain, or increase treatment; appendix, pp 6–8).

Patients in both groups received instructions at each clinic visit from dedicated study coordinators trained in the study procedures and working closely with study investigators; those in the biomarker strategy group received a default advisory to maintain treatment and those in the control group had their treatment adjusted according to the steps indicated by the trial algorithm. All participants were contacted once biomarker results were available, thereby ensuring patients remained masked to which treatment group they were in; patients in the control group received a default advisory to maintain their treatment, whereas treatment advice in the biomarker group was based on the result generated by the trial algorithm. A default advisory to maintain treatment was provided if, for practical reasons, any individual biomarker measure was not available. When patients reported an asthma exacerbation, it was managed according to the patient's self-management plan with no adjustment in background treatment. Planned therapy adjustments were deferred until the next scheduled study visit. Patients remained in the study irrespective of whether or not they followed their treatment advisories. Patients were seen by the clinical investigator in the clinic if there was any concern




Figure 1: Trial profile

FENO=fractional exhaled nitric oxide. PV=protocol violation. FVC=forced vital capacity.

about persistent poorly controlled asthma or treatment adjustments.

On completion of the study at week 48, all patients underwent final assessment. In addition to standard sample collection, patients were asked to identify which treatment strategy they thought they had been assigned to.

Outcomes

The primary outcome was the proportion of patients with a reduction in ICS or OCS dose from baseline to week 48. Secondary outcomes were the ICS dose at the end of the study, cumulative dose of ICS during the study, and proportion of patients on maintenance OCS at study end. Asthma outcomes were the rate of protocol-defined severe exacerbations per patient year, time to first severe exacerbation, number of hospital admissions for asthma, changes in lung function, ACQ-7 score, Asthma Quality of Life Questionnaire (AQLQ) score, and T2 biomarkers from baseline to week 48. We expected that some patients would be reluctant to progress to OCS irrespective of their biomarker or symptom score status, so this reluctance was a prespecified secondary outcome. To facilitate the analysis, detailed information was collected on whether treatment advisories were followed at each study visit with reasons for non-compliance recorded

where possible. A severe asthma exacerbation was recorded as new or as increased asthma symptoms if it led to at least one of the following: a doubling of daily OCS dose (for patients on maintenance OCS); the prescription of a course of rescue OCS for 3 or more consecutive days; administration of intravenous or intramuscular corticosteroid for asthma; or hospital visit for asthma. Patients were asked to manage exacerbations according to their written personalised self-management plans that were provided as part of their routine clinical care. We did a post-hoc analysis to explore the effects of biomarker-based management in patients with uncontrolled asthma.

Statistical analysis

It was assumed that a 20% difference between the study groups in the proportion of patients receiving a lower dose of corticosteroid treatment would be clinically meaningful. This assumption was supported by the findings of a subsequent Delphi exercise of international severe asthma experts.¹⁵ Assuming a study dropout rate of approximately 20% and a 10% reduction in the number of patients achieving a lower dose of corticosteroid in the control group, we estimated that 300 patients, randomly assigned in a 4:1 ratio, would provide the study with 80% power to show a 20% difference in corticosteroid reduction between the treatment groups.

Analysis was done according to a prespecified statistical analysis plan under the intention-to-treat (ITT) principle; two-tailed hypothesis tests were done at the 5% α -level with 95% CIs used throughout. For the primary outcome, reductions in corticosteroid doses between the start and end of the study were analysed by means of logistic regression models, adjusted for age, sex, smoking status, treatment centre, use of rescue steroids in the year before randomisation (categorised as <2 and ≥ 2 courses) and ACQ-7 score (categorised as <1.5 and ≥ 1.5) at baseline. Multiple imputation with chained equations was used, which assumed that data were missing at random with imputation models including treatment group, age, sex, smoking status, treatment centre, rescue steroids in the year before randomisation (categorised as <2 and ≥ 2 courses), ACQ-7 score (categorised as <1.5 and ≥ 1.5) at baseline, and change in corticosteroid dose between baseline and the first visit.¹⁶

All patients who attended at least one follow-up visit were included in the primary outcome analysis. Secondary outcomes were analysed by means of regression. Linear models were used for ACQ-7, AQLQ, FEV₁, log (FENO), log [blood eosinophils], periostin, ICS dose, and OCS dose. Negative binomial models were used for protocol-defined exacerbation and hospitalisation counts. Logistic models were used for the probability of refusing to start OCS, and Cox proportional hazards models were used for the time to first protocol-defined exacerbation. All models were adjusted for age, sex, smoking status, treatment centre, rescue steroid use in

	Biomarker strategy group (n=240)	Control group (n=61)
Age at inclusion, years	55.2 (13.4)	57.5 (11.9)
Sex		
Female	151 (63%)	43 (70%)
Male	89 (37%)	18 (30%)
Ethnicity*		
Caucasian	221 (92%)	58 (95%)
Non-Caucasian	19 (8%)	3 (5%)
Body-mass index, kg/m ²	31.4 (7.2)	32.5 (7.2)
Smoking status		
Never smoked	183 (76%)	41 (67%)
Ex-smoker	57 (24%)	20 (33%)
Working status		
Not working owing to asthma-related ill health	39 (16%)	13 (21%)
Not working owing to other cause	76 (32%)	26 (43%)
Working part-time owing to asthma-related ill health	12 (5%)	1 (2%)
Working part-time owing to other cause	38 (16%)	1 (2%)
Student	2 (1%)	0
Full time	72 (30%)	20 (33%)
Atopic disease	167 (70%)	40 (66%)
Hospital admissions for asthma in last year	0.0 (0.0–0.0)	0.0 (0.0–0.0)
Accident and emergency department visits in last year	0.0 (0.0–0.0)	0.0 (0.0–0.0)
General practitioner visits for asthma in last year	1.0 (0.0–3.0)	1.0 (0.0–3.0)
Rescue courses of oral corticosteroids in past year	2.0 (1.0–4.0)	2.0 (1.0–4.0)
Previous admission for asthma to intensive therapy unit	55 (22.9%)	9 (14.8%)
Number of previous admissions for asthma to intensive therapy unit	1.0 (1.0–2.0)	1.0 (1.0–2.0)
Ever been ventilated	27 (11%)	4 (7%)
Rhinitis	168 (70%)	40 (66%)
Eczema	85 (35%)	15 (25%)
Nasal polyps	58 (24%)	15 (25%)
Previous nasal surgery	57 (24%)	13 (21%)
Oesophageal reflux	137 (57%)	42 (69%)
Aspirin sensitivity	33 (14%)	14 (23%)
Depression or anxiety	73 (30%)	19 (31%)
Hypertension	70 (29%)	24 (39%)
Osteoporosis or osteopenia	57 (24%)	9 (15%)
Osteoarthritis	64 (27%)	14 (23%)
Hypercholesterolaemia	39 (16%)	14 (23%)
Diabetes	28 (12%)	6 (10%)
Cataracts	26 (11%)	7 (11%)
Obstructive sleep apnoea	15 (6%)	2 (3%)
Ischaemic heart disease	8 (3%)	4 (7%)
Peptic ulcer	8 (3%)	0

(Table 3 continues in next column)

	Biomarker strategy group (n=240)	Control group (n=61)
(Continued from previous column)		
Stroke	4 (2%)	2 (3%)
Chronic kidney disease	4 (2%)	3 (5%)
Glaucoma	4 (2%)	0
Myocardial infarction	2 (1%)	1 (2%)
FEV ₁	2.2 (0.7)	2.1 (0.7)
% predicted FEV ₁	75.3% (19.2)	76.5% (19.8)
Forced vital capacity	3.3 (0.9)	3.2 (0.9)
% predicted forced vital capacity	90.6% (16.3)	93.1% (18.9)
FEV ₁ /forced vital capacity	0.66 (0.11)	0.65 (0.13)
PEF	380.4 (126.4)	359.6 (133.2)
Sputum eosinophils†	1.5% (0.4–9.5)	1.3% (0.2–4.3)
Sputum neutrophils	64.8% (33.5–80.4)	57.8% (30.8–79.3)
Sputum lymphocytes	0.4% (0.0–1.5)	0.5% (0.0–1.5)
Sputum macrophage	22.1% (10.6–40.1)	27.1% (10.3–49.0)
FENO, ppb	21 (13–29)	19 (12–28)
Blood eosinophil count, 10 ³ cells per L	0.20 (0.11–0.32)	0.26 (0.15–0.40)
Periostin, ng/mL	52.8 (15.7)	53.5 (18.2)
Composite biomarker score		
0	55 (23%)	13 (21%)
1	135 (56%)	37 (61%)
2	48 (20%)	10 (16%)
Maintenance oral corticosteroids user	87 (36%)	24 (39%)
Oral corticosteroid dose (mg)	10 (5–10)	9 (5–10)
Inhaled oral corticosteroids dose (beclomethasone dipropionate µg equivalent)	2000 (2000–2000)	2000 (2000–2000)
LAMA user	109 (45%)	35 (57%)
Patient reported outcomes		
Asthma Control Questionnaire-7 score	2.0 (1.1)	2.0 (1.3)
Asthma Quality of Life Questionnaire total score	4.9 (1.3)	4.7 (1.6)

Data are n (%), median (IQR), or mean (SD). LAMA=long-acting muscarinic antagonist. FENO=fractional exhaled nitric oxide. PEF=peak expiratory flow rate. ppb=parts per billion. *Ethnicity as per Global Lung Initiative 2012. †Sputum data was available in 123 subjects at baseline.

Table 3: Demographics, medical history, lung function, biomarkers, corticosteroid treatment, and patient reported outcomes in the randomised population

the year before randomisation (categorised as <2 and ≥2 courses), and ACQ-7 score at baseline (categorised as <1.5 and ≥1.5). Data for outcomes measured at each study visit (ACQ-7, AQLQ, FEV₁, FENO, blood eosinophil count, periostin, ICS dose, and OCS dose) were further adjusted for the baseline measurement of the outcome. Secondary analysis was only done on data from patients who completed the study without imputation of missing data.




Figure 2: Proportion of patients who reduced corticosteroid treatment dose over 48 weeks
(A) In the intention-to-treat population. (B) In the per-protocol population. (C) In the group with uncontrolled asthma at baseline (Asthma Control Questionnaire-7 ≥ 1.5).

The prespecified per-protocol (PP) analysis excluded data from patients who did not attend a study visit or who did not follow any study treatment advisory (except in instances where patients had low cortisol or if patients were already on the lowest ICS dose allowed according to the protocol). The PP analysis adjusted for the same set of variables as the primary analyses. Exploratory post-hoc analyses were done to understand the observed difference between the results seen in the ITT population versus the PP population: analysis restricted to patients who had uncontrolled asthma at study entry (defined as an ACQ-7 ≥ 1.5) and exacerbation rate in patients not following treatment advisories compared with those following treatment advisories. We assessed the effect of the imputation on our primary outcome by repeating our analysis under a complete-case framework. Additionally, we assessed the robustness of our findings

to missing-not-at-random mechanisms by assuming a best-case scenario (assuming all withdrawals would have reduced their corticosteroid dose) and worst-case scenario (assuming all withdrawals would not have reduced their corticosteroid dose). All analyses were done with the STATA 16 software package (StataCorp, TX, USA). Conduct of the trial was monitored by an independent trial steering committee. This study is registered with ClinicalTrials.gov, NCT02717689.

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. All authors had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Results

Patients were recruited from Jan 8, 2016, to July 12, 2018 (figure 1). Of 549 patients assessed, 301 were randomly assigned to the biomarker strategy group ($n=240$) or to the control group ($n=61$). Baseline demographics, medical history, comorbidities, lung function, and corticosteroid treatment at baseline are shown in table 3.

For the primary outcome in the ITT population, 28.4% of patients in the biomarker strategy group reduced their dose of corticosteroid at 48 weeks compared with 18.5% of patients in the control group (adjusted odds ratio [aOR] 1.71 [95% CI 0.80–3.63]; $p=0.17$; figure 2A; table 4). Our estimates were similar when doing a complete-case analysis (aOR 1.70 [0.79–3.67]), and were robust to missing-not-at-random assumptions under both the best-case scenario (aOR 1.74 [0.81–3.74]) and worst-case scenario (aOR 1.69 [0.78–3.65]). There was no significant difference in secondary outcomes between study groups in the ITT analysis (table 4) and no difference in lung function, symptom scores, quality of life, or biomarkers in patients who reduced their corticosteroid dose (appendix p 9).

In the prespecified PP analysis of 121 patients, a significantly greater proportion of patients were on a lower dose of corticosteroid at week 48 in the biomarker study group (30.7% of patients) compared with the control group (5.0% of patients; aOR 11.48 [95% CI 1.35–97.83]; $p=0.026$; figure 2B; table 4). The baseline characteristics of the PP and ITT populations were similar except that in the PP population, patients in the control group were more symptomatic compared with the biomarker strategy group (mean ACQ-7 score 2.6 [SD 1.1] vs 2.0 [1.1]; $p=0.022$) and had greater impairment of AQLQ (mean total score 4.2 [SD 1.2] vs 5.0 [1.2]; $p=0.0086$; appendix p 11). The estimated number of patients needed to treat to achieve one patient on lower corticosteroid dose was four (95% CI 3–9). As with the ITT analysis, other than FENO, there was no significant difference in secondary outcomes (table 4).

and no worsening of lung function, symptom scores, quality of life, or biomarkers in patients who had a reduction in corticosteroid dose (appendix p 9).

The PP population was 40% of the ITT population as a consequence of patients not following at least one treatment advisory (n=124), withdrawing from the study

	Control group		Biomarker strategy group		Unadjusted analysis		Adjusted analysis	
	n	Effect estimate (95% CI)	n	Effect estimate (95% CI)	Odds ratio (95% CI)	p value	Odds ratio (95% CI)	p value
Intention-to-treat analysis								
Decrease in corticosteroid dose*	54	18.5% (9.3 to 31.4)	208	28.4% (22.3 to 35.0)	1.73 (0.83 to 3.61)	0.14	1.71 (0.80 to 3.63)	0.17
Inhaled corticosteroid dose (beclometasone dipropionate, µg equivalent)	54	2011 (1893 to 2129)	208	1955 (1820 to 2090)	-56 (-325 to 213)	0.68	-96 (-287 to 94)	0.32
Cumulative inhaled corticosteroid dose (beclometasone dipropionate, µg equivalent)	54	724 763 (684 245 to 765 281)	209	718 298 (676 416 to 760 180)	-6465 (-90 905 to 77 976)	0.88	-4348 (-87 909 to 79 212)	0.92
Oral corticosteroid dose, mg	54	6 (4 to 8)	208	5 (4 to 6)	-1 (-3 to 2)	0.58	-1 (-2 to 1)	0.46
Cumulative oral corticosteroid dose, mg	54	1626 (1109 to 2142)	209	1631 (1327 to 1935)	6 (-643 to 654)	0.99	55 (-581 to 691)	0.87
Oral corticosteroid use*	54	55.6% (41.4 to 69.1)	208	49.5% (42.5 to 56.5)	0.78 (0.43 to 1.43)	0.43	0.79 (0.36 to 1.74)	0.56
ACQ-7 score	54	2.2 (1.9 to 2.6)	206	2.0 (1.8 to 2.1)	-0.3 (-0.6 to 0.1)	0.16	-0.1 (-0.3 to 0.2)	0.47
AQLQ total score	52	4.7 (4.2 to 5.1)	206	5.0 (4.8 to 5.2)	0.4 (-0.1 to 0.8)	0.108	0.2 (-0.2 to 0.5)	0.36
% predicted FEV ₁	54	72.1 (66.5 to 77.7)	204	73.2 (70.6 to 75.8)	1.1 (-4.7 to 6.8)	0.72	0.4 (-2.7 to 3.6)	0.80
FENO ₁	54	19.5 (17.5 to 25.0)	205	22.0 (19.0 to 25.0)	1.06 (0.88 to 1.29)	0.53	1.02 (0.87 to 1.19)	0.85
Blood eosinophil count, 10 ³ cells per L†	51	0.31 (0.26 to 0.39)	206	0.20 (0.18 to 0.23)	0.66 (0.49 to 0.89)	0.0062	0.79 (0.61 to 1.03)	0.078
Periostin, ng/mL	51	54.2 (48.7 to 59.7)	201	51.4 (49.5 to 53.4)	-2.8 (-7.5 to 2.0)	0.25	-2.0 (-5.0 to 1.0)	0.19
Annual exacerbation rate‡	54	1.67 (1.35 to 2.06)	208	1.36 (1.21 to 1.54)	0.82 (0.58 to 1.16)	0.26	0.84 (0.61 to 1.15)	0.28
Time to first exacerbation§	0.72 (0.50 to 1.05)	0.085	0.77 (0.53 to 1.13)	0.18
Annual hospitalisation rate‡	54	0.16 (0.08 to 0.32)	208	0.10 (0.06 to 0.15)	0.62 (0.25 to 1.50)	0.29	0.77 (0.32 to 1.84)	0.56
Refused to initiate oral corticosteroid*	16	43.8% (19.8 to 70.1)	101	56.4% (46.2 to 66.3)	1.67 (0.58 to 4.82)	0.35	0.80 (0.23 to 2.76)	0.72
Per-protocol analysis								
Decrease in corticosteroid dose*	20	5.0% (0.1 to 24.9)	101	30.7% (21.9 to 40.7)	8.41 (1.08 to 65.68)	0.042	11.48 (1.35 to 97.83)	0.026
Inhaled corticosteroid dose (beclometasone dipropionate µg equivalent)	20	1990 (1814 to 2166)	101	1825 (1680 to 1970)	-165 (-497 to 166)	0.33	-210 (-480 to 61)	0.13
Cumulative inhaled corticosteroid dose (beclometasone dipropionate µg equivalent)	20	706 610 (650 247 to 762 973)	101	665 711 (620 861 to 710 561)	-40 899 (-143 437 to 61 639)	0.43	-52 599 (-162 431 to 57 234)	0.35
Oral corticosteroid dose, mg	20	7 (3-10)	101	5 (4-7)	-1 (-5 to 2)	0.39	-2 (-5 to 0)	0.088
Cumulative oral corticosteroid dose, mg	20	1721 (812 to 2629)	101	1644 (1217 to 2071)	-77 (-1099 to 945)	0.88	13 (-1070 to 1096)	0.98
Oral corticosteroid use*	20	65.0% (40.8 to 84.6)	101	53.5% (43.3 to 63.5)	0.62 (0.23 to 1.68)	0.35	0.43 (0.10 to 1.78)	0.24
ACQ-7 score	20	2.4 (1.9 to 3.0)	100	1.9 (1.7 to 2.2)	-0.5 (-1.0 to 0.0)	0.071	0.1 (-0.3 to 0.5)	0.52
AQLQ total score	18	4.2 (3.5 to 4.9)	99	5.1 (4.8 to 5.4)	0.9 (0.2 to 1.6)	0.010	0.5 (-0.1 to 1.1)	0.11
% predicted FEV ₁	20	70.2 (59.0 to 81.4)	99	72.2 (68.0 to 76.5)	2.0 (-8.4 to 12.5)	0.71	-0.9 (-6.8 to 5.0)	0.76
FENO ₁	20	13.5 (9.0 to 25.0)	100	21.0 (19.0 to 24.0)	1.50 (1.14 to 1.97)	0.0037	1.39 (1.07 to 1.80)	0.014
Blood eosinophil count, 10 ³ cells per L†	19	0.35 (0.26 to 0.47)	100	0.19 (0.16 to 0.21)	0.63 (0.39 to 1.01)	0.053	0.95 (0.62 to 1.47)	0.83
Periostin, ng/mL	19	49.1 (44.5 to 53.7)	97	50.7 (48.2 to 53.1)	1.5 (-4.3 to 7.4)	0.61	-0.8 (-4.9 to 3.3)	0.71
Annual exacerbation rate‡	20	1.29 (0.86 to 1.92)	101	1.04 (0.86 to 1.27)	0.82 (0.45 to 1.48)	0.50	0.89 (0.49 to 1.61)	0.69
Time to first exacerbation§	0.86 (0.46 to 1.60)	0.63	0.80 (0.40 to 1.61)	0.54
Annual hospitalisation rate‡	20	0.11 (0.03 to 0.43)	101	0.12 (0.06 to 0.21)	1.09 (0.23 to 5.13)	0.91	2.15 (0.38 to 12.27)	0.39
Refused to initiate oral corticosteroids*	3	0.0% (0.0 to 70.8)	..	0.0% (0.0 to 18.5)

(Table 4 continues on next page)

	Control group		Biomarker strategy group		Unadjusted analysis		Adjusted analysis	
	n	Effect estimate (95% CI)	n	Effect estimate (95% CI)	Odds ratio (95% CI)	p value	Odds ratio (95% CI)	p value
(Continued from previous page)								
Uncontrolled asthma (ACQ-7 ≥ 1.5)								
Decrease in corticosteroid dose*	35	5.7% (0.7 to 19.2)	125	26.4% (18.9 to 35.0)	5.92 (1.35 to 26.04)	0.019	5.78 (1.27 to 26.23)	0.023
Inhaled corticosteroid dose (beclomethasone dipropionate μg equivalent)	35	2137 (2004 to 2270)	125	2041 (1860 to 2223)	-96 (-443 to 251)	0.59	-280 (-488 to -72)	0.0084
Cumulative inhaled corticosteroid dose (beclomethasone dipropionate μg equivalent)	35	746 503 (701 511 to 791 494)	126	742 762 (684 624 to 800 901)	-3741 (-115 557 to 108 076)	0.95	8525 (-103 243 to 120 292)	0.88
Oral corticosteroid dose, mg	35	7 (4 to 9)	125	6 (5 to 8)	-0 (-3 to 3)	0.76	-2 (-4 to 1)	0.15
Cumulative oral corticosteroid dose, mg	35	1864 (1159 to 2570)	126	1927 (1487 to 2366)	63 (-838 to 963)	0.89	39 (-890 to 968)	0.93
Oral corticosteroid use*	35	62.9% (44.9 to 78.5)	125	55.2% (46.0 to 64.1)	0.73 (0.34 to 1.57)	0.42	0.49 (0.17 to 1.36)	0.17
ACQ-7 score	35	2.7 (2.3 to 3.1)	123	2.5 (2.3 to 2.7)	-0.2 (-0.7 to 0.2)	0.28	0.0 (-0.3 to 0.4)	0.82
AQLQ total score	34	4.0 (3.5 to 4.5)	123	4.4 (4.1 to 4.6)	0.4 (-0.2 to 0.9)	0.17	0.2 (-0.3 to 0.7)	0.42
% predicted FEV ₁	35	66.2 (59.4 to 72.9)	122	70.5 (67.1 to 73.9)	4.4 (-2.8 to 11.5)	0.24	0.8 (-2.8 to 4.3)	0.68
FeNO†	35	18.0 (12.0 to 25.0)	122	22.0 (21.0 to 26.0)	1.38 (1.10 to 1.73)	0.0057	1.24 (1.02 to 1.51)	0.032
Blood eosinophil count, 10 ⁹ cells per L†	33	0.27 (0.22 to 0.36)	123	0.18 (0.16 to 0.20)	0.64 (0.43 to 0.93)	0.019	0.81 (0.57 to 1.14)	0.23
Periostin (ng/mL)	32	50.7 (45.4 to 56.0)	120	50.0 (47.8 to 52.2)	-0.7 (-5.6 to 4.2)	0.78	-0.1 (-3.1 to 2.9)	0.96
Annual exacerbation rate‡	35	1.92 (1.50 to 2.46)	125	1.70 (1.48 to 1.95)	0.88 (0.60 to 1.29)	0.52	0.93 (0.65 to 1.33)	0.69
Time to first exacerbation§	0.76 (0.49 to 1.16)	0.20	0.83 (0.53 to 1.32)	0.44
Annual hospitalisation rate‡	35	0.21 (0.10 to 0.45)	125	0.13 (0.08 to 0.21)	0.60 (0.24 to 1.50)	0.27	0.67 (0.26 to 1.71)	0.40
Refused to initiate oral corticosteroid*	11	27.3% (6.0 to 61.0)	43	41.9% (27.0 to 57.9)	1.92 (0.45 to 8.26)	0.38	1.18 (0.20 to 6.94)	0.85

Data are mean (95% CI), effect estimate shown as mean difference unless otherwise stated. ACQ-7=Asthma Control Questionnaire 7; AQLQ=Asthma Quality of Life Questionnaire; FENO=fractional exhaled nitric oxide. *Data are proportion (95% CI), effect estimate shown as odds ratio. †Data are median (bootstrapped 95% CI), effect estimate shown as ratio of geometric means. ‡Value shown as rate (95% CI), effect estimate shown as rate ratio. §Effect estimate shown as hazard ratio.

Table 4: Effects on corticosteroid treatment dose, lung function, asthma symptoms, asthma related quality of life, and type-2 biomarkers

(n=32), or missing at least one study visit (n=18). Overall, 1041 (85%) of 1224 treatment advisories were followed in the biomarker-based group and 257 (81%) of 318 in the control group, though this was lower in both groups when a treatment change was mandated (273 [65%] of 419 in the biomarker strategy group and 77 [58%] of 133 in the control group (appendix p 13).

There was a wide variation in the number of deviations from the protocol seen at individual clinical centres (appendix p 12). The predominant reason for not following a treatment advisory was the patient's choice not to do so (appendix p 13) and adherence to treatment advice reduced as the study progressed (appendix p 13). Patients showed a reluctance to initiate regular OCS (56% in the biomarker strategy group and 44% in the control group). We also observed a reluctance to reduce corticosteroid treatment and notably, in the biomarker strategy group, these patients were more symptomatic than those who did follow a reduce advisory (ACQ-7 2.5 [SD 1.2] vs 1.8 [1.0]; $p<0.0001$). Those who were reluctant to reduce corticosteroid treatment also had lower lung function (FEV₁ 72.4% [SD 17.6] vs 82.8% [19.2]; $p=0.0002$) with worse airflow obstruction

(FEV₁/forced vital capacity [FVC] 65% [SD 12] vs 70% [12]; $p=0.0016$). There was no difference in the biomarker profile of patients who reduced treatment and those who did not.

The odds ratio of patients reducing corticosteroid dose in the biomarker strategy group remained significant when the PP definition was relaxed to include patients who were only non-adherent to the study protocol on one (n=200; aOR 4.54 [95% CI 1.30–15.87]; $p=0.018$) or two (n=237; aOR 2.64 [1.04–6.72]; $p=0.041$) study visits or treatment advisories.

In the biomarker strategy group, exacerbation rates were significantly higher in those patients who did not follow treatment advisories compared with those who did (adjusted hazard ratio [aHR] 1.64 [95% CI 1.13–2.38]; $p=0.010$; figure 3A) but this was not seen in the control group (aHR 1.07 [0.64–1.79]; $p=0.80$; figure 3B). This increased exacerbation rate was noted in patients who did not reduce treatment as well as those who did not increase treatment in the biomarker strategy group (appendix p 14). Overall, 60% of those patients who completed the study had a protocol-defined exacerbation.

Given the greater symptom burden in patients not following advice to reduce treatment in the biomarker strategy group, we did an exploratory post-hoc analysis in those patients with uncontrolled asthma at baseline (ACQ-7 ≥ 1.5). These patients were predominantly female, with obesity, not in current employment owing to asthma-related ill health, reported clinical depression or anxiety, and had more unscheduled asthma care, with twice as many on maintenance OCS. They also had significantly lower FEV₁ and FVC but no difference in FEV₁/FVC (appendix p 15). Analysis of outcomes in this subgroup indicated that a greater proportion of patients were taking a lower dose of corticosteroid treatment at week 48 in the biomarker study group compared with management based on standard care (26.4% of patients vs 5.7% of patients; aOR 5.78 [95% CI 1.27–26.23]; $p=0.023$; figure 2C; table 4). Despite this reduction in corticosteroid, symptoms, lung function, and exacerbation rates did not change (table 4). There was also no difference in lung function, symptom scores, quality of life, or biomarker profile in patients who reduced their corticosteroid treatment (appendix p 9).

The median corticosteroid dose reduction in those patients who reduced their corticosteroid treatment in the ITT, PP, and ACQ-7 ≥ 1.5 analyses was 1000 µg beclomethasone dipropionate equivalent consistently across all analyses (appendix p 9). There was no reduction in the median OCS dose in any analysis.

Of the 209 patients at the end of the study who felt they could identify which treatment strategy they had been assigned, 60 (36%) of 165 patients in the biomarker strategy group thought that they had been assigned to the control group, despite being told in the study information at the outset that they had a four out of five chance of being assigned to the biomarker strategy group. In the control group, 23 (52%) of 44 patients thought that they were receiving biomarker-directed treatment. There was no significant difference in whether they correctly identified their treatment group, suggesting that the strategy employed to mask patients to their treatment strategy had been effective.

An important secondary aim of the study was to identify how many patients could reduce their corticosteroid dose to the lowest dose of ICS and remain composite T2 biomarker-low. Of the patients recruited to the biomarker strategy group who completed the study, only nine (4%) of 209 patients successfully withdrew their corticosteroid treatment so that they achieved the lowest dose of ICS and remained biomarker-low (ie, a composite score of zero). Two other patients received serial advisories to reduce their corticosteroid dose because they were biomarker-low, but they did not successfully reduce to the lowest ICS dose. Taken together, 11 (5%) of 209 patients were composite biomarker-low.

There was no difference in mild or moderate adverse events between study groups (rate ratio [symptom:biomarker] 1.10, 95% CI 0.96–1.26) and




Figure 3: Proportion of patients who did or did not follow treatment advisories and had a severe asthma exacerbation

(A) In the biomarker strategy group. (B) In the control group.

there was no difference in severe adverse events between study groups (rate ratio [symptom:biomarker] 1.26, 95% CI 0.78–2.05). There was an increased rate of reported serious adverse events in the control group compared with the biomarker strategy group (rate ratio [symptom:biomarker] 1.64, 95% CI 1.04–2.60). There was no difference between study groups in the adjusted rate of hospitalisations for asthma (table 4).

Discussion

This study compared T2 biomarker-directed corticosteroid treatment with a standardised clinical strategy based on current symptoms and recent exacerbation history in people with severe asthma treated with high-dose ICS. We found no significant difference between these strategies in the proportion of patients able to lower the dose of their corticosteroids in the ITT study population. The aim of the study was to reduce corticosteroid dose in a population of T2-biomarker-low patients and importantly, in terms of effect on the patients, there was no evidence of worsening of asthma control or changes in T2-associated biomarkers in those patients who reduced their dose.

In contrast to the findings in the ITT population, the prespecified PP analysis showed a significantly greater proportion of patients reducing their corticosteroid treatment in the biomarker strategy group compared with the control group. The larger effect size seen in the PP analysis was principally because fewer patients reduced corticosteroid treatment in the control group, which might in part have been due to patients being more symptomatic compared with the ITT control population. The predominant reason for the low number of patients in the PP control group was that patients chose not to follow treatment advisories, including (in some situations) patients not increasing treatment after loss of asthma control. In both treatment groups, as expected, patients were reluctant to initiate regular OCS, but the reluctance to reduce ICS treatment was not expected, particularly since this intention had been communicated to patients as a core aim of the study before recruitment. We noted that biomarker-low patients in the biomarker strategy group who were more symptomatic and had lower lung function were less likely to agree to reduce their corticosteroid treatment. This finding suggests that some of these patients might have benefited from a more thorough explanation of the dissociation between their symptoms and corticosteroid dose when T2 biomarkers are low, which might have assisted them in agreeing to corticosteroid reduction.

There was a broad difference between clinical centres in the proportion of participants adhering to the protocol and specifically those following treatment advisories. The reason for this is unclear and will require further research. Treatment reductions seen in the PP analysis mirrored the assumptions in our power calculations estimating the potential benefit of a biomarker-based approach.

Given that participants who chose not to reduce corticosteroid treatment in the biomarker strategy group were more symptomatic, we did a post-hoc analysis to explore the effects of biomarker-based management in patients with uncontrolled asthma ($ACQ-7 \geq 1.5$). These findings were in line with our PP analysis, where a biomarker-directed approach to treatment resulted in a significant proportion of these symptomatic patients being able to reduce their corticosteroid treatment with no evidence of worsening in control of their asthma. As some biomarker-low patients with high symptom scores were reluctant to reduce treatment, we believe the scale of the benefit, in terms of the proportion who could reduce treatment, might have been underestimated in this analysis. As there were fewer opportunities for symptom-based treatment reduction in this sub-population, they serve as a useful treatment-stable comparator group, showing little difference in clinical outcomes in the biomarker strategy group who reduced their treatment significantly compared with the control group who did not. The uncontrolled asthma analysis is particularly relevant since it describes

a common clinical problem: a highly symptomatic patient potentially overtreated with corticosteroids. By means of an asthma guideline-based approach to treatment, this group could often see their corticosteroid dose increased owing to high symptom burden, but this analysis suggests that a biomarker-based strategy in this group might successfully and substantially reduce corticosteroid treatment with no deterioration in asthma control or lung function or increased exacerbation of T2 inflammation based on biomarker profile.

This study had a high retention of patients, including those who did not follow treatment advisories, allowing the opportunity to compare the risk of severe exacerbation in participants who did and did not follow treatment advice in either study group. However, given the pragmatic nature of the study, we did not track patient adherence to treatment advisories in real time (ie, treatment adjustment was reviewed at next study visit) and it was not possible to monitor differences in the proportion of patients not following advisories at the different sites. Increased risk of exacerbation in patients with severe asthma who were biomarker-high was not unexpected but the increased risk observed in biomarker-low patients who did not reduce their corticosteroid dose was not expected. The use of both OCS and ICS has been associated with increased risk of infections and azithromycin has been shown to reduce severe exacerbations in both eosinophilic and non-eosinophilic patients.^{6,7,17} The NOVEL-START study showed a tendency to increased severe exacerbations in patients with mild asthma and low blood eosinophil counts (<150 cells per μ L) initiating regular ICS treatment compared with treatment with as-needed salbutamol.¹⁸ Taken together, these data suggest that ICS in doses that suppress substantially T2 biomarkers might be harmful in some patients. If so, this makes the ability to carefully and safely titrate corticosteroid dose to T2-associated biomarker activity all the more important. However, given the difficulty that some symptomatic patients had in this study to reduce treatment when established on high-dose treatment, it might be better to use biomarkers to identify the right patient before increasing corticosteroid treatment.

The median daily ICS reduction in patients who reduced treatment was consistent across all analyses at a beclomethasone dipropionate equivalent dose of 1000 μ g. Our expert Delphi consensus advised that at least a 250–500 μ g reduction in daily ICS treatment would be regarded as a clinically meaningful dose reduction in a severe asthma population on high-dose ICS, suggesting that severe asthma specialists will recognise this as a substantial treatment benefit.¹⁵

The current study also aimed to establish the prevalence of T2 biomarker-low asthma in our severe asthma population when the corticosteroid dose was reduced to a minimum, because it is recognised that exposure to corticosteroids downregulates T2 biomarkers and overtreatment might potentially overestimate the

prevalence of the T2 low phenotype.¹⁹ At study entry, 23% of participants were composite biomarker-low, which was in line with our analysis of the placebo groups of clinical trials with lebrikizumab and omalizumab (appendix p 6).¹³ However, only 11 (5%) of 209 patients managed either to down-titrate successfully to the lowest dose of ICS and remained biomarker-low or remained serially biomarker-low and did not reduce their corticosteroid treatment. These data suggest that ICS can be reduced cautiously to relatively low maintenance levels in T2 biomarker-low patients, and this would seem important before assigning any phenotypic label. It further implies that T2 biomarker-low severe asthma, as defined in this study, is uncommon. We propose that detailed clinical assessment including corticosteroid withdrawal is important before investigational research studies and testing of clinical interventions in this T2 biomarker-low population.²⁰

In summary, this study did not meet its primary outcome of showing a greater proportion of patients on a lower corticosteroid dose by means of a biomarker-based strategy in severe asthma compared with corticosteroid adjustment based on symptoms and previous exacerbation history in the ITT population. We noted a large number of patients who did not follow treatment advisories to reduce corticosteroid treatment in both treatment groups; understanding the reasons for this finding is an important area for future research. In the study population who followed treatment advice per protocol or who had uncontrolled asthma at study entry, meaningful corticosteroid reduction was achieved. A biomarker-adjusted corticosteroid strategy seems particularly beneficial in those patients where symptoms and T2 biomarker profile are discordant and, as seen in up to 50% of patients with severe asthma;⁸ we suggest that before progression to high-dose inhaled and systemic corticosteroid treatment, predictive biomarkers of therapeutic response should be assessed. Once established on this dose, biomarker-driven dose reduction can be particularly challenging in patients with high levels of asthma symptoms.

Contributors

LGH, JB, CEH, RD, AW, SMW, TCH, JRA, DFC, PB, CEB, RC, DCC, AHM, SJF, RMN, PHH, JLL, AM-G, TWH, DSR, CTJH, JGM, and IDP developed the initial research concept, grant application, addressed feedback, reworked the full grant and ethical submission, contributed to protocol development, study procedures, and delivery of the study as well as manuscript preparation. All authors read and approved the final manuscript.

Declaration of interests

LGH has received grant funding and has taken part in advisory boards and given lectures at meetings supported by Hoffmann-La Roche–Genentech and Aerocrine. He has taken part in asthma clinical trials sponsored by Hoffmann-La Roche–Genentech, for which his institution was remunerated. LGH is Academic Lead for the Medical Research Council Stratified Medicine UK Consortium in Severe Asthma, which involves industrial partnerships with a number of pharmaceutical companies, including Hoffmann-La Roche and Aerocrine. RD reports receiving fees for lectures at symposia organised by Novartis, AstraZeneca, and TEVA, consultation for TEVA and Novartis as a member of advisory boards, and participation in a scientific discussion

about asthma organised by GSG. He is a co-founder and consultant, and has shares in Synairgen, a University of Southampton spin-out company. RD has given lectures at symposia organised by pharma companies and has consulted companies as a member of advisory boards and received non-financial support from GlaxoSmithKline, Chiesi, Novartis, and Napp Pharmaceuticals. AW reports personal fees from GlaxoSmithKline, personal fees from Novartis, personal fees from Chiesi, other from Reacta Biotech, and other services from Axalabion and the Medicines Evaluation Unit. JRA is an employee of Genentech, holds stock and stock options in the Roche Group, and is a named inventor on patents pending relating to diagnosis and treatment of asthma. DFC is an employee and holds stocks and options in Genentech and is a member of the Roche Group. He is an inventor of planned or filed patents related to the diagnosis and treatment of respiratory diseases. PB has received grant funding from Genentech, acts as a consultant for Roche and GlaxoSmithKline, and has received speaker fees from AstraZeneca and support to attend conferences from GlaxoSmithKline. CEB has received grants or consultancy fees via his institution from AstraZeneca and Roche–Genentech. CEB reports grants and personal fees from GlaxoSmithKline, AstraZeneca, Sanofi, Novartis, Chiesi, Genentech, Gossamer, Mologic, and 4DPharma for services outside of the submitted work. RC has been paid to attend Advisory Board meetings by AstraZeneca and received educational grants for research from Aerocrine. AM-G has attended advisory boards and/or received lecture fees from AstraZeneca and Hoffmann-La Roche, and has participated in clinical studies for which his institution has been reimbursed by Hoffmann-La Roche. TWH reports grants, personal fees, and non-financial support from AstraZeneca, grants and personal fees from GlaxoSmithKline, personal fees from Vectura, Synairgen, and Chiesi, outside of the submitted work. SJF reports personal fees from AstraZeneca, Chiesi, GlaxoSmithKline, Novartis, and Teva and grants and personal fees from Boehringer Ingelheim, outside of the submitted work. RMN has received funds for attending advisory boards, speaking on a fee-paid basis, and received travel support for attending internal education conferences from AstraZeneca. PHH has participated on advisory boards for Roche. He has received sponsorship to attend international scientific meetings from AstraZeneca and is an employee of GlaxoSmithKline. AHM reports personal fees, non-financial support, and other from GlaxoSmithKline, AstraZeneca, Novartis, Sanofi, and Teva, and other for services outside of the submitted work. CTJH is an employee of Genentech and a member of the Roche Group. IDP has received speaker's honoraria for speaking at sponsored meetings from AstraZeneca and Aerocrine, and payment for organising an educational event from AstraZeneca. He has received honoraria for attending advisory panels with Genentech and AstraZeneca. The remaining authors declare that they have no competing interests.

Data sharing

Following publication of the primary and secondary analyses, individual de-identified patient data, including the data dictionary, will be made available via our data sharing portal eTRIKS indefinitely delivered via Imperial College. This will allow for maximum use of the data to improve patient care and advance medical knowledge. The trial protocol and statistical analysis plan will also be made available.

Acknowledgments

This study was funded jointly by the Medical Research Council (MRC) UK (MR/M016579/1) and industrial partners within the MRC Refractory Asthma Stratification Programme consortium (appendix pp 2–3). This study was part of the MRC UK Refractory Asthma Stratification Programme and programme support was obtained from Hoffmann-La Roche–Genentech (periostin assay and sample biobanking) and Circassia (FENO measurements—reduced pricing for machines and test kits) for in-kind support within that Consortium. We would like to thank the members of the Trial Steering Committee for all their support and assistance with study delivery; Martyn Partridge (Chair), Mike Morgan, Anne Millar, Mark Stafford-Watson (patient representative—during his tenure on the Trial Steering Committee, Mark sadly passed away and we gratefully acknowledge his significant contribution to this programme and other patient involvement in research projects) and Gabriella Cooper. We are grateful to Niche Science & Technology for

For eTRIKS see <https://www.eatriks.org/>

assistance with study delivery and to all the patients who volunteered for the study and clinical and research teams at all the participating clinical and academic centres. We thank Sofia Mosesova, Chris Patterson, and Nicola Gallagher for statistical advice during study set-up and analysis. We also thank Amgen (Thousand Oaks, California, USA), Astra Zeneca (London, UK), Janssen Research & Development LLC (London, UK), GSK (UK), Boehringer Ingelheim (Germany), and Vitalograph (Ennis, Ireland) for supporting the RASP-UK Consortium.

References

- Global Initiative for Asthma 2019. <https://ginasthma.org/wp-content/uploads/2019/06/GINA-2019-main-report-June-2019-wms.pdf> (accessed Dec 12, 2019).
- Pavord ID, Brightling CE, Woltmann G, Wardlaw AJ. Non-eosinophilic corticosteroid unresponsive asthma. *Lancet* 1999; **353**: 2213–14.
- Woodruff PG, Modrek B, Choy DF, et al. T-helper type 2-driven inflammation defines major subphenotypes of asthma. *Am J Respir Crit Care Med* 2009; **180**: 388–95.
- McGrath KW, Icitovic N, Boushey HA, et al. A large subgroup of mild-to-moderate asthma is persistently noneosinophilic. *Am J Respir Crit Care Med* 2012; **185**: 612–19.
- Sweeney J, Patterson CC, Menzies-Gow A, et al. Comorbidity in severe asthma requiring systemic corticosteroid therapy: cross-sectional data from the Optimum Patient Care Research Database and the British Thoracic Difficult Asthma Registry. *Thorax* 2016; **71**: 339–46.
- Bleecker ER, Menzies-Gow AN, Price DB, et al. Systematic literature review of systemic corticosteroid use for asthma management. *Am J Respir Crit Care Med* 2019; **201**: 276–93.
- Heffler E, Madeira LNG, Ferrando M, et al. Inhaled corticosteroids safety and adverse effects in patients with asthma. *J Allergy Clin Immunol Pract* 2018; **6**: 776–81.
- Haldar P, Pavord ID, Shaw DE, et al. Cluster analysis and clinical asthma phenotypes. *Am J Respir Crit Care Med* 2008; **178**: 218–24.
- Pavord ID, Beasley R, Agusti A, et al. After asthma: redefining airways diseases. *Lancet* 2018; **391**: 350–400.
- Green RH, Brightling CE, McKenna S, et al. Asthma exacerbations and sputum eosinophil counts: a randomised controlled trial. *Lancet* 2002; **360**: 1715–21.
- Jayaram L, Pizzichini MM, Cook RJ, et al. Determining asthma treatment by monitoring sputum cell counts: effect on exacerbations. *Eur Respir J* 2006; **27**: 483–94.
- Petsky HL, Cates CJ, Lasserson TJ, et al. A systematic review and meta-analysis: tailoring asthma treatment on eosinophilic markers (exhaled nitric oxide or sputum eosinophils). *Thorax* 2012; **67**: 199–208.
- Heaney LG, Djukanovic R, Woodcock A, et al. Research in progress: Medical Research Council United Kingdom Refractory Asthma Stratification Programme (RASP-UK). *Thorax* 2016; **71**: 187–89.
- Hanratty CE, Matthews JG, Arron JR, et al. A randomised pragmatic trial of corticosteroid optimization in severe asthma using a composite biomarker algorithm to adjust corticosteroid dose versus standard care: study protocol for a randomised trial. *Trials* 2018; **19**: 5.
- Gallagher NE, Hanratty CE, Humbert M, et al. Biomarker-based corticosteroid adjustment in severe asthma: a modified Delphi consensus. *ERJ Open Res* 2018; **4**: 00081–02018.
- White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* 2011; **30**: 377–99.
- Gibson PG, Yang IA, Upham JW, et al. Effect of azithromycin on asthma exacerbations and quality of life in adults with persistent uncontrolled asthma (AMAZES): a randomised, double-blind, placebo-controlled trial. *Lancet* 2017; **390**: 659–68.
- Pavord ID, Holliday M, Reddel HK, et al. Predictive value of blood eosinophils and exhaled nitric oxide in adults with mild asthma: a prespecified subgroup analysis of an open-label, parallel-group, randomised controlled trial. *Lancet Respir Med* 2020; **8**: 671–80.
- Busby J, Holweg CTJ, Chai A, et al. Change in type-2 biomarkers and related cytokines with prednisolone in uncontrolled severe oral corticosteroid dependent asthmatics: an interventional open-label study. *Thorax* 2019; **74**: 806–09.
- McDowell PJ, Heaney LG. Different endotypes and phenotypes drive the heterogeneity in severe asthma. *Allergy* 2019; published online July 2. <https://doi.org/10.1111/all.13966>.