

Five papers extracted from the reviews identified parental deployment as having a negative effect on school attainment.

Nine studies extracted from the review papers found a positive correlation between having a deployed parent and a greater chance of experiencing depressive symptoms and feelings of anxiety.

Strong correlations of increased prevalence of both externalising & internalising behaviours were conclusively found in 7 of the reviews.

Increased resilience was detailed in only one study featured in multiple reviews.

Just one study featured across the reviews reported on physiological measures - adolescents with deployed parents had higher blood pressures and significantly higher heart rates and stress scores than civilian children.

Conclusion. More research obtaining the viewpoint of the child directly and observation of such children is required to properly understand the effects on children with a deployed parent, without the interference of parent or teacher reporting bias. Additionally, with only one study reporting on increased offspring resilience there has been limited exploration of potential positive correlates, so further research regarding these is important.

Landscape of childhood and adolescent depression in Pakistan: experience from a tertiary care hospital in Karachi, Pakistan

Maria Khan*, Mohummad Hassan Raza Raja, Fatima Gauhar and Tania Nadeem

The Aga Khan University Hospital

*Corresponding author.

doi: 10.1192/bjo.2021.703

Aims. Depression is highly prevalent in children and adolescents in Pakistan, yet, factors affecting depression have not been widely studied. This study aims to assess the demographic and clinical characteristics of depression in children and adolescents and identify associations between parental marital status and confounding factors for depression.

Method. A descriptive retrospective study was undertaken at the Aga Khan University Hospital in Karachi, Pakistan. Patient records of children and adolescents (aged under 18 years), presenting to the psychiatry clinic with depression from 2015-2019 were reviewed. The diagnosis of clinical depression was made based on clinical assessment according to international guidelines. Patients whose medical records had missing information were excluded. Data were analysed using IBM SPSS Statistics for Windows, version 23.0 (IBM Corp., Armonk, N.Y., USA). Continuous data are presented as mean \pm standard deviation, whereas categorical data are presented as percentages (%). Pearson Chi-square test of association has been used to assess the association between parental factors and confounding factors. In instances where Pearson's Chi-square test could not be applied, Fisher's exact test is used instead. Associations at $p < 0.05$ (95% confidence limit) are considered statistically significant.

Result. A total of 133 participants were included, of which 78 (58.6%) were female, and 55 (41.4%) were male, with a mean age of 15.5 \pm 2.4 (Range: Ages 4-18). The population had a 50.4% prevalence of suicidal ideation, 21.1% of self-harm, 15% of substance abuse and 14.3% of suicide attempts. Academic stress (54.9%), inter-parental conflict (30.1%) and child abuse (29.3%) were the most common confounding factors reported. Other confounding factors include a family history of depression (20.3%), experience of bullying

(16.5%) witnessing domestic violence (16.5%), substance abuse (15.0%) and experiencing sexual abuse (6.0%). There is a statistically significant association between children having parents with non-intact marriages and experiencing sexual abuse ($p < 0.001$, Odds Ratio (OR) = 21.48), having a family history of depression ($p < 0.001$, OR = 7.04), child abuse (OR = 3.78). Children of non-traditional (not living with both parents) families were more likely to witness domestic violence ($p < 0.001$, OR = 4.28), have a family history of depression ($p < 0.001$, OR = 3.44), abuse substances (OR = 3.20) and experience child abuse (OR = 2.48).

Conclusion. This study identifies factors that may put children at an increased risk of developing depression and performing high-risk behaviours. The findings can help develop better screening programs and counselling for children and adolescents, allowing prevention and ensuring early diagnosis and care.

Survey of remote consultations in psychiatry during the SARS-CoV-2 outbreak

Nusra Khodabux*, Satheesh Gangadharan, Samuel Tromans and Avinash Hiremath

Leicestershire partnership NHS Trust

*Corresponding author.

doi: 10.1192/bjo.2021.704

Aims. To compare the usage of remote consultations before and after the first wave of the SARS-CoV-2 outbreak and explore mental health workers' views on the usage of telemedicine.

Method. An online questionnaire survey was developed, and disseminated to mental healthcare professionals via e-mail and social media. Quantitative data were analysed using descriptive statistics and qualitative data were analysed using Braun and Clarke's six step procedure for thematic analysis.

Result. There were 40 responses from mental healthcare professionals of varying grades from different sub-specialties, predominantly from the UK. Compared to before the SARS-CoV-2 outbreak, there was an increase in usage of telephone (9(22.5% to 29(72.5%)) and video consultations (4(10%) to 17(42.5%)). Respondents reported an increase in virtual MDTs (35(87.5%) during the pandemic, 9(22.5%) pre-pandemic).

Based on a 5-point Likert scale, the mean technical quality of telephone consultations was 3.56/5 (Range 2-5), with 75% rating telephone consultations as not being as good as face-to-face consultations. The mean technical quality of video consultations was 3.58/5 (Range 2-5), with 63% rating video consultations as not being as good as face-to-face consultations. 25 (62.5%) respondents felt comfortable using telephone consultations during the pandemic, 20(50%) felt comfortable using video consultations. Recurring themes identified from the qualitative data regarding reasons for the technical quality ratings were: connection issues, poor infrastructure and security concerns.

Nine (23%) respondents felt that using video conferencing consultations had a detrimental impact on the mental health of patients while 14(35%) felt that telephone consultations had a detrimental impact on patients' mental health. Recurring themes for health practitioners' perceived effect of the use of telemedicine on patients' mental health were the loss of personal touch and reduced patient engagement.

Conclusion. There was a substantial increase in usage of remote consultations during the first wave of the SARS-CoV-2 pandemic among mental healthcare professionals. The results reported in the present study suggest there are numerous barriers to the use of telemedicine in psychiatry, which require future exploration, ideally through interview or ethnographic studies.

References

Braun, V, Clarke V. Using thematic analysis in psychology. *Qualitative research psychology* 2006; 3(2): 77-101

Generalized anxiety disorder among diabetic patients visiting gharyan-polyclinic in Libya during COVID-19 pandemic

Ahmed Khrwat^{1*} and Abduraof Saadawi²

¹faculty of medicine, University of Gharyan and ²Gharyan polyclinic

*Corresponding author.

doi: 10.1192/bjo.2021.705

Aims. To estimate the prevalence of Generalized anxiety disorder (GAD) in adult patients with diabetes mellitus (T1DM or T2DM) during COVID-19 pandemic.

Method. Random sample of 115 Adult Libyan patients (≥18 years) were drawn from 1200 Medical records of diabetic patients previously diagnosed in a primary care clinic (Gharyan polyclinic, South of Tripoli, West of Libya). Patients were recruited and diagnostically interviewed through outpatient visits and through Phone calls. Anxiety was assessed using Generalized Anxiety Disorder 7-item instrument (GAD-7), personal information, Co morbidities and History of COVID-19 infection within period of 3 weeks.

Result. The statistical analysis done by SPSS version 23, using ANOVA test. The GAD-7 scores ranged from 0 to 19 for the diabetic patient, 82 patients scores ranged from 0 to 4 with varying degrees of non-significant to subsyndromal symptoms of Generalized anxiety disorder, 24 patient with Mild GAD, 7 patients with moderate GAD and 2 patients with severe GAD. (P value = 0.000)

Conclusion. GAD is present in 28% of the patients who participated in the study. Additional epidemiological studies are needed to determine the prevalence of anxiety in the broader population of persons with diabetes.

Impact of online group studying for the MRCPsych A exam amongst international doctors logging-in from 7 countries

Praveen Kumar^{1*}, Sara Mohsen¹, Oksana Zinchenko¹, Philip Verde¹, Kathleen Breslin¹, Alice Judge¹ and Sridevi Shanmugam²

¹NHS Highland and ²Essex Partnership University NHS Foundation Trust

*Corresponding author.

doi: 10.1192/bjo.2021.706

Aims. Recently, global-remote group studying has been made possible via digital video conferencing platforms. In preparation for the December 2020 MRCPsych part A exam, a study group was formed comprising 30 International Medical Graduates (IMG) logging-in from different countries via 3 hour Zoom-study sessions hosted daily from 28th September until 12th December 2020 (1800-2100 GMT time). This study demonstrates the impact of online group study in preparation for the MRCPsych A exam for s via data collected through questionnaires.

Method. The data of the study were collected through the questionnaires given to the group study members containing a total of 17 questions, 5 of which were open-ended.

The participants totalled 30 International Doctors who responded to an advertisement to form an online study group

on Facebook. They logged-in for the sessions from seven different countries: Malaysia, India, Bangladesh, Ireland, Nigeria, Saudi Arabia, and the United Kingdom. The participants represented different working grades including experiences in psychiatry ranging from 0 to 5 years.

Data were analysed using percentage. The answers given to the open-ended questions were each examined using descriptive interpretation methods.

Result. Thematic analysis demonstrated that online group study made learning faster and easier. 96.6% support using online study sessions for future exams citing that they fostered cooperation, respect for diverse opinions and motivation for regular studying. 93.1% and partly 6.9% found the experience enjoyable and enabled the cultivation of different ideas. Indeed, 89.7% relied on it as a big part of their preparation with 26 saying it contributed to their passing of the exam success.

Almost three quarter of participants in the group also forged friendships and a sense of trust. It also became a platform for expressing opinions comfortably and developing communication and interpersonal skills.

Different working hours and time zones represented a challenge with most linking in at odd hours. Cultural differences were ultimately accepted including aspects of delivery of information which made a few participants appear abrupt.

Conclusion. With the ease in which social media connects us on a global scale, online study groups connecting IMGs from various backgrounds and diverse cultures not only makes exam preparations stimulating and easier to pass but also fosters interpersonal skills and connections that would be an asset in the long run.

Glucose transporters in Alzheimer's disease

Natalia Kyrtata^{1*}, Ben Dickie², Hedley Emsley³ and Laura Parkes²

¹The University Hospitals of Morecambe Bay Trust, The University of Manchester; ²The University of Manchester and ³Lancaster University

*Corresponding author.

doi: 10.1192/bjo.2021.707

Background. Physiological brain function depends on tight glucose regulation, including transport and phosphorylation, the first step in its metabolism. Impaired glucose regulation is increasingly implicated in the pathophysiology of Alzheimer's disease (AD). Glucose hypometabolism in AD may be at least partly due to impaired glucose transport at the blood-brain barrier (BBB). Glucose transporters (GLUTs) are an integral component of the BBB. There is evidence of a significant reduction in vascular and non-vascular forms of GLUT1 and GLUT3 in AD brains compared to age-matched controls. Glucose transport, as well as phosphorylation, appears to be a rate limiting step for glucose metabolism in the brain. We have reviewed the literature on glucose transport abnormalities in AD and the effect such abnormalities have on the brain.

Method. Published literature between 1st January 1946 and 1st November 2019 was identified using EMBASE and MEDLINE databases and titles and abstracts were scanned. Human studies (autopsy and imaging) and data from animal models were included while reviews, letters and cellular or molecular studies were excluded from the search.

Result. Autopsy studies in AD patients show significant reductions in GLUT3 in areas of the brain closely associated with AD pathology. Patients with AD and diabetes showed greater reductions of GLUT1 and GLUT3. A longitudinal study showed